


MEDICAL ELIGIBILITY CRITERIA FOR CONTRACEPTIVE USE



Third edition, 2004



RH Reproductive Health and Research
World Health Organization, Geneva

WHO Library Cataloguing-in-Publication Data

World Health Organization.

Medical eligibility criteria for contraceptive use. -- 3rd ed.

1.Contraception - methods 2.Family planning - methods 3.Eligibility determination - standards 4.Quality assurance, Health care 5.Health services accessibility 6.Guidelines I.Title.

ISBN 92 4 156266 8

(NLM classification: WP 630)

© World Health Organization 2004

All rights reserved. Publications of the World Health Organization can be obtained from Marketing and Dissemination, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 2476; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to Publications, at the above address (fax: +41 22 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

Printed in

Table of contents

Acknowledgements

Executive summary and Overview

Tables

Low-dose combined oral contraceptives (COCs)

Combined injectable contraceptives, patch and ring (CICs/P/R)

Progestogen-only contraceptives (POCs)

Emergency contraceptive pills (ECPs)

Intrauterine devices (IUDs)

Copper IUD for emergency contraception (E-IUD)

Barrier methods (BARR)

Fertility awareness-based methods (FAB)

Lactational amenorrhoea method (LAM)

Coitus interruptus (CI)

Surgical sterilization procedures (STER)

Summary tables (SUMM)

Annex

Annex 1. COCs and antiretroviral therapies

Annex 2. List of participants

Acknowledgements

This document is the result of collaboration between the World Health Organization's Department of Reproductive Health and Research and a large number of international agencies and organizations active in the field of family planning policies and programmes. Funding and other support for this project was provided by the Government of the United States of America (through the US Agency for International Development, the Centers for Disease Control and Prevention, and the National Institute of Child Health and Human Development), the International Planned Parenthood Federation and the United Nations Population Fund. This support is gratefully acknowledged.

Representatives of 10 agencies and organizations, along with 19 other individuals, served as experts at the meeting that achieved consensus on these recommendations for contraceptive use. We would like to express our deep appreciation to all of them for contributing their time and expertise towards the consensus-building process.

The evidence on which the decisions in this document were based was in large part obtained from systematic reviews of the literature conducted and summarized by Dr KM Curtis, Dr ME Gaffield, Ms AP Mohllajee, Dr K Nanda, and Dr JS Smith, who also provided substantial support to Secretariat. Dr H Peterson was overall coordinator of the project for the WHO Secretariat, which included Ms K Church, Ms K Curran, Ms S Johnson and Ms G Lamptey. Ms C Hamill, who was also a part of Secretariat, contributed substantially to the meeting and was responsible for the design and layout of the publication. Ms M NíMhearáin was responsible for the cover design. We would like to express our deep appreciation to these individuals as well as to Drs L Edouard, C Huezco and J Shelton for their strong support of this endeavour.

We are grateful to the following individuals who served as peer reviewers for the Continuous Identification of Research Evidence (CIRE) system: Dr T Chipato, Dr P Corfman, Dr M Cravioto, Dr V Cullins, Dr J Diaz, Dr S Diaz, Dr A Glasier, Dr J Guillebaud, Dr M Gulmezoglu, Dr K Hagenfeldt, Dr P Hannaford, Dr R Hatcher, Dr C Huezco, Dr V Jennings, Dr P Lynam, Dr P Marchbanks, Dr O Meirik, Dr S Mittal, Dr K Nanda, Dr E Olorin, Dr A Pollack, Dr H Rees, Dr R Rivera, Dr D Skegg, Dr C Smith, Dr B Sood, and Dr E Weisberg.

Partial funding for the printing and dissemination of this publication was provided by the United Nations Population Fund and is gratefully acknowledged.

For any further information on this publication, please contact the Department of Reproductive Health and Research, World Health Organization, 1211 Geneva 27, Switzerland. Direct fax: + 41 22 791 4189; e-mail: reproductivehealth@who.int

Further copies may be obtained from: Documentation Centre, Department of Reproductive Health and Research, World Health Organization, 1211 Geneva 27, Switzerland. Direct fax: + 41 22 791 4189; telephone: + 41 22 791 4447; e-mail: rhrpublications@who.int. The document is also available through WHO's reproductive health web site at www.who.int/reproductive-health. Any updates of the information contained in this document will appear in the first instance on this site.

Executive summary	1
Overview	2
Goal	3
Background	3
Reproductive and sexual health care	3
Issues of service quality and access that affect method use	4
Effectiveness of method	5
Conditions that expose a woman to increased risk as a result of unintended pregnancy	8
Return to fertility	8
STIs and contraception: Dual protection	9
Method of work	9
How to use this document	11
Using the tables	11
Classification of categories.....	12
Using the categories in practice	13
Programmatic implications	13
Clients with special needs	14
Adolescents	14
Summary of changes from the second edition	15

Executive summary

This document is one important step in a process for improving access to quality of care in family planning by reviewing the medical eligibility criteria for selecting methods of contraception. It updates the second edition of *Improving access to quality care in family planning: medical eligibility criteria for contraceptive use*, published in 2000, and summarizes the main recommendations of an expert Working Group meeting held at the World Health Organization, Geneva, 21–24 October 2003. (Please see Annex 2 for the list of participants.) The Working Group brought together 36 participants from 18 countries, including representatives of many agencies and organizations. The document provides recommendations for appropriate medical eligibility criteria based on the latest clinical and epidemiological data and is intended to be used by policy-makers, family planning programme managers and the scientific community. It aims to provide guidance to national family planning/reproductive health programmes in the preparation of guidelines for service delivery of contraceptives. It should not be seen or used as the actual guidelines but rather as a reference.


The document covers the following family planning methods: low-dose combined oral contraceptives (COCs), combined injectable contraceptives (CICs), combined patch (P), combined vaginal ring (R), progestogen-only pills (POPs), depot medroxyprogesterone acetate (DMPA), norethisterone enantate (NET-EN), levonorgestrel (LNG) and etonogestrel (ETG) implants, emergency contraceptive pills (ECPs), copper intrauterine devices (Cu-IUDs), levonorgestrel-releasing IUDs (LNG-IUDs), copper-IUD for emergency contraception (E-IUD), barrier methods (BARR), fertility awareness-based methods (FAB), lactational amenorrhoea method (LAM), coitus interruptus (CI), and female and male sterilization (STER).

WHO will update and add to the recommendations in this document at appropriate intervals through expert Working Group meetings every three to four years and through input from its family planning Guidelines Steering Group on an as-needed basis. These recommendations will be made available on the WHO web site (www.who.int/reproductive-health). The web site will also provide additional information determined by WHO to be relevant to these recommendations, pending the next formal consensus Working Group meeting. Such updates may be particularly warranted for issues where the evidence base may change rapidly. WHO encourages research to address key unresolved issues for establishing medical eligibility criteria for contraceptive use. WHO also invites comments and suggestions for improving this guidance.

Overview

In 1999, WHO reviewed its family planning guidance and determined that the creation of new evidence-based guidelines was warranted. Accordingly, WHO initiated a new series of evidence-based family planning guidelines beginning with the second edition of *Improving access to quality care in family planning. Medical eligibility criteria for contraceptive use*, published in 2000. The first two cornerstones of this evidence-based series (Figure 1) are this document, the *Medical eligibility criteria for contraceptive use*, which provides guidance regarding “who” can use contraceptive methods safely and the *Selected practice recommendations for contraceptive use*, which provides guidance regarding “how” to use contraceptive methods safely and effectively. These two documents provide evidence-based guidance for choosing (the *Medical eligibility criteria for contraceptive use*) and for using (the *Selected practice recommendations for contraceptive use*) contraceptive methods. The third and fourth cornerstones, a decision-making tool for family planning clients and providers and a handbook for family planning providers, are being prepared as practical tools to improve the quality of family planning counselling and service delivery. These two tools incorporate the *Medical eligibility criteria for contraceptive use* and the *Selected practice recommendations for contraceptive use*. All four cornerstones are best interpreted and used in a broader context of reproductive and sexual health care.

Figure 1. The four cornerstones of family planning guidance



Goal

The goal of this document is to provide policy- and decision-makers and the scientific community with a set of recommendations that can be used for developing or revising national guidelines on medical eligibility criteria for contraceptive use.

The document does not provide rigid guidelines but rather gives recommendations that provide a basis for rationalizing the provision of various contraceptives in view of the most up-to-date information available on the safety of the methods for people with certain health conditions.

Because country situations and programme environments vary so greatly, it is inappropriate to set firm international guidelines on criteria for contraceptive use. However, it is expected that national programmes will use these for updating or developing their own contraceptive eligibility guidelines in the light of their national health policies, needs, priorities and resources. The intent is to help improve access to, and quality of, family planning services. These improvements must be made within the context of users' informed choices and medical safety. Adaptation is not always an easy task and is best done by those well-acquainted with prevailing health conditions, behaviours, and cultures.

Background

Over the past 30 years, there have been significant advances in the development of new contraceptive technologies, including transitions from high-dose to low-dose combined oral contraceptives, and from inert to copper- and levonorgestrel-releasing vaginal IUDs. In addition, combined injectable contraceptives, a combined hormonal patch and ring, and progestogen-only injectables and implants have been introduced. However, current policies and health care practices in some countries are based on scientific studies of contraceptive products that are no longer in wide use, on long-standing theoretical concerns that have never been substantiated, or on the personal preference or bias of service providers. These outdated policies or practices often result in limitations to both the quality of, and the access to, family planning services for clients. This document is intended to update the medical eligibility criteria used in the provision of all hormonal contraceptives, IUDs, barrier methods, fertility awareness-based methods, coitus interruptus, lactational amenorrhoea method, male and female sterilization, and emergency contraception.

Reproductive and sexual health care

“Reproductive rights embrace certain human rights that are already recognised in national laws, international human rights documents and other relevant consensus documents. These rights rest on the recognition of the basic right of all couples and individuals to decide freely and responsibly the number and spacing and timing of their children and to have the information and means to do so, and the right to attain the highest standard of sexual and reproductive health.” (para. 95, Beijing Platform for Action, 1995)

Reproductive and sexual health care including family planning services and information is recognized not only as a key intervention for improving the health of men, women and children but also as a human right. All individuals have the right to access, choice, and the benefits of scientific progress in the selection of family planning methods. A rights-based approach to the provision of contraceptives assumes a holistic view of clients, which includes taking into account clients' sexual and reproductive health care needs and considering all appropriate eligibility criteria in helping clients choose and use a family planning method.

While this document primarily addresses medical eligibility criteria for contraceptive use, considerations of social, behavioural, and other non-medical criteria, particularly client preference, must be taken into account. To provide contraceptive choices to clients in a way that respects and fulfils their human rights necessitates enabling clients to make informed choices for themselves. Women's choices, however, are often imposed or limited by direct or indirect social, economic and cultural factors. From the women's point of view, choices are made in a particular time, societal and cultural context; choices are complex, multifactorial and subject to change. Decision-making for contraceptive methods usually requires the need to make trade-offs among the different methods, with advantages and disadvantages of specific contraceptive methods varying according to individual circumstances, perceptions, and interpretations.

Delivery of care in accordance with the client's human and reproductive rights is fundamental to quality of care. The development of international norms for medical eligibility criteria and practice recommendations for contraceptive use is only one aspect of improving the quality of reproductive health care. Many family planning programmes have included screening, treatment and follow-up procedures that reflect high standards of public health and clinical practice but should not be seen as eligibility requirements for specific contraceptive methods. These procedures include the screening and treatment of cervical cancer, anaemia and sexually transmitted infections (STIs), and the promotion of breastfeeding and cessation of smoking. Such procedures should be strongly encouraged if the human and material resources are available to carry them out, but they should not be seen as prerequisites for the acceptance and use of family planning methods when they are not necessary to establish eligibility for the use or continuation of a particular method.

Issues of service quality and access that affect method use

While this document chiefly addresses medical eligibility criteria, there are many other considerations in the appropriate provision of contraceptive methods including the following service delivery criteria which are universally relevant to the initiation and follow-up of all contraceptive method use.

- a) Clients should be given adequate information in order to make an informed, voluntary choice of a contraceptive method. Information given to clients to help them make this choice should at least include: understanding of the relative effectiveness of the method; correct use of the method; how it works; common side-effects; health risks and benefits of the method; signs and symptoms that would necessitate a return to the clinic; information on return to fertility after discontinuing method use; and information on STI protection.

- b) For those methods that require surgical approaches, insertion, fitting and/or removal by a trained health provider (sterilization, implants, IUDs, diaphragms, cervical caps), appropriately trained personnel in adequately equipped facilities must be available in order for those methods to be offered, and appropriate infection prevention procedures must be followed.
- c) Adequate and appropriate equipment and supplies need to be maintained and held in stock (for example, contraceptive commodities, equipment and supplies for infection prevention procedures).
- d) Service providers should be provided with guidelines (or client cards or other screening tools) to enable them to screen clients appropriately for conditions in which use of certain contraceptive methods would carry unacceptable health risks.
- e) Service providers must be trained in providing family planning counselling to help clients make informed and voluntary decisions about their fertility. Counselling is a key element in quality of care and is also an important part of both initiation and follow-up visits and should respond to clients' needs not only in contraception but also related to sexuality and the prevention of STIs, including infection with the human immunodeficiency virus (HIV).

Effectiveness of method

Contraceptive choice is in part dependent on the effectiveness of the contraceptive method in preventing unplanned pregnancy, which, in turn, is dependent for some methods not only on the protection afforded by the method itself, but also on how consistently and correctly it is used. Table 1 compares the percentage of women experiencing an unintended pregnancy during the first year of contraceptive method use when the method is used perfectly (consistently and correctly) and when it is used typically. Both consistent and correct use can vary greatly with such characteristics as age, income, users' desires to prevent or delay pregnancy, and culture. Methods that depend on consistent and correct use by clients have a wide range of effectiveness. Most men and women tend to be more effective users as they become more experienced with a method. However, programmatic aspects also have a profound effect on how effectively the method will be used.

Table 1. Percentage of women experiencing an unintended pregnancy during the first year of use and the percentage continuing use at the end of the first year, United States of America.

Method (1)	% of women experiencing an unintended pregnancy within the first year of use		% of women continuing use at one year ³
	Typical use ¹ (2)	Perfect use ² (3)	(4)
No method ⁴	85	85	
Spermicides ⁵	29	18	42
Withdrawal	27	4	43
Periodic abstinence	25		51
Calendar		9	
Ovulation method		3	
Sympto-thermal ⁶		2	
Post-ovulation		1	
Cap ⁷			
Parous women	32	26	46
Nulliparous women	16	9	57
Sponge			
Parous women	32	20	46
Nulliparous women	16	9	57
Diaphragm ⁷	16	6	57
Condom ⁸			
Female (Reality)	21	5	49
Male	15	2	53
Combined pill and minipill	8	0.3	68
Combined hormonal patch (Evra)	8	0.3	68
Combined hormonal ring (NuvaRing)	8	0.3	68
DMPA (Depo-Provera)	3	0.3	56
Combined injectable (Lunelle)	3	0.05	56
IUD			
ParaGard (copper T)	0.8	0.6	78
Mirena (LNG-IUS)	0.1	0.1	81
LNG implants (Norplant and Norplant-2)	0.05	0.05	84
Female sterilization	0.5	0.5	100
Male sterilization	0.15	0.10	100

Emergency contraceptive pills: Treatment initiated within 72 hours after unprotected intercourse reduces the risk of pregnancy by at least 75%.

Lactational amenorrhea method: LAM is a highly effective, *temporary* method of contraception.⁹

Source: Trussell J. Contraceptive efficacy. In Hatcher RA, Trussell J, Stewart F, Nelson A, Cates W, Guest F, Kowal D. *Contraceptive Technology: Eighteenth Revised Edition*. New York NY: Ardent Media, 2004.

Note: This table has been adapted from the source document by changing the title, changing the trade names of methods to generic names and by modifying footnotes.

Notes:

- ¹ Among *typical* couples who initiate use of a method (not necessarily for the first time), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason. Estimates of the probability of pregnancy during the first year of typical use for spermicides, withdrawal, periodic abstinence, the diaphragm, the male condom, the pill, and Depo-Provera are taken from the 1995 National Survey of Family Growth corrected for underreporting of abortion; see original source (Trussel J, 2004) cited above for the derivation of the estimates for the other methods.
- ² Among couples who initiate use of a method (not necessarily for the first time) and who use it *perfectly* (both consistently and correctly), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason; see original source (Trussel J, 2004) cited above for the derivation of the estimates for each method.
- ³ Among couples attempting to avoid pregnancy, the percentage who continue to use a method for 1 year.
- ⁴ The percentages becoming pregnant in columns (2) and (3) are based on data from populations where contraception is not used and from women who cease using contraception in order to become pregnant. Among populations where contraception is not used, about 89% become pregnant within 1 year. This estimate was lowered slightly (to 85%) to represent the percentage who would become pregnant within 1 year among women now relying on reversible methods of contraception if they abandoned contraception altogether.
- ⁵ Foams, creams, gels, vaginal suppositories, and vaginal film.
- ⁶ Cervical mucus (ovulation) method supplemented by calendar in the pre-ovulatory and basal body temperature in the post-ovulatory phases.
- ⁷ With spermicidal cream or jelly.
- ⁸ Without spermicides.
- ⁹ However, to maintain effective protection against pregnancy, another method of contraception must be used as soon as menstruation resumes, the frequency or duration of breastfeeds is reduced, bottle feeds are introduced, or the baby reaches 6 months of age.

Conditions that expose a woman to increased risk as a result of unintended pregnancy

Women with conditions that may make pregnancy an unacceptable health risk should be advised that, because of their relatively higher typical-use failure rates, sole use of barrier methods for contraception and behaviour-based methods of contraception may not be the most appropriate choice for them. These conditions are noted in Table 2.

Table 2. Conditions that expose a woman to increased risk as a result of unintended pregnancy

Breast cancer
Complicated valvular heart disease
Diabetes: insulin-dependent; with nephropathy/retinopathy/neuropathy or other vascular disease; or of > 20 years' duration
Endometrial or ovarian cancer
High blood pressure (systolic >160 mm Hg or diastolic >100 mm Hg) [†]
HIV/AIDS*
Ischaemic heart disease
Malignant gestational trophoblastic disease
Malignant liver tumours (hepatoma)
Schistosomiasis with fibrosis of the liver
Severe (decompensated) cirrhosis
Sickle cell disease
STI*
Stroke
Thrombogenic mutations
Tuberculosis

[†] Throughout this document, blood pressure measurements are given in mm Hg. To convert to kPa, multiply by 0.1333. For example, 120/80 mm Hg = 16.0/10.7 kPa.

* Dual protection is strongly recommended for protection against HIV/AIDS and other STIs when a risk of STI/HIV transmission exists. This can be achieved through the simultaneous use of condoms with other methods or the consistent and correct use of condoms alone.

Return to fertility

The use of contraceptive methods, with the exception of male and female sterilization, does not result in an irreversible change in fertility. Return to fertility is prompt with all methods, with the exception of DMPA and NET-EN; the median delay in return to fertility with these methods is 10 and 6 months, respectively, from the date of the last injection, regardless of the duration of their use. Male and female sterilization should be regarded as permanent methods and all individuals and couples considering these methods should be counselled accordingly. No other methods result in permanent infertility.

STIs and contraception: Dual protection

While the development of international norms for contraceptive provision is essential for quality of care in services, the social, cultural and behavioural context of each client must also be considered. In this regard, the problems of exposure to STIs, including HIV, deserve special consideration because of the equal importance of preventing pregnancy and preventing transmission of infection. When a risk of STI/HIV transmission exists, it is important that health care providers strongly recommend dual protection to all persons at significant risk, either through the simultaneous use of condoms with other methods or through the consistent and correct use of condoms alone for both pregnancy prevention and disease prevention. Women and men seeking contraceptive advice must always be reminded of the importance of condom use for preventing the transmission of STI/HIV and such use should be encouraged and facilitated where appropriate. Male latex condoms are proven to be highly effective against STI/HIV when used consistently and correctly.

Method of work

This document builds on a process initiated in 1994 that culminated in the 1996 publication of the document, *Improving access to quality care in family planning: medical eligibility criteria for contraceptive use*. In the initial process, which was created to reach agreement on appropriate eligibility criteria for widely used contraceptive methods, a number of agencies and organizations collaborated in an in-depth review of the epidemiological and clinical evidence relevant to medical eligibility criteria of well established contraceptive methods. The process involved comparing the eligibility criteria used by different agencies for various contraceptives, preparing summaries of published medical and epidemiological literature relevant to medical eligibility criteria, and preparing a draft classification for review by a larger group of experts and agencies. Two expert Working Group meetings were organized by WHO, in March 1994 and May 1995, to review the background classifications and to formulate recommendations; publication of the document followed in 1996.

The first revision of the 1996 document was based on the recommendations of an expert Working Group meeting held at WHO on 8-10 March 2000, that brought together 32 participants from 17 countries, including representatives of many agencies and organizations. The Working Group reviewed new evidence since the last Working Group meetings in 1994 and 1995. This new evidence was primarily obtained from a systematic review of the most recent literature, which was conducted to identify and summarize new evidence for medical eligibility criteria of contraceptive methods.

This third edition of the document is based on the recommendations of an expert Working Group meeting held at WHO on 21–24 October 2003, that brought together 36 participants from 18 countries, including representatives of many agencies and organizations. The Working Group was comprised of international family planning experts, including clinicians, epidemiologists, policy-makers and programme experts. The Working Group also included experts in evidence identification and synthesis and users of the guideline. A Guideline Steering Group was established for this edition. All members of the Working Group were asked to declare conflicts of interests and none were declared.

Using a system to identify new evidence on an ongoing basis (the Continuous Identification of Research Evidence or CIRE, www.infoforhealth.org/cire/cire_pub.pl), WHO identified 151

current recommendations for which new evidence was available since the second edition. WHO also decided to develop recommendations for 3 new conditions and 3 new contraceptive methods in this third edition. Systematic reviews were conducted to appraise the complete body of evidence for these 151 recommendations and for the new conditions and new methods. A systematic, comprehensive search of bibliographic databases, such as MEDLINE, yielded all primary studies, through August 2003, that described use of contraceptive methods among women with certain conditions (e.g., the risk of stroke for women with migraines who use COCs). The purpose of these systematic reviews was to identify direct evidence for the appropriateness of contraceptive method use by women with selected conditions. Information on indirect evidence or theoretical considerations was obtained for these recommendations when direct evidence was sought but not available. The strength and quality of the evidence was graded using the Grades of Recommendation Assessment, Development, and Evaluation (GRADE) system (www.gradeworkinggroup.org). The grading of the evidence was provided to the Working Group as each relevant recommendation was considered. Issues of cost were considered primarily in terms of availability and access to contraceptive services, as well as potential resource constraints. Programmatic implications of the recommendations were also considered by the Working Group. The recommendations primarily concern safety issues and these issues were considered in light of their applicability in a variety of settings.

For most recommendations (method/condition combinations), there are a limited number of studies that address the use of a specific method by women with a specific condition. Thus, most of the decisions regarding eligibility criteria using evidence were often necessarily based on extrapolations from studies that primarily included healthy women, as well as on theoretical considerations and expert opinion. Evidence was particularly limited for newer products and for those with limited usage. The total body of evidence considered by the Working Group included:

- evidence based on direct studies or observations of the contraceptive method used by women (or men) with the condition;
- evidence derived from effects of the contraceptive method used by women (or men) without the condition;
- indirect evidence or theoretical concerns based on studies of suitable animal models, human laboratory studies, or analogous clinical situations.

Where the Working Group had a systematic review of the evidence to consider as they made a recommendation, the evidence is cited in this document alongside the recommendation. The recommendations for which no evidence is cited were based on expert opinion and/or evidence obtained from sources other than systematic reviews. As noted below, over 1000 of the recommendations in this edition are unchanged from those made in the first edition. The evidence for the first edition was provided to the 1994 and 1995 Working Groups in a series of background papers prepared for the project.

The second edition included 1287 recommendations. These recommendations are widely used globally and, therefore, WHO determined that any changes should be based on new evidence unless there was a compelling reason to do otherwise. The Guideline Steering Group, which convened on 21 October 2003, proposed that the expert Working Group consider only those recommendations from the second edition for which there was new evidence or for which a compelling case had been made. The Working Group concurred

with this proposal on 22 October and, thus, the remainder of the Working Group meeting was focused on 151 current recommendations, three new conditions and three new contraceptive methods.

The Working Group was charged with determining the eligibility criteria for each condition and method of contraception by selecting a category (1 through 4, as described below). Where the Working Group determined that guidance in addition to the category was required, that guidance was provided by the Working Group as a "Clarification". Where new evidence was considered by the Working Group, this evidence has been summarized and presented under the heading "Evidence," in the column labelled "Clarifications/Evidence". In addition to the clarifications of guidance and the summaries of the evidence, comments have been provided at the end of each contraceptive method section by the WHO Secretariat for selected methods/conditions.

The final list of 1705 recommendations was approved by all members of the Guideline Steering Group and the Working Group at the completion of the meeting on 24 October 2003.

How to use this document

The present document is intended to be used by policy-makers, family planning programme managers and the scientific community. It aims to provide guidance to national family planning/reproductive health programmes in the preparation of guidelines for service delivery of contraceptives. It should not be seen or used as the actual guidelines but rather as a reference.

The guidance in this document is intended for interpretation at country and programme levels in a manner that reflects the diversity of situations and settings in which contraceptives are provided. While it is unlikely that the classification of categories in this document would change during this process, it is very likely that the application of these categories at country level will vary. In particular, the level of clinical knowledge and experience of various types of providers and the resources available at the service delivery point will have to be taken into consideration.

Using the tables

The Working Group addressed medical criteria for the initiation and continuation of use of all methods evaluated. The issue of continuation criteria is clinically relevant whenever a woman develops the condition while she is using the method. When the Working Group determined that categories for initiation and continuation were different, these differences are noted in the columns 'I=Initiation' and 'C=Continuation'. Where I and C are not denoted, the category is the same for initiation and continuation of use.

On the basis of this classification system, the eligibility criteria for initiating and continuing use of a specific contraceptive method are presented in this document in a set of tables. The first column indicates the condition. Several conditions were subdivided to differentiate between varying degrees of the condition. The second column classifies the condition for

initiation and/or continuation into one of the four categories described below. If necessary, the third column gives clarification or evidence regarding the classification, as described in the section above.

A summary table is included at the end of the document covering medical eligibility criteria by condition for hormonal methods and IUDs. A summary of the conditions or categories that were revised for this edition is included at the end of this section.

TYPE OF CONTRACEPTIVE		
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/ EVIDENCE
Condition	Condition classified from 1 to 4 The categories for fertility awareness-based methods and surgical sterilization are described at the beginning of the relevant section.	Clarifications and evidence regarding the classification

NA denotes a condition for which a ranking was not given by the Working Group but for which clarifications have been provided.

Classification of categories

The medical eligibility criteria in this document were based on the method of work described above and aim to ensure an adequate margin of safety.

Each condition was defined as representing either an individual's characteristics (e.g., age, history of pregnancy) or a known pre-existing medical/pathological condition (e.g., diabetes, hypertension). It is expected that national and institutional health and service delivery environments will decide the most suitable means for screening for conditions according to their public health importance. Client history will often be the most appropriate approach.

The conditions affecting eligibility for the use of each contraceptive method were classified under one of the following four categories:

- | |
|---|
| <ol style="list-style-type: none"> 1. A condition for which there is no restriction for the use of the contraceptive method. 2. A condition where the advantages of using the method generally outweigh the theoretical or proven risks. 3. A condition where the theoretical or proven risks usually outweigh the advantages of using the method. 4. A condition which represents an unacceptable health risk if the contraceptive method is used. |
|---|

Using the categories in practice

Categories 1 and 4 are self-explanatory. Classification of a method/condition as category 2 indicates the method can generally be used, but careful follow-up may be required. However, provision of a method to a woman with a condition classified as category 3 requires careful clinical judgement and access to clinical services; for such a woman, the severity of the condition and the availability, practicality, and acceptability of alternative methods should be taken into account. For a method/condition classified as category 3, use of that method is not usually recommended unless other more appropriate methods are not available or acceptable. Careful follow-up will be required.

Where resources for clinical judgement are limited, such as in community-based services, the four-category classification framework can be simplified into two categories. With this simplification, a classification of Category 3 indicates that a woman is not medically eligible to use the method.

CATEGORY	WITH CLINICAL JUDGEMENT	WITH LIMITED CLINICAL JUDGEMENT
1	Use method in any circumstances	Yes (Use the method)
2	Generally use the method	
3	Use of method not usually recommended unless other more appropriate methods are not available or not acceptable	No (Do not use the method)
4	Method not to be used	

Programmatic implications

Programmatic issues that need to be addressed include:

- informed choice,
- elements of quality of care,
- essential screening procedures for administering the methods,
- provider training and skills,
- referral and follow-up for contraceptive use as appropriate.

In the application of the eligibility criteria to programmes, service delivery practices that are essential for the safe use of the contraceptive should be distinguished from practices that may be appropriate for good health care but are not related to use of the method. The promotion of good health care practices unrelated to safe contraception should be considered neither as a prerequisite nor as an obstacle to the provision of a contraceptive method, but as complementary to it.

As a next step, the recommendations on eligibility criteria need to be considered in light of country circumstances, so as to be applicable to providers at all levels of the service delivery system. Countries will need to determine how far and by what means it may be possible to extend their services to the more peripheral levels. This may involve upgrading both staff and facilities where feasible and affordable, or may require the extension of the skills of certain categories of health personnel or a modest addition of equipment and supplies, and redeployment of space. It will also be necessary to address questions of misperceptions sometimes held by providers and users on the risks and side-effects of the methods and to look closely at the needs and perspectives of women and men in the context of informed choice.

Clients with special needs

Medical eligibility criteria address contraceptive use by people with specific medical conditions. In addition, contraceptive provision to people with special needs requires further consideration. Individuals with a physical disability represent such a group. Decisions on appropriate contraception must take into account the nature of the disability, the expressed desires of the individual and the nature of the method. Decisions must be based on informed choice. Similar considerations should be given to individuals with mental disability or with serious psychiatric disease. Where the nature of the condition does not allow for informed choice, contraceptives should be provided only after full discussion with all parties including guardians or care-givers. The reproductive rights of the individual must be considered in any such decisions.

Adolescents

In general, adolescents are eligible to use any method of contraception and must have access to a variety of contraceptive choices. Age alone does not constitute a medical reason for denying any method to adolescents. While some concerns have been expressed regarding the use of certain contraceptive methods in adolescents (e.g., the use of progestogen-only injectables by those below 18 years), these concerns must be balanced against the advantages of avoiding pregnancy. It is clear that many of the same eligibility criteria that apply to older clients apply to young people. However, some conditions (e.g., cardiovascular disorders) that may limit use of some methods in older women do not generally affect young people since these conditions are rare in this age group. Social and behavioural issues should be important considerations in the choice of contraceptive methods by adolescents. For example, in some settings, adolescents are also at increased risk for STIs, including HIV. While adolescents may choose to use any one of the contraceptive methods available in their communities, in some cases, using methods that do not require a daily regimen may be more appropriate. Adolescents, married or unmarried, have also been shown to be less tolerant of side-effects and therefore have high discontinuation rates. Method choice may also be influenced by factors such as sporadic patterns of intercourse and the need to conceal sexual activity and contraceptive use. For instance, sexually active adolescents who are unmarried have very different needs from those who are married and want to postpone, space or limit pregnancy. Expanding the number of method choices offered can lead to improved satisfaction, increased acceptance and increased prevalence of contraceptive use. Proper education and counselling both before and at the time of method selection can help adolescents address their specific

problems and make informed and voluntary decisions. Every effort should be made to prevent service and method costs from limiting the options available.

Summary of changes from the second edition

A summary of the classification changes or major condition modifications from the second edition is given in Table 3.

It is recognized that some of the eligibility criteria in this report will need to be reviewed in the light of new research findings from studies being completed and/or currently in progress. It is intended that this document will be updated on a continual basis in order to reflect the latest scientific evidence and findings.

Table 3. Summary of changes from the second edition

(Conditions for which there was a classification change for one or more methods or a major modification to the condition description. Changed classifications are highlighted in blue.)

CONDITION	COC	CIC	POP	DMPA NET-EN	LNG/ ETG	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation							
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY							
OBESITY ≥30 kg/m ² body mass index (BMI)	2	2	1	1	1	1	1
CARDIOVASCULAR DISEASE							
KNOWN THROMBOGENIC MUTATIONS (e.g. Factor V Leiden; Prothrombin mutation; Protein S, Protein C and Antithrombin deficiencies)	4	4	2	2	2	1	2
DEPRESSIVE DISORDERS							
DEPRESSIVE DISORDERS	1	1	1	1	1	1	1
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS							
UTERINE FIBROIDS							
a) Without distortion of the uterine cavity	1	1	1	1	1	1	1
b) With distortion of the uterine cavity	1	1	1	1	1	4	4
PELVIC INFLAMMATORY DISEASE (PID)							
a) Past PID (assuming no current risk factors of STIs)						I	C
(i) with subsequent pregnancy	1	1	1	1	1	1	1
ii) without subsequent pregnancy	1	1	1	1	1	2	2
b) PID - current	1	1	1	1	1	4	2

CONDITION	COC	CIC	POP	DMPA NET-EN	LNG/ ETG	Cu-IUD		LNG-IUD	
I = Initiation, C = Continuation									
STIs						I	C	I	C
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	1	1	1	1	4	2	4	2
b) Other STIs (excluding HIV and hepatitis)	1	1	1	1	1	2	2	2	2
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	1	1	1	1	2	2	2	2
d) Increased risk of STIs	1	1	1	1	1	2/3	2	2/3	2
HIV/AIDS									
HIGH RISK OF HIV						I	C	I	C
	1	1	1	1	1	2	2	2	2
HIV-INFECTED	1	1	1	1	1	2	2	2	2
AIDS	1	1	1	1	1	3	2	3	2
Clinically well on ARV therapy	See ANTIRETROVIRAL THERAPY below					2	2	2	2
DRUG INTERACTIONS									
DRUGS WHICH AFFECT LIVER ENZYMES									
a) Rifampicin	3	2	3	2	3	1		1	
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	3	2	3	2	3	1		1	
ANTIBIOTICS (excluding rifampicin)									
a) Griseofulvin	2	1	2	1	2	1		1	
b) Other antibiotics	1	1	1	1	1	1		1	
ANTIRETROVIRAL THERAPY						I	C	I	C
	2	2	2	2	2	2/3	2	2/3	2

In addition, the following changes were made which are not included in the summary of changes:

1. Patch, ring and etonogestrel implants

Three new methods (patch, ring and etonogestrel implants) were added. The patch and ring are grouped with CICs but are given the same category ratings as COCs. The etonogestrel implants are grouped with, and are given the same category ratings as, the levonorgestrel implants.

2. Barrier methods

For the conditions of high risk of HIV, HIV-infected, and AIDS, spermicide use is a Category 4.

For the conditions of high risk of HIV, HIV-infected, and AIDS, diaphragm (with spermicide) and cervical cap are Category 3.

3. Female surgical sterilization

The condition of known thrombogenic mutations (e.g. Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies) has been added for female surgical sterilization and is Category A.

The condition of depressive disorders has been added for female surgical sterilization and is a Category C.

For the condition of other STIs (excluding HIV and hepatitis), female surgical sterilization is Category A.

For the condition of previous abdominal or pelvic surgery, female surgical sterilization is Category C.

4. Male surgical sterilization

The condition of young age has been added for male surgical sterilization and is Category C.

The condition of depressive disorders has been added for male surgical sterilization and is Category C.

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	1
Pregnancy.....	1
Age	1
Parity.....	1
Breastfeeding.....	1
Postpartum	1
Post-abortion.....	1
Past ectopic pregnancy.....	1
History of pelvic surgery.....	1
Smoking.....	2
Obesity.....	2
Blood pressure measurement unavailable.....	2
CARDIOVASCULAR DISEASE.....	2
Multiple risk factors for arterial cardiovascular disease.....	2
Hypertension.....	2
History of high blood pressure during pregnancy.....	3
Deep venous thrombosis (DVT)/pulmonary embolism (PE).....	3
Known thrombogenic mutations.....	4
Superficial venous thrombosis.....	4
Current and history of ischaemic heart disease.....	4
Stroke	4
Known hyperlipidaemias.....	4
Valvular heart disease.....	4
NEUROLOGIC CONDITIONS.....	5
Headaches.....	5
Epilepsy	5
DEPRESSIVE DISORDERS	5
Depressive disorders	5
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS.....	5
Vaginal bleeding patterns	5
Unexplained vaginal bleeding	6
Endometriosis	6
Benign ovarian tumours.....	6
Severe dysmenorrhoea.....	6
Trophoblast disease.....	6
Cervical ectropion.....	6
Cervical intraepithelial neoplasia (CIN).....	6
Cervical cancer.....	6
Breast disease.....	6
Endometrial cancer.....	7
Ovarian cancer.....	7
Uterine fibroids.....	7
Pelvic inflammatory disease (PID).....	7
STIs	7
HIV/AIDS	8
High risk of HIV	8
HIV-infected	8
AIDS	8
OTHER INFECTIONS	8
Schistosomiasis	8
Tuberculosis.....	8
Malaria	8

ENDOCRINE CONDITIONS	9
Diabetes.....	9
Thyroid disorders	9
GASTROINTESTINAL CONDITIONS	9
Gall-bladder disease	9
History of cholestasis	9
Viral hepatitis	9
Cirrhosis.....	10
Liver tumours	10
ANAEMIAS.....	10
Thalassaemia.....	10
Sickle cell disease.....	10
Iron-deficiency anaemia.....	10
DRUG INTERACTIONS	10
Drugs which affect liver enzymes.....	10
Antibiotics	10
Antiretroviral therapy.....	11
Additional comments	12
References for low-dose combined oral contraceptives.....	14

LOW-DOSE COMBINED ORAL CONTRACEPTIVES

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) \leq 35 μg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY		
PREGNANCY	NA	Clarification: Use of COCs is not required. There is no known harm to the woman, the course of her pregnancy, or the fetus if COCs are accidentally used during pregnancy.
AGE*		
a) Menarche to < 40 years	1	
b) \geq 40 years	2	
PARITY		
a) Nulliparous	1	
b) Parous	1	
BREASTFEEDING*		
a) < 6 weeks postpartum	4	
b) \geq 6 weeks to < 6 months postpartum (primarily breastfeeding)	3	
c) \geq 6 months postpartum	2	
POSTPARTUM* (in non-breastfeeding women)		
a) < 21 days	3	
b) \geq 21 days	1	
POST-ABORTION		
a) First trimester	1	Clarification: COCs may be started immediately post-abortion.
b) Second trimester	1	
c) Immediate post-septic abortion	1	
PAST ECTOPIC PREGNANCY*	1	
HISTORY OF PELVIC SURGERY	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
SMOKING a) Age < 35 years b) Age ≥ 35 years (i) <15 cigarettes/day (ii) ≥15 cigarettes/day	2 3 4	Evidence: COC users who smoked were at increased risk of cardiovascular diseases, especially myocardial infarction, compared with those who did not smoke. Studies also showed an increased risk of myocardial infarction with increasing number of cigarettes smoked per day. ¹⁻¹²
OBESITY ≥ 30 kg/m ² body mass index (BMI)	2	Evidence: Obese women who used COCs were at increased risk of VTE compared with non-users. The absolute risk of VTE remained small. Data are limited regarding the impact of obesity on COC effectiveness. ^{6, 13, 14}
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA	Clarification: It is desirable to have blood pressure measurements taken before initiation of COC use. However, in some settings blood pressure measurements are unavailable. In many of these settings pregnancy morbidity and mortality risks are high, and COCs are one of the few methods widely available. In such settings, women should not be denied use of COCs simply because their blood pressure cannot be measured.
CARDIOVASCULAR DISEASE		
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	3/4	Clarification: When a woman has multiple major risk factors, any of which alone would substantially increase the risk of cardiovascular disease, use of COCs may increase her risk to an unacceptable level. However, a simple addition of categories for multiple risk factors is not intended; for example, a combination of two risk factors assigned a category 2 may not necessarily warrant a higher category.
HYPERTENSION*		
For all categories of hypertension, classifications are based on the assumption that no other risk factors for cardiovascular disease exist. When multiple risk factors do exist, risk of cardiovascular disease may increase substantially. A single reading of blood pressure level is not sufficient to classify a woman as hypertensive.		
a) History of hypertension, where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	3	Clarification: Evaluation of cause and level of hypertension is recommended, as soon as feasible. Evidence: Women who did not have a blood pressure check before COC use had an increased risk of acute myocardial infarction and stroke. ¹⁵⁻¹⁹

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) \leq 35 μg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
HYPERTENSION (Cont'd)		
b) Adequately controlled hypertension, where blood pressure CAN be evaluated	3	Clarification: Women adequately treated for hypertension are at reduced risk of acute myocardial infarction and stroke as compared with untreated women. Although there are no data, COC users with adequately controlled and monitored hypertension should be at reduced risk of acute myocardial infarction and stroke compared with untreated hypertensive COC users.
c) Elevated blood pressure levels (properly taken measurements)		Evidence: Among women with hypertension, COC users were at increased risk of stroke, acute myocardial infarction, and peripheral arterial disease compared with non-users. ^{1,3,9-11, 15-31}
(i) systolic 140-159 or diastolic 90-99	3	
(ii) systolic \geq 160 or diastolic \geq 100	4	
d) Vascular disease	4	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	2	Evidence: Women who had a history of high blood pressure in pregnancy, who also used COCs, had an increased risk of myocardial infarction and venous thromboembolism, compared with COC users who did not have a history of high blood pressure during pregnancy. The absolute risks of acute myocardial infarction and venous thromboembolism in this population remained small. ^{11, 17-19, 21, 32-37}
DEEP VEIN THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)*		
a) History of DVT/PE	4	
b) Current DVT/PE	4	
c) Family history of DVT/PE (first-degree relatives)	2	
d) Major surgery		
(i) with prolonged immobilization	4	
(ii) without prolonged immobilization	2	
e) Minor surgery without immobilization	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) \leq 35 μg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	4	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening. Evidence: Among women with thrombogenic mutations, COC users had a two to twenty-fold higher risk of thrombosis than non-users. ³⁸⁻⁵¹
SUPERFICIAL VENOUS THROMBOSIS* a) Varicose veins b) Superficial thrombophlebitis	1 2	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE*	4	
STROKE* (history of cerebrovascular accident)	4	
KNOWN HYPERLIPIDAEMIAS	2/3	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening. While some types of hyperlipidaemias are risk factors for vascular disease, the category should be assessed according to the type, its severity, and the presence of other cardiovascular risk factors.
VALVULAR HEART DISEASE* a) Uncomplicated b) Complicated (pulmonary hypertension, risk of atrial fibrillation, history of subacute bacterial endocarditis)	2 4	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
NEUROLOGIC CONDITIONS		
HEADACHES*	I	C
a) Non-migrainous (mild or severe)	1	2
b) Migraine (i) without aura <i>Age < 35</i> <i>Age ≥ 35</i> (ii) with aura, at any age	2	3
	3	4
	4	4
EPILEPSY	1	
DEPRESSIVE DISORDERS		
DEPRESSIVE DISORDERS	1	
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS		
VAGINAL BLEEDING PATTERNS*		
a) Irregular pattern <i>without</i> heavy bleeding	1	
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	1	
	Clarification: Unusually heavy bleeding should raise the suspicion of a serious underlying condition.	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
UNEXPLAINED VAGINAL BLEEDING* (suspicious for serious condition) Before evaluation	2	Clarification: If pregnancy or an underlying pathological condition (such as pelvic malignancy) is suspected, it must be evaluated and the category adjusted after evaluation.
ENDOMETRIOSIS*	1	
BENIGN OVARIAN TUMOURS (including cysts)	1	
SEVERE DYSMENORRHOEA	1	Evidence: There was no increased risk of side-effects with COC use among women with dysmenorrhoea compared to women not using COCs. Some COC users had a reduction in pain and bleeding. ^{62, 63}
TROPHOBLAST DISEASE a) Benign gestational trophoblastic disease b) Malignant gestational trophoblastic disease	1 1	Evidence: Among women with benign or malignant gestational trophoblastic disease, there was no difference in mean times to hCG normalization or incidence of postmolar trophoblastic disease for COC users compared to non-hormonal users. ⁶⁴⁻⁷¹
CERVICAL ECTROPION*	1	
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	2	Evidence: Among women with persistent HPV infection, long-term COC use (≥ 5 years) may increase the risk of carcinoma in situ and invasive carcinoma. ⁷²
CERVICAL CANCER* (awaiting treatment)	2	
BREAST DISEASE* a) Undiagnosed mass b) Benign breast disease c) Family history of cancer	2 1 1	Clarification: Evaluation should be pursued as early as possible. Evidence: Among COC users with a family history of breast cancer, there was no increased risk of breast cancer compared with non-COC users with a family history of breast cancer. ⁷³⁻⁸⁰ Among women with BRCA1 mutations, COC users may have a small increased risk of breast cancer compared with non-users. ⁸¹⁻⁸³

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) \leq 35 μg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
BREAST DISEASE (Cont'd)		
d) Breast cancer		
(i) current	4	
(ii) past and no evidence of current disease for 5 years	3	
ENDOMETRIAL CANCER*	1	
OVARIAN CANCER*	1	
UTERINE FIBROIDS*		
a) Without distortion of the uterine cavity	1	
b) With distortion of the uterine cavity	1	
PELVIC INFLAMMATORY DISEASE (PID)*		
a) Past PID (assuming no current risk factors for STIs)		
(i) with subsequent pregnancy	1	
(ii) without subsequent pregnancy	1	
b) PID - current	1	
STIs*		
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	
b) Other STIs (excluding HIV and hepatitis)	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
STIs (Cont'd)		
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	
d) Increased risk of STIs	1	Evidence: Evidence suggests that there may be an increased risk of chlamydial cervicitis among COC users at high risk of STIs. For other STIs, there is either evidence of no association between COC use and STI acquisition or limited evidence to draw any conclusions. ⁸⁴⁻¹⁶⁰
HIV/AIDS		
HIGH RISK OF HIV*	1	Evidence: Overall, evidence is inconsistent regarding whether there is any increased risk of HIV acquisition among COC users compared with non-users. ¹⁶¹⁻¹⁹⁸
HIV-INFECTED	1	Evidence: Limited evidence suggests no association between COC use and changes in RNA levels or CD4 counts among HIV-infected women. There is also limited evidence showing no association between COC use and female to male HIV transmission, and mixed results regarding increased risk of HIV and herpes simplex virus (HSV) shedding among HIV-infected women using hormonal contraception. ^{161, 199-204}
AIDS On ARV therapy	1 2	Clarification: If a woman is taking antiretroviral (ARV) therapy, refer to the section on drug interactions. Because there may be drug interactions between hormonal contraceptives and ARVs, AIDS with ARV therapy is classified as Category 2.
OTHER INFECTIONS		
SCHISTOSOMIASIS		
a) Uncomplicated	1	Evidence: Among women with uncomplicated schistosomiasis, COC use had no adverse effects on liver function. ²⁰⁵⁻²¹¹
b) Fibrosis of liver (if severe, see cirrhosis)	1	
TUBERCULOSIS		
a) Non-pelvic	1	Clarification: If a woman is taking rifampicin, refer to the section on drug interactions. Rifampicin is likely to decrease COC effectiveness.
b) Known pelvic	1	
MALARIA	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) \leq 35 μg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
ENDOCRINE CONDITIONS		
DIABETES*		
a) History of gestational disease	1	
b) Non-vascular disease		
(i) non-insulin dependent	2	
(ii) insulin dependent	2	
c) Nephropathy/ retinopathy/ neuropathy	3/4	Clarification: The category should be assessed according to the severity of the condition.
d) Other vascular disease or diabetes of > 20 years' duration	3/4	Clarification: The category should be assessed according to the severity of the condition.
THYROID DISORDERS		
a) Simple goitre	1	
b) Hyperthyroid	1	
c) Hypothyroid	1	
GASTROINTESTINAL CONDITIONS		
GALL-BLADDER DISEASE*		
a) Symptomatic		
(i) treated by cholecystectomy	2	
(ii) medically treated	3	
(iii) current	3	
b) Asymptomatic	2	
HISTORY OF CHOLESTASIS*		
a) Pregnancy-related	2	
b) Past COC-related	3	
VIRAL HEPATITIS*		
a) Active	4	
b) Carrier	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
CIRRHOSIS*		
a) Mild (compensated)	3	
b) Severe (decompensated)	4	
LIVER TUMOURS*		
a) Benign (adenoma)	4	
b) Malignant (hepatoma)	4	
ANAEMIAS		
THALASSAEMIA*	1	
SICKLE CELL DISEASE	2	
IRON-DEFICIENCY ANAEMIA*	1	
DRUG INTERACTIONS		
DRUGS WHICH AFFECT LIVER ENZYMES		
a) Rifampicin	3	Clarification: Although the interaction of rifampicin or certain anticonvulsants with COCs is not harmful to women, it is likely to reduce the effectiveness of COCs. Use of other contraceptives should be encouraged for women who are long-term users of any of these drugs. Whether increasing the hormone dose of COCs is of benefit remains unclear. Evidence: Use of rifampicin and certain anticonvulsants decreased the contraceptive effectiveness of COCs. ²¹²⁻²³⁷
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	3	
ANTIBIOTICS (excluding rifampicin)		
a) Griseofulvin	2	Evidence: The contraceptive effectiveness of COCs was not affected by coadministration of most broad-spectrum antibiotics. ²³⁸⁻²⁹⁰
b) Other antibiotics	1	

* See also additional comments at end of table

LOW-DOSE COMBINED ORAL CONTRACEPTIVES (COCs) ≤ 35 µg of ethinylestradiol	COCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY I=Initiation C=Continuation	CLARIFICATIONS/EVIDENCE
ANTIRETROVIRAL THERAPY	<p style="text-align: center;">2</p>	<p>Clarification: It is important to note that antiretroviral drugs (ARV) have the potential to either decrease or increase the bioavailability of steroid hormones in hormonal contraceptives. The limited data available (outlined in Annex 1) suggest that potential drug interactions between many ARVs (particularly some non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs)) and hormonal contraceptives may alter safety and effectiveness of both the hormonal contraceptives and the ARVs. It is not known whether the contraceptive effectiveness of progestogen-only injectable contraceptives (such as depot medroxyprogesterone acetate and norethisterone enantate) would be compromised, as these methods provide higher blood hormone levels than other progestogen-only hormonal contraceptives, as well as than combined oral contraceptives. Studies are underway to evaluate potential interactions between depot medroxyprogesterone acetate and selected PI and NNRTI drugs. Thus, if a woman on ARV treatment decides to initiate or continue hormonal contraceptive use, the consistent use of condoms is recommended for preventing HIV transmission and may also compensate for any possible reduction in the effectiveness of the hormonal contraceptive.</p> <p>Evidence: See Annex 1.</p>

* See also additional comments at end of table

Additional comments

AGE

Menarche to < 40 years: Theoretical concerns about the use of combined hormonal contraceptives among young adolescents have not been substantiated.

≥ 40 years: The risk of cardiovascular disease increases with age and may also increase with combined hormonal contraceptive use. In the absence of other adverse clinical conditions, combined hormonal contraceptives can be used until menopause.

BREASTFEEDING

< 6 weeks postpartum: There is some theoretical concern that the neonate may be at risk due to exposure to steroid hormones during the first 6 weeks postpartum.

≥ 6 weeks to < 6 months (primarily breastfeeding): Use of COCs during breastfeeding diminishes the quantity of breast milk, decreases the duration of lactation, and may thereby adversely affect the growth of the infant.

POSTPARTUM

< 21 days: There is some theoretical concern regarding the association between combined hormonal contraceptive use up to 3 weeks postpartum and risk of thrombosis in the mother. Blood coagulation and fibrinolysis are essentially normalized by 3 weeks postpartum.

PAST ECTOPIC PREGNANCY

The risk of future ectopic pregnancy is increased among women who have had an ectopic pregnancy in the past. Combined hormonal contraceptives provide protection against pregnancy in general, including ectopic gestation.

HYPERTENSION

Vascular disease: Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

DEEP VEIN THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)

Family history of DVT/PE (first-degree relatives): Some conditions which increase the risk of DVT/PE are heritable.

Major surgery: The degree of risk of DVT/PE associated with major surgery varies depending on the length of time that a woman is immobilized. There is no need to stop combined hormonal contraceptives prior to female surgical sterilization.

SUPERFICIAL VEIN THROMBOSIS

Varicose veins: Varicose veins are not risk factors for DVT/PE.

CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE

Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

STROKE

Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

VALVULAR HEART DISEASE

Among women with valvular heart disease, combined hormonal contraceptive use may further increase the risk of arterial thrombosis; women with complicated valvular heart disease are at greatest risk.

HEADACHES

Aura is a specific focal neurologic symptom. For more information on this and other diagnostic criteria, see: Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders, 2nd Edition. Cephalalgia. 2004; 24 (Suppl 1): 1- 150.

http://216.25.100.131/ihscommon/guidelines/pdfs/ihc_II_main_no_print.pdf

VAGINAL BLEEDING PATTERNS

Irregular menstrual bleeding patterns are common among healthy women.

UNEXPLAINED VAGINAL BLEEDING

There are no conditions that cause vaginal bleeding that will be worsened in the short term by use of combined hormonal contraceptives.

ENDOMETRIOSIS

Combined hormonal contraceptives do not worsen, and may alleviate, the symptoms of endometriosis.

CERVICAL ECTROPION

Cervical ectropion is not a risk factor for cervical cancer, and there is no need for restriction of combined hormonal contraceptive use.

CERVICAL CANCER (awaiting treatment)

There is some theoretical concern that combined hormonal contraceptive use may affect prognosis of the existing disease. While awaiting treatment, women may use combined hormonal contraceptives. In general, treatment of this condition renders a woman sterile.

BREAST DISEASE

Family history of cancer: Women with BRCA1 or BRCA2 mutations have a much higher baseline risk of breast cancer than women who do not have these mutations. Most women with a family history of breast cancer do not have these mutations.

Breast cancer: Breast cancer is a hormonally sensitive tumour, and the prognosis of women with current or recent breast cancer may worsen with combined hormonal contraceptive use.

ENDOMETRIAL CANCER

COC use reduces the risk of developing endometrial cancer. While awaiting treatment, women may use COCs. In general, treatment of this condition renders a woman sterile.

OVARIAN CANCER

COC use reduces the risk of developing ovarian cancer. While awaiting treatment, women may use COCs. In general, treatment of this condition renders a woman sterile.

UTERINE FIBROIDS

COCs do not appear to cause growth of uterine fibroids.

PELVIC INFLAMMATORY DISEASE (PID)

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs.

STIs

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs.

HIGH RISK OF HIV

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs.

DIABETES

Although carbohydrate tolerance may change with combined hormonal contraceptive use, the major concerns are vascular disease due to diabetes and additional risk of arterial thrombosis due to combined hormonal contraceptive use.

GALL-BLADDER DISEASE

COCs may cause a small increased risk of gall-bladder disease. There is also concern that COCs may worsen existing gall-bladder disease.

HISTORY OF CHOLESTASIS

Pregnancy-related: History of pregnancy-related cholestasis may predict an increased risk of developing COC-associated cholestasis.

Past COC-related: History of COC-related cholestasis predicts an increased risk with subsequent COC use.

VIRAL HEPATITIS

Active: COCs are metabolized by the liver, and their use may adversely affect women whose liver function is compromised.

CIRRHOSIS

COCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised.

LIVER TUMOURS

COCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. In addition, COC use may enhance the growth of tumours.

THALASSAEMIA

There is anecdotal evidence from countries where thalassaemia is prevalent that COC use does not worsen the condition.

IRON-DEFICIENCY ANAEMIA

Combined hormonal contraceptive use may decrease menstrual blood loss.

References for low-dose combined oral contraceptives

1. Gillum LA, Mamidipudi SK, Johnston SC. Ischemic stroke risk with oral contraceptives: a meta-analysis. *JAMA*, 2000, 284:72-8.
2. Jick SS et al. Oral contraceptives and breast cancer. *British Journal of Cancer*, 1989, 59:618-21.
3. Khader YS et al. Oral contraceptives use and the risk of myocardial infarction: a meta-analysis. *Contraception*, 2003, 68:11-7.
4. Lawson DH, Davidson JF, Jick H. Oral contraceptive use and venous thromboembolism: absence of an effect of smoking. *BMJ*, 1977, 2:729-30.
5. Lidegaard O, Edstrom B, Kreiner S. Oral contraceptives and venous thromboembolism. A case-control study. *Contraception*, 1998, 57:291-301.
6. Nightingale AL et al. The effects of age, body mass index, smoking and general health on the risk of venous thromboembolism in users of combined oral contraceptives. *European Journal of Contraception & Reproductive Health Care*, 2000, 5:265-74.
7. Petitti DB et al. Risk of vascular disease in women. Smoking, oral contraceptives, noncontraceptive estrogens, and other factors. *JAMA*, 1979, 242:1150-4.
8. Straneva P et al. Smoking, oral contraceptives, and cardiovascular reactivity to stress. *Obstetrics & Gynecology*, 2000, 95:78-83.
9. Tanis BC et al. Oral contraceptives and the risk of myocardial infarction. *New England Journal of Medicine*, 2001, 345:1787-93.
10. Van den Bosch MA et al. The RATIO study: oral contraceptives and the risk of peripheral arterial disease in young women. *Journal of Thrombosis and Haemostasis*, 2003, 1:439-44.
11. WHO. Venous thromboembolic disease and combined oral contraceptives: results of international multicentre case-control study. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *Lancet*, 1995, 346:1575-82.
12. Rosenberg L et al. Low-dose oral contraceptive use and the risk of myocardial infarction. *Archives of Internal Medicine*, 2001, 161:1065-70.
13. Abdollahi MC. Obesity: risk of venous thrombosis and the interaction with coagulation factor levels and oral contraceptive use. *Thrombosis & Haemostasis*, 2003, 89:493-8.
14. Holt VL, Cushing-Haugen KL, Daling JR. Body weight and risk of oral contraceptive failure. *Obstetrics & Gynecology*, 2002, 99:820-7.
15. Heinemann LA et al. Thromboembolic stroke in young women. A European case-control study on oral contraceptives. Transnational Research Group on Oral Contraceptives and the Health of Young Women. *Contraception*, 1998, 57:29-37.
16. Lewis MA et al. The use of oral contraceptives and the occurrence of acute myocardial infarction in young women. Results from the Transnational Study on Oral Contraceptives and the Health of Young Women. *Contraception*, 1997, 56:129-40.
17. WHO. Haemorrhagic stroke, overall stroke risk, and combined oral contraceptives: results of an international, multicentre, case-control study. WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *Lancet*, 1996, 348:505-10.
18. WHO. Ischaemic stroke and combined oral contraceptives: results of an international, multicentre, case-control study. WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *Lancet*, 1996, 348:498-505.
19. WHO. Acute myocardial infarction and combined oral contraceptives: results of an international multicentre case-control study. WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *Lancet*, 1997, 349:1202-9.
20. Collaborative Group for the Study of Stroke in Young Women. Oral contraceptives and stroke in young women: associated risk factors. *JAMA*, 1975, 231:718-22.
21. Croft P, Hannaford PC. Risk factors for acute myocardial infarction in women: evidence from the Royal College of General Practitioners' Oral Contraception Study. *BMJ*, 1989, 298:165-8.
22. D'Avanzo B et al. Oral contraceptive use and risk of myocardial infarction: an Italian case-control study. *Journal of Epidemiology and Community Health*, 1994, 48:324-5.
23. Dunn NR et al. Risk of myocardial infarction in young female smokers. *Heart (British Cardiac Society)*, 1999, 82:581-3.
24. Hannaford PC, Croft PR, Kay CR. Oral contraception and stroke. Evidence from the Royal College of General Practitioners' Oral Contraception Study. *Stroke*, 1994, 25:935-42.
25. Kemmeren JM et al. Risk of Arterial Thrombosis in Relation to Oral Contraceptives (RATIO) study: oral contraceptives and the risk of ischemic stroke. *Stroke*, 2002, 33:1202-8.
26. Lidegaard O. Oral contraception and risk of a cerebral thromboembolic attack: results of a case-control study. *BMJ*, 1993, 306:956-63.

27. Lidegaard O. Oral contraceptives, pregnancy and the risk of cerebral thromboembolism: the influence of diabetes, hypertension, migraine and previous thrombotic disease. *British Journal of Obstetrics & Gynaecology*, 1995, 102:153-9.
28. Lidegaard O, Kreiner S. Contraceptives and cerebral thrombosis: a five-year national case-control study. *Contraception*, 2002, 65:197-205.
29. Lubianca JN, Faccin CS, Fuchs FD. Oral contraceptives: a risk factor for uncontrolled blood pressure among hypertensive women. *Contraception*, 2003, 67:19-24.
30. Narkiewicz K et al. Ambulatory blood pressure in mild hypertensive women taking oral contraceptives. A case-control study. *American Journal of Hypertension*, 1995, 8:249-53.
31. Siritho S et al. Risk of ischemic stroke among users of the oral contraceptive pill: The Melbourne Risk Factor Study (MERFS) Group. *Stroke*, 2003, 34:1575-80.
32. Aberg H, Karlsson L, Melander S. Studies on toxemia of pregnancy with special reference to blood pressure. II. Results after 6-11 years' follow-up. *Uppsala Journal of Medical Sciences*, 1978, 83:97-102.
33. Carmichael SM, Taylor MM, Ayers CR. Oral contraceptives, hypertension, and toxemia. *Obstetrics & Gynecology*, 1970, 35:371-6.
34. Meinel H, Ihle R, Laschinski M. [Effect of hormonal contraceptives on blood pressure following pregnancy-induced hypertension]. [German]. *Zentralblatt für Gynäkologie*, 1987, 109:527-31.
35. Pritchard JA, Pritchard SA. Blood pressure response to estrogen-progestin oral contraceptive after pregnancy-induced hypertension. *American Journal of Obstetrics & Gynecology*, 1977, 129:733-9.
36. Sibai BM et al. Maternal-perinatal outcome associated with the syndrome of hemolysis, elevated liver enzymes, and low platelets in severe preeclampsia-eclampsia. *American Journal of Obstetrics & Gynecology*, 1986, 155:501-9.
37. Sibai BM et al. Pregnancies complicated by HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets): subsequent pregnancy outcome and long-term prognosis. *American Journal of Obstetrics & Gynecology*, 1995, 172:125-9.
38. Andersen BS et al. Third generation oral contraceptives and heritable thrombophilia as risk factors of non-fatal venous thromboembolism. *Thrombosis & Haemostasis*, 1998, 79:28-31.
39. Andersen BS, Olsen J. Oral contraception and factor V Leiden mutation in relation to localization of deep vein thrombosis. *Thrombosis Research*, 1998, 90:191-4.
40. Bloemenkamp KW et al. Enhancement by factor V Leiden mutation of risk of deep-vein thrombosis associated with oral contraceptives containing a third-generation progestagen. *Lancet*, 1995, 346:1593-6.
41. Bloemenkamp KW et al. Higher risk of venous thrombosis during early use of oral contraceptives in women with inherited clotting defects. *Archives of Internal Medicine*, 2000, 160:49-52.
42. de Bruijn SF et al. Case-control study of risk of cerebral sinus thrombosis in oral contraceptive users and in [correction of who are] carriers of hereditary prothrombotic conditions. The Cerebral Venous Sinus Thrombosis Study Group.[erratum appears in *BMJ* 1998 Mar 14;316(7134):822]. *BMJ*, 1998, 316:589-92.
43. Emmerich J et al. Combined effect of factor V Leiden and prothrombin 20210A on the risk of venous thromboembolism--pooled analysis of 8 case-control studies including 2310 cases and 3204 controls. Study Group for Pooled-Analysis in Venous Thromboembolism.[erratum appears in *Thromb Haemost* 2001 Dec;86(6):1598]. *Thrombosis & Haemostasis*, 2001, 86:809-16.
44. Legnani C et al. Venous thromboembolism in young women; role of thrombophilic mutations and oral contraceptive use. *European Heart Journal*, 2002, 23:984-90.
45. Martinelli I et al. High risk of cerebral-vein thrombosis in carriers of a prothrombin-gene mutation and in users of oral contraceptives. *New England Journal of Medicine*, 1998, 338:1793-7.
46. Martinelli I et al. Interaction between the G20210A mutation of the prothrombin gene and oral contraceptive use in deep vein thrombosis. *Arteriosclerosis, Thrombosis & Vascular Biology*, 1999, 19:700-3.
47. Middeldorp S et al. A prospective study of asymptomatic carriers of the factor V Leiden mutation to determine the incidence of venous thromboembolism. *Annals of Internal Medicine*, 2001, 135:322-7.
48. Santamaria A et al. Risk of thrombosis associated with oral contraceptives of women from 97 families with inherited thrombophilia: high risk of thrombosis in carriers of the G20210A mutation of the prothrombin gene. *Haematologica*, 2001, 86:965-71.
49. Spannagl M, Heinemann LA, Schramm W. Are factor V Leiden carriers who use oral contraceptives at extreme risk for venous thromboembolism? *European Journal of Contraception & Reproductive Health Care*, 2000, 5:105-12.

50. Vandenbroucke JP et al. Increased risk of venous thrombosis in oral-contraceptive users who are carriers of factor V Leiden mutation. *Lancet*, 1994, 344:1453-7.
51. Vaya AM. Prothrombin G20210A mutation and oral contraceptive use increase upper-extremity deep vein thrombotic risk. *Thrombosis & Haemostasis*, 2003, 89:452-7.
52. Carolei A, Marini C, De Matteis G. History of migraine and risk of cerebral ischaemia in young adults. The Italian National Research Council Study Group on Stroke in the Young. *Lancet*, 1996, 347:1503-6.
53. Chang CL, Donaghy M, Poulter N. Migraine and stroke in young women: case-control study. The World Health Organisation Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *BMJ*, 1999, 318:13-8.
54. Tzourio C et al. Case-control study of migraine and risk of ischaemic stroke in young women. *BMJ*, 1995, 310:830-3.
55. Donaghy M, Chang CL, Poulter N. Duration, frequency, recency, and type of migraine and the risk of ischaemic stroke in women of childbearing age. *Journal of Neurology, Neurosurgery, and Psychiatry*, 2002, 73:747-50.
56. Lidegaard O. Oral contraceptives, pregnancy and the risk of cerebral thromboembolism: the influence of diabetes, hypertension, migraine and previous thrombotic disease (Letter). *British Journal of Obstetrics & Gynaecology*, 1996, 103:94.
57. MacGregor EA, Guillebaud J. Combined oral contraceptives, migraine and ischaemic stroke. Clinical and Scientific Committee of the Faculty of Family Planning and Reproductive Health Care and the Family Planning Association. *British Journal of Family Planning*, 1998, 24:55-60.
58. Schwartz SM et al. Stroke and use of low-dose oral contraceptives in young women: a pooled analysis of two US studies. *Stroke*, 1998, 29:2277-84.
59. Cromer BA et al. A prospective study of adolescents who choose among levonorgestrel implant (Norplant), medroxyprogesterone acetate (Depo-Provera), or the combined oral contraceptive pill as contraception. *Pediatrics*, 1994, 94:687-94.
60. Deijen JB et al. Use of a monophasic, low-dose oral contraceptive in relation to mental functioning. *Contraception*, 1992, 46:359-67.
61. Herzberg BN et al. Oral contraceptives, depression, and libido. *BMJ*, 1971, 3:495-500.
62. Hendrix SL, Alexander NJ. Primary dysmenorrhea treatment with a desogestrel-containing low-dose oral contraceptive. *Contraception*, 2002, 66:393-9.
63. Proctor ML, Roberts H, Farquhar CM. Combined oral contraceptive pill (OCP) as treatment for primary dysmenorrhoea. *Cochrane Database of Systematic Reviews*, 2001, CD002120.
64. Berkowitz RS et al. Oral contraceptives and postmolar trophoblastic disease. *Obstetrics & Gynecology*, 1981, 58:474-7.
65. Curry SL et al. Hormonal contraception and trophoblastic sequelae after hydatidiform mole (a Gynecologic Oncology Group Study) *American Journal of Obstetrics & Gynecology*, 1989, 160:805-9.
66. Deicas RE et al. The role of contraception in the development of postmolar gestational trophoblastic tumor. *Obstetrics & Gynecology*, 1991, 78:221-6.
67. Eddy GL et al. Postmolar trophoblastic disease in women using hormonal contraception with and without estrogen. *Obstetrics & Gynecology*, 1983, 62:736-40.
68. Goldberg GL et al. Medroxyprogesterone acetate in non-metastatic gestational trophoblastic disease. *British Journal of Obstetrics & Gynaecology*, 1987, 94:22-5.
69. Morrow P et al. The influence of oral contraceptives on the postmolar human chorionic gonadotropin regression curve. *American Journal of Obstetrics & Gynecology*, 1985, 151:906-14.
70. Stone M et al. Relationship of oral contraception to development of trophoblastic tumour after evacuation of a hydatidiform mole. *British Journal of Obstetrics & Gynaecology*, 1976, 83:913-6.
71. Ho Yuen B, Burch P. Relationship of oral contraceptives and the intrauterine contraceptive devices to the regression of concentrations of the beta subunit of human chorionic gonadotropin and invasive complications after molar pregnancy. *American Journal of Obstetrics & Gynecology*, 1983, 145:214-7.
72. Smith JS et al. Cervical cancer and use of hormonal contraceptives: a systematic review. *Lancet*, 2003, 361:1159-67.
73. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53 297 women with breast cancer and 100 239 women without breast cancer from 54 epidemiological studies. Collaborative Group on Hormonal Factors in Breast Cancer. *Lancet*, 1996, 347:1713-27.
74. Brinton LA et al. Modification of oral contraceptive relationships on breast cancer risk by selected factors among younger women. *Contraception*, 1997, 55:197-203.

75. Colditz GA, Rosner BA, Speizer FE for the Nurses' Health Study Research Group. Risk factors for breast cancer according to family history of breast cancer. *Journal of the National Cancer Institute*, 1996, 88:365-71.
76. Egan KM et al. Risk factors for breast cancer in women with a breast cancer family history. *Cancer Epidemiology, Biomarkers & Prevention*, 1998, 7:359-64.
77. Grabrick DM et al. Risk of breast cancer with oral contraceptive use in women with a family history of breast cancer. *JAMA*, 2000, 284:1791-8.
78. Harris RE, Zang EA, Wynder EL. Oral contraceptives and breast cancer risk: a case-control study. *International Journal of Epidemiology*, 1990, 19:240-6.
79. Marchbanks PA et al. Oral contraceptives and the risk of breast cancer. *New England Journal of Medicine*, 2002, 346:2025-32.
80. Rosenberg L et al. Case-control study of oral contraceptive use and risk of breast cancer. *American Journal of Epidemiology*, 1996, 143:25-37.
81. Heimdal K, Skovlund E, Moller P. Oral contraceptives and risk of familial breast cancer. *Cancer Detection & Prevention*, 2002, 26:23-7.
82. Narod SA et al. Oral contraceptives and the risk of breast cancer in BRCA1 and BRCA2 mutation carriers. *Journal of the National Cancer Institute*, 2002, 94:1773-9.
83. Ursin G et al. Does oral contraceptive use increase the risk of breast cancer in women with BRCA1/BRCA2 mutations more than in other women? *Cancer Research*, 1997, 57:3678-81.
84. The Italian MEGIC Group. Determinants of cervical Chlamydia trachomatis infection in Italy. *Genitourinary Medicine*, 1993, 69:123-5.
85. Ackers JP et al. Antitrichomonal antibody in the vaginal secretions of women infected with *T. vaginalis*. *British Journal of Venereal Diseases*, 1975, 51:319-23.
86. Acosta-Cazares B, Ruiz-Maya L, Escobedo de la Pena J. Prevalence and risk factors for Chlamydia trachomatis infection in low-income rural and suburban populations of Mexico. *Sexually Transmitted Diseases*, 1996, 23:283-8.
87. Addiss DG et al. Selective screening for Chlamydia trachomatis infection in nonurban family planning clinics in Wisconsin. *Family Planning Perspectives*, 1987, 19:252-6.
88. Arya OP, Mallinson H, Goddard AD. Epidemiological and clinical correlates of chlamydial infection of the cervix. *British Journal of Venereal Diseases*, 1981, 57:118-24.
89. Austin H, Louv WC, Alexander WJ. A case-control study of spermicides and gonorrhea. *JAMA*, 1984, 251:2822-4.
90. Avonts D et al. Incidence of uncomplicated genital infections in women using oral contraception or an intrauterine device: a prospective study. *Sexually Transmitted Diseases*, 1990, 17:23-9.
91. Baeten JM et al. Hormonal contraception and risk of sexually transmitted disease acquisition: results from a prospective study. *American Journal of Obstetrics & Gynecology*, 2001, 185:380-5.
92. Barbone F et al. A follow-up study of methods of contraception, sexual activity, and rates of trichomoniasis, candidiasis, and bacterial vaginosis. *American Journal of Obstetrics & Gynecology*, 1990, 163:510-4.
93. Barnes RC et al. Quantitative culture of endocervical Chlamydia trachomatis. *Journal of Clinical Microbiology*, 1990, 28:774-80.
94. Berger GS, Keith L, Moss W. Prevalence of gonorrhoea among women using various methods of contraception. *British Journal of Venereal Diseases*, 1975, 51:307-9.
95. Bhattacharyya MN, Jephcott AE. Diagnosis of gonorrhoea in women--Influence of the contraceptive pill. *Journal of the American Venereal Disease Association*, 1976, 2:21-4.
96. Blum M, Pery J, Kitai E. The link between contraceptive methods and Chlamydia trachomatis infection. *Advances in Contraception*, 1988, 4:233-9.
97. Bontis J et al. Detection of Chlamydia trachomatis in asymptomatic women: relationship to history, contraception, and cervicitis. *Advances in Contraception*, 1994, 10:309-15.
98. Bramley M, Kinghorn G. Do oral contraceptives inhibit *Trichomonas vaginalis*? *Sexually Transmitted Diseases*, 1979, 6:261-3.
99. Bro F, Juul S. Predictors of Chlamydia trachomatis infection in women in general practice. *Family Practice*, 1990, 7:138-43.
100. Burns DC et al. Isolation of Chlamydia from women attending a clinic for sexually transmitted disease. *British Journal of Venereal Diseases*, 1975, 51:314-8.
101. Ceruti M et al. Methods of contraception and rates of genital infections. *Clinical & Experimental Obstetrics & Gynecology*, 1994, 21:119-23.
102. Chacko MR, Lovchik JC. Chlamydia trachomatis infection in sexually active adolescents: prevalence and risk factors. *Pediatrics*, 1984, 73:836-40.

103. Cottingham J, Hunter D. Chlamydia trachomatis and oral contraceptive use: a quantitative review. *Genitourinary Medicine*, 1992, 68:209-16.
104. Crowley T et al. Hormonal factors and the laboratory detection of Chlamydia trachomatis in women: implications for screening? *International Journal of STD & AIDS*, 1997, 8:25-31.
105. Edwards D, Phillips D, Stancombe S. Chlamydia trachomatis infection at a family planning clinic. *New Zealand Medical Journal*, 1985, 98:333-5.
106. Evans BA et al. Predictors of seropositivity to herpes simplex virus type 2 in women. *International Journal of STD & AIDS*, 2003, 14:30-6.
107. Evans DL et al. Detection of Chlamydia trachomatis in adolescent females using direct immunofluorescence. *Clinical Pediatrics*, 1988, 27:223-8.
108. Fish AN et al. Chlamydia trachomatis infection in a gynaecology clinic population: identification of high-risk groups and the value of contact tracing. *European Journal of Obstetrics, Gynecology, & Reproductive Biology*, 1989, 31:67-74.
109. Fouts AC, Kraus SJ. Trichomonas vaginalis: reevaluation of its clinical presentation and laboratory diagnosis. *Journal of Infectious Diseases*, 1980, 141:137-43.
110. Fraser JJ, Jr., Rettig PJ, Kaplan DW. Prevalence of cervical Chlamydia trachomatis and Neisseria gonorrhoeae in female adolescents. *Pediatrics*, 1983, 71:333-6.
111. Gertig DM et al. Risk factors for sexually transmitted diseases among women attending family planning clinics in Dar-es-Salaam, Tanzania. *Genitourinary Medicine*, 1997, 73:39-43.
112. Green J. Human papillomavirus infection and use of oral contraceptives. *British Journal of Cancer*, 2003, 88:1713-20.
113. Griffiths M, Hindley D. Gonococcal pelvic inflammatory disease, oral contraceptives, and cervical mucus. *Genitourinary Medicine*, 1985, 61:67.
114. Han Y et al. Risk profile for Chlamydia infection in women from public health clinics in New York State. *Journal of Community Health*, 1993, 18:1-9.
115. Handsfield HH et al. Criteria for selective screening for Chlamydia trachomatis infection in women attending family planning clinics. *JAMA*, 1986, 255:1730-4.
116. Hanna NF et al. The relation between vaginal pH and the microbiological status in vaginitis. *British Journal of Obstetrics & Gynaecology*, 1985, 92:1267-71.
117. Harrison HR et al. Cervical Chlamydia trachomatis infection in university women: relationship to history, contraception, ectopy, and cervicitis. *American Journal of Obstetrics & Gynecology*, 1985, 153:244-51.
118. Hart G. Factors associated with genital chlamydial and gonococcal infection in females. *Genitourinary Medicine*, 1992, 68:217-20.
119. Herrmann B et al. Genital chlamydial infection among women in Nicaragua: validity of direct fluorescent antibody testing, prevalence, risk factors and clinical manifestations. *Genitourinary Medicine*, 1996, 72:20-6.
120. Hewitt AB. Oral contraception among special clinic patients. With particular reference to the diagnosis of gonorrhoea. *British Journal of Venereal Diseases*, 1970, 46:106-7.
121. Hilton AL et al. Chlamydia A in the female genital tract. *British Journal of Venereal Diseases*, 1974, 50:1-10.
122. Hiltunen-Back E et al. A nationwide sentinel clinic survey of chlamydia trachomatis infection in Finland. *Sexually Transmitted Diseases*, 2001, 28:252-8.
123. Jacobson DL et al. Relationship of hormonal contraception and cervical ectopy as measured by computerized planimetry to chlamydial infection in adolescents. *Sexually Transmitted Diseases*, 2000, 27:313-9.
124. Jaffe LR et al. Chlamydia trachomatis detection in adolescents. A comparison of direct specimen and tissue culture methods. *Journal of Adolescent Health Care*, 1986, 7:401-4.
125. Jick H et al. Vaginal spermicides and gonorrhoea. *JAMA*, 1982, 248:1619-21.
126. Johannisson G, Karamustafa A, Brorson J. Influence of copper salts on gonococci. *British Journal of Venereal Diseases*, 1976, 52:176-7.
127. Keith L, Berer GS, Moss W. Cervical gonorrhoea in women using different methods of contraception. *Journal of the American Venereal Disease Association*, 1976, 3:17-9.
128. Kinghorn GR, Waugh MA. Oral contraceptive use and prevalence of infection with Chlamydia trachomatis in women. *British Journal of Venereal Diseases*, 1981, 57:187-90.
129. Lavreys L et al. Human herpesvirus 8: seroprevalence and correlates in prostitutes in Mombasa, Kenya. *Journal of Infectious Diseases*, 2003, 187:359-63.
130. Lefevre JC et al. Lower genital tract infections in women: comparison of clinical and epidemiologic findings with microbiology. *Sexually Transmitted Diseases*, 1988, 15:110-3.

131. Louv WC et al. Oral contraceptive use and the risk of chlamydial and gonococcal infections. *American Journal of Obstetrics & Gynecology*, 1989, 160:396-402.
132. Lowe TL, Kraus SJ. Quantitation of *Neisseria gonorrhoeae* from women with gonorrhoea. *Journal of Infectious Diseases*, 1976, 133:621-6.
133. Lycke E et al. The risk of transmission of genital *Chlamydia trachomatis* infection is less than that of genital *Neisseria gonorrhoeae* infection. *Sexually Transmitted Diseases*, 1980, 7:6-10.
134. Macaulay ME et al. A prospective study of genital infections in a family-planning clinic. 2. *Chlamydia* infection--the identification of a high-risk group. *Epidemiology & Infection*, 1990, 104:55-61.
135. Magder LS et al. Factors related to genital *Chlamydia trachomatis* and its diagnosis by culture in a sexually transmitted disease clinic. *American Journal of Epidemiology*, 1988, 128:298-308.
136. Magder LS et al. Effect of patient characteristics on performance of an enzyme immunoassay for detecting cervical *Chlamydia trachomatis* infection. *Journal of Clinical Microbiology*, 1990, 28:781-4.
137. Masse R et al. *Chlamydia trachomatis* cervical infection: prevalence and determinants among women presenting for routine gynecologic examination. *CMAJ Canadian Medical Association Journal*, 1991, 145:953-61.
138. McCormack WM, Reynolds GH. Effect of menstrual cycle and method of contraception on recovery of *Neisseria gonorrhoeae*. *JAMA*, 1982, 247:1292-4.
139. Nayyar KC et al. Isolation of *Chlamydia trachomatis* from women attending a clinic for sexually transmitted diseases. *British Journal of Venereal Diseases*, 1976, 52:396-8.
140. Oh MK et al. *Chlamydia trachomatis* cervical infection and oral contraceptive use among adolescent girls. *Journal of Adolescent Health Care*, 1989, 10:376-81.
141. Oriol JD et al. Infection of the uterine cervix with *Chlamydia trachomatis*. *Journal of Infectious Diseases*, 1978, 137:443-51.
142. Paavonen J, Vesterinen E. *Chlamydia trachomatis* in cervicitis and urethritis in women. *Scandinavian Journal of Infectious Diseases - Supplementum*, 1982, 32:45-54.
143. Park BJ et al. Contraceptive methods and the risk of *Chlamydia trachomatis* infection in young women. *American Journal of Epidemiology*, 1995, 142:771-8.
144. Pereira LH et al. Cytomegalovirus infection among women attending a sexually transmitted disease clinic: association with clinical symptoms and other sexually transmitted diseases. *American Journal of Epidemiology*, 1990, 131:683-92.
145. Reed BD, Huck W, Zazove P. Differentiation of *Gardnerella vaginalis*, *Candida albicans*, and *Trichomonas vaginalis* infections of the vagina. *Journal of Family Practice*, 1989, 28:673-80.
146. Ripa KT et al. *Chlamydia trachomatis* cervicitis in gynecologic outpatients. *Obstetrics & Gynecology*, 1978, 52:698-702.
147. Ruijs GJ et al. Direct immunofluorescence for *Chlamydia trachomatis* on urogenital smears for epidemiological purposes. *European Journal of Obstetrics, Gynecology & Reproductive Biology*, 1988, 27:289-97.
148. Schachter J, Stoner E, Moncada J. Screening for chlamydial infections in women attending family planning clinics. *Western Journal of Medicine*, 1983, 138:375-9.
149. Sellors JW et al. Incidence, clearance and predictors of human papillomavirus infection in women. *CMAJ Canadian Medical Association Journal*, 2003, 168:421-5.
150. Sessa R et al. Epidemiology of urogenital infections caused by *Chlamydia trachomatis* and outline of characteristic features of patients at risk. *Journal of Medical Microbiology*, 1994, 41:168-72.
151. Shafer MA et al. *Chlamydia trachomatis*: important relationships to race, contraception, lower genital tract infection, and Papanicolaou smear. *Journal of Pediatrics*, 1984, 104:141-6.
152. Smith JS et al. Prevalence and risk factors for herpes simplex virus type 2 infection among middle-age women in Brazil and the Philippines. *Sexually Transmitted Diseases*, 2001, 28:187-94.
153. Staerfelt F et al. A survey of genital infections in patients attending a clinic for sexually transmitted diseases. *Scandinavian Journal of Infectious Diseases - Supplementum*, 1983, 40:53-7.
154. Svensson L, Westrom L, Mardh PA. *Chlamydia trachomatis* in women attending a gynaecological outpatient clinic with lower genital tract infection. *British Journal of Venereal Diseases*, 1981, 57:259-62.
155. Tait IA et al. Chlamydial infection of the cervix in contacts of men with nongonococcal urethritis. *British Journal of Venereal Diseases*, 1980, 56:37-45.

156. Willmott FE, Mair HJ. Genital herpesvirus infection in women attending a venereal diseases clinic. *British Journal of Venereal Diseases*, 1978, 54:341-3.
157. Winer RL et al. Genital human papillomavirus infection: incidence and risk factors in a cohort of female university students.[erratum appears in Am J Epidemiol. 2003 May 1;157(9):858]. *American Journal of Epidemiology*, 2003, 157:218-26.
158. Winter L, Goldy AS, Baer C. Prevalence and epidemiologic correlates of Chlamydia trachomatis in rural and urban populations. *Sexually Transmitted Diseases*, 1990, 17:30-6.
159. Wolinska WH, Melamed MR. Herpes genitalis in women attending Planned Parenthood of New York City. *Acta Cytologica*, 1970, 14:239-42.
160. Woolfitt JM, Watt L. Chlamydial infection of the urogenital tract in promiscuous and non-promiscuous women. *British Journal of Venereal Diseases*, 1977, 53:93-5.
161. European Study Group on Heterosexual Transmission of HIV. Comparison of female to male and male to female transmission of HIV in 563 stable couples. *BMJ*, 1992, 304:809-13.
162. Akililu M et al. Factors associated with HIV-1 infection among sex workers of Addis Ababa, Ethiopia. *AIDS*, 2001, 15:87-96.
163. Allen S et al. Pregnancy and contraception use among urban Rwandan women after HIV testing and counseling. *American Journal of Public Health*, 1993, 83:705-10.
164. Carael M et al. Human immunodeficiency virus transmission among heterosexual couples in Central Africa. *AIDS*, 1988, 2:201-5.
165. Chao A et al. National University of Rwanda-Johns Hopkins University AIDS Research Team. Risk factors associated with prevalent HIV-1 infection among pregnant women in Rwanda. *International Journal of Epidemiology*, 1994, 23:371-80.
166. Cohen E, Navaline H, Metzger D. High-risk behaviors for HIV: a comparison between crack-abusing and opioid-abusing African-American women. *Journal of Psychoactive Drugs*, 1994, 26:233-41.
167. de Vincenzi I. European Study Group on Heterosexual Transmission of HIV. A longitudinal study of human immunodeficiency virus transmission by heterosexual partners. *New England Journal of Medicine*, 1994, 331:341-6.
168. Ellerbrock TV et al. Heterosexually transmitted human immunodeficiency virus infection among pregnant women in a rural Florida community. *New England Journal of Medicine*, 1992, 327:1704-9.
169. Gray JA et al. HIV-1 infection among female commercial sex workers in rural Thailand. *AIDS*, 1997, 11:89-94.
170. Guimaraes MD et al. Rio de Janeiro Heterosexual Study Group. HIV infection among female partners of seropositive men in Brazil. *American Journal of Epidemiology*, 1995, 142:538-47.
171. Hira SK et al. Oral contraceptive use and HIV infection. *International Journal of STD & AIDS*, 1990, 1:447-8.
172. Kapiga SH et al. Risk factors for HIV infection among women in Dar-es-Salaam, Tanzania. *Journal of Acquired Immune Deficiency Syndromes*, 1994, 7:301-9.
173. Kapiga SH et al. The incidence of HIV infection among women using family planning methods in Dar es Salaam, Tanzania. *AIDS*, 1998, 12:75-84.
174. Kiddugavu M et al. Hormonal contraceptive use and HIV-1 infection in a population-based cohort in Rakai, Uganda. *AIDS*, 2003, 17:233-40.
175. Kunanusont C et al. HIV-1 subtypes and male-to-female transmission in Thailand. *Lancet*, 1995, 345:1078-83.
176. Laga M et al. Non-ulcerative sexually transmitted diseases as risk factors for HIV-1 transmission in women: results from a cohort study. *AIDS*, 1993, 7:95-102.
177. Latif AS et al. Genital ulcers and transmission of HIV among couples in Zimbabwe. *AIDS*, 1989, 3:519-23.
178. Limpakarnjanarat K et al. HIV-1 and other sexually transmitted infections in a cohort of female sex workers in Chiang Rai, Thailand. *Sexually Transmitted Infections*, 1999, 75:30-5.
179. Martin HL, Jr. et al. Hormonal contraception, sexually transmitted diseases, and risk of heterosexual transmission of human immunodeficiency virus type 1. *Journal of Infectious Diseases*, 1998, 178:1053-9.
180. Mati JK et al. Contraceptive use and the risk of HIV infection in Nairobi, Kenya. *International Journal of Gynaecology & Obstetrics*, 1995, 48:61-7.
181. Moss GB et al. Association of cervical ectopy with heterosexual transmission of human immunodeficiency virus: results of a study of couples in Nairobi, Kenya. *Journal of Infectious Diseases*, 1991, 164:588-91.

182. Nagachinta T et al. Risk factors for HIV-1 transmission from HIV-seropositive male blood donors to their regular female partners in northern Thailand. *AIDS*, 1997, 11:1765-72.
183. Nicolosi A et al. Italian Study Group on HIV Heterosexual Transmission. The efficiency of male-to-female and female-to-male sexual transmission of the human immunodeficiency virus: a study of 730 stable couples. *Epidemiology*, 1994, 5:570-5.
184. Nzila N et al. HIV and other sexually transmitted diseases among female prostitutes in Kinshasa. *AIDS*, 1991, 5:715-21.
185. Pineda JA et al. HIV-1 infection among non-intravenous drug user female prostitutes in Spain. No evidence of evolution to pattern II. *AIDS*, 1992, 6:1365-9.
186. Plourde PJ et al. Human immunodeficiency virus type 1 infection in women attending a sexually transmitted diseases clinic in Kenya. *Journal of Infectious Diseases*, 1992, 166:86-92.
187. Plummer FA et al. Cofactors in male-female sexual transmission of human immunodeficiency virus type 1. *Journal of Infectious Diseases*, 1991, 163:233-9.
188. Rehle T et al. Risk factors of HIV-1 infection among female prostitutes in Khon Kaen, Northeast Thailand. *Infection*, 1992, 20:328-31.
189. Saracco A et al. Man-to-woman sexual transmission of HIV: longitudinal study of 343 steady partners of infected men. *Journal of Acquired Immune Deficiency Syndromes*, 1993, 6:497-502.
190. Simonsen JN et al. HIV infection among lower socioeconomic strata prostitutes in Nairobi. *AIDS*, 1990, 4:139-44.
191. Sinei SK et al. Contraceptive use and HIV infection in Kenyan family planning clinic attenders. *International Journal of STD & AIDS*, 1996, 7:65-70.
192. Siraprasasiri T et al. Risk factors for HIV among prostitutes in Chiangmai, Thailand. *AIDS*, 1991, 5:579-82.
193. Spence MR et al. Seroprevalence of human immunodeficiency virus type I (HIV-1) antibodies in a family-planning population. *Sexually Transmitted Diseases*, 1991, 18:143-5.
194. Stephenson JM. Systematic review of hormonal contraception and risk of HIV transmission: when to resist meta-analysis. *AIDS*, 1998, 12:545-53.
195. Taneepanichskul S, Phuapradit W, Chaturachinda K. Association of contraceptives and HIV-1 infection in Thai female commercial sex workers. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 1997, 37:86-8.
196. Temmerman M et al. Maternal human immunodeficiency virus-1 infection and pregnancy outcome. *Obstetrics & Gynecology*, 1994, 83:495-501.
197. Ungchusak K et al. Determinants of HIV infection among female commercial sex workers in northeastern Thailand: results from a longitudinal study.[erratum appears in J Acquir Immune Defic Syndr Hum Retrovirol 1998 Jun 1;18(2):192]. *Journal of Acquired Immune Deficiency Syndromes & Human Retrovirology*, 1996, 12:500-7.
198. Wang CC, Reilly M, Kreiss JK. Risk of HIV infection in oral contraceptive pill users: a meta-analysis.[erratum appears in J Acquir Immune Defic Syndr 1999 Aug 15;21(5):428]. *Journal of Acquired Immune Deficiency Syndromes: JAIDS*, 1999, 21:51-8.
199. Cejtin HEJ. Effect of hormonal contraceptive use on plasma HIV-1-RNA levels among HIV-infected women. *AIDS*, 2003, 17:1702-4.
200. Clemetson DB et al. Detection of HIV DNA in cervical and vaginal secretions. Prevalence and correlates among women in Nairobi, Kenya. *JAMA*, 1993, 269:2860-4.
201. Kreiss J et al. Association between cervical inflammation and cervical shedding of human immunodeficiency virus DNA. *Journal of Infectious Diseases*, 1994, 170:1597-601.
202. McClelland RS et al. A prospective study of hormonal contraceptive use and cervical shedding of herpes simplex virus in human immunodeficiency virus type 1-seropositive women. *Journal of Infectious Diseases*, 2002, 185:1822-5.
203. Mostad SB et al. Hormonal contraception, vitamin A deficiency, and other risk factors for shedding of HIV-1 infected cells from the cervix and vagina. *Lancet*, 1997, 350:922-7.
204. Mostad SB et al. Cervical shedding of herpes simplex virus in human immunodeficiency virus-infected women: effects of hormonal contraception, pregnancy, and vitamin A deficiency. *Journal of Infectious Diseases*, 2000, 181:58-63.
205. el Raghy I et al. Contraceptive steroid concentrations in women with early active schistosomiasis: lack of effect of antischistosomal drugs. *Contraception*, 1986, 33:373-7.
206. Gad-el-Mawla N, Abdallah A. Liver function in bilharzial females receiving contraceptive pills. *Acta Hepato-Splenologica*, 1969, 16:308-10.
207. Gad-el-Mawla N et al. Plasma lipids and lipoproteins in bilharzial females during oral contraceptive therapy. *Journal of the Egyptian Medical Association*, 1972, 55:137-47.

208. Shaaban MM et al. Effects of oral contraception on liver function tests and serum proteins in women with active schistosomiasis. *Contraception*, 1982, 26:75-82.
209. Shaaban MM et al. Effect of oral contraception on serum bile acid. *International Journal of Gynaecology & Obstetrics*, 1984, 22:111-5.
210. Sy FS et al. Effect of oral contraceptive on liver function tests of women with schistosomiasis in the Philippines. *Contraception*, 1986, 34:283-94.
211. Tagy AH et al. The effect of low-dose combined oral contraceptive pills versus injectable contraceptive (Depot Provera) on liver function tests of women with compensated bilharzial liver fibrosis. *Contraception*, 2001, 64:173-6.
212. Back DJ et al. The effect of rifampicin on norethisterone pharmacokinetics. *European Journal of Clinical Pharmacology*, 1979, 15:193-7.
213. Back DJ et al. The effect of rifampicin on the pharmacokinetics of ethinylestradiol in women. *Contraception*, 1980, 21:135-43.
214. Barditch-Crovo P et al. The effects of rifampin and rifabutin on the pharmacokinetics and pharmacodynamics of a combination oral contraceptive. *Clinical Pharmacology & Therapeutics*, 1999, 65:428-38.
215. Bessot JC et al. [Rifampicin interference with oral contraceptives]. *Journal de Médecine de Strasbourg*, 1977, 8:131-3.
216. Bolt HM, Bolt M, Kappus H. Interaction of rifampicin treatment with pharmacokinetics and metabolism of ethinyloestradiol in man. *Acta Endocrinologica*, 1977, 85:189-97.
217. Gelbke HP, Gethmann U, Knuppen R. Influence of rifampicin treatment on the metabolic fate of [4-¹⁴C] mestranol in women. *Hormone and Metabolic Research*, 1977, 9:415-9.
218. Gupta KC, Ali MY. Failure of oral contraceptive with rifampicin. *Medical Journal of Zambia*, 1980, 15:23.
219. Hirsch A. [Letter: Sleeping pills]. [French]. *Nouvelle Presse Médicale*, 1973, 2:2957.
220. Hirsch A, Tillement JP, Chretien J. Effets contrariais de la rifampicine sur les contraceptifs oraux: à propos de trois grossesses non désirées chez trois malades. *Revue Française des Maladies Respiratoires*, 1975, 2:174-82.
221. Joshi JV et al. A study of interaction of a low-dose combination oral contraceptive with anti-tubercular drugs. *Contraception*, 1980, 21:617-29.
222. Kropp R. [Rifampicin and oral contraceptives (author's transl)]. [German]. *Praxis der Pneumologie vereinigt mit der Tuberkulosearzt*, 1974, 28:270-2.
223. LeBel M et al. Effects of rifabutin and rifampicin on the pharmacokinetics of ethinylestradiol and norethindrone. *Journal of Clinical Pharmacology*, 1998, 38:1042-50.
224. Meyer B et al. A model to detect interactions between roxithromycin and oral contraceptives. *Clinical Pharmacology & Therapeutics*, 1990, 47:671-4.
225. Nocke-Finck L, Breuer H, Reimers D. [Effects of rifampicin on the menstrual cycle and on oestrogen excretion in patients taking oral contraceptives]. [German]. *Deutsche Medizinische Wochenschrift*, 1973, 98:1521-3.
226. Piguet B, Muglioni JF, Chaline G. [Letter: Oral contraception and rifampicin]. [French]. *Nouvelle Presse Médicale*, 1975, 4:115-6.
227. Reimers D, Jezek A. [The simultaneous use of rifampicin and other antitubercular agents with oral contraceptives]. [German]. *Praxis der Pneumologie vereinigt mit der Tuberkulosearzt*, 1971, 25:255-62.
228. Skolnick JL et al. Rifampin, oral contraceptives, and pregnancy. *JAMA*, 1976, 236:1382.
229. Szoka PR, Edgren RA. Drug interactions with oral contraceptives: compilation and analysis of an adverse experience report database. *Fertility & Sterility*, 1988, 49:31S-8S.
230. Back DJ et al. The interaction of phenobarbital and other anticonvulsants with oral contraceptive steroid therapy. *Contraception*, 1980, 22:495-503.
231. Bartoli A et al. A double-blind, placebo-controlled study on the effect of vigabatrin on in vivo parameters of hepatic microsomal enzyme induction and on the kinetics of steroid oral contraceptives in healthy female volunteers. *Epilepsia*, 1997, 38:702-7.
232. Crawford P et al. The lack of effect of sodium valproate on the pharmacokinetics of oral contraceptive steroids. *Contraception*, 1986, 33:23-9.
233. Eldon MA et al. Gabapentin does not interact with a contraceptive regimen of norethindrone acetate and ethinyl estradiol. *Neurology*, 1998, 50:1146-8.
234. Fattore C et al. Induction of ethinylestradiol and levonorgestrel metabolism by oxcarbazepine in healthy women. *Epilepsia*, 1999, 40:783-7.
235. Ragueneau-Majlessi I, Levy RH, Janik F. Levetiracetam does not alter the pharmacokinetics of an oral contraceptive in healthy women. *Epilepsia*, 2002, 43:697-702.

236. Rosenfeld WE et al. Effect of topiramate on the pharmacokinetics of an oral contraceptive containing norethindrone and ethinyl estradiol in patients with epilepsy. *Epilepsia*, 1997, 38:317-23.
237. Sabers A et al. Lamotrigine plasma levels reduced by oral contraceptives. *Epilepsy Research*, 2001, 47:151-4.
238. Back DJ et al. The effects of ampicillin oral contraceptive steroids in women. *British Journal of Clinical Pharmacology*, 1982, 14:43-8.
239. Back DJ et al. Pharmacokinetics of oral contraceptive steroids following the administration of the antimalarial drugs primaquine and chloroquine. *Contraception*, 1984, 30:289-95.
240. Back DJ et al. Evaluation of Committee on Safety of Medicines yellow card reports on oral contraceptive-drug interactions with anticonvulsants and antibiotics. *British Journal of Clinical Pharmacology*, 1988, 25:527-32.
241. Back DJ et al. The lack of interaction between temafloxacin and combined oral contraceptive steroids. *Contraception*, 1991, 43:317-23.
242. Bacon JF, Shenfield GM. Pregnancy attributable to interaction between tetracycline and oral contraceptives. *BMJ*, 1980, 280:293.
243. Bainton R. Interaction between antibiotic therapy and contraceptive medication. *Oral Surgery, Oral Medicine, Oral Pathology*, 1986, 61:453-5.
244. Biron CR. Questions surface over whether antibiotics neutralized 'the pill,' resulting in pregnancy. *RDH*, 1996, 16:34-6.
245. Bollen M. Use of antibiotics when taking the oral contraceptive pill. *Australian Family Physician*, 1995, 24:928-9.
246. Bromham DR, Cartmill RS. Knowledge and use of secondary contraception among patients requesting termination of pregnancy. *BMJ*, 1993, 306:556-7.
247. Cote J. Interaction of griseofulvin and oral contraceptives. *Journal of the American Academy of Dermatology*, 1990, 22:124-5.
248. Csemiczky G, Alvendal C, Landgren BM. Risk for ovulation in women taking a low-dose oral contraceptive (Microgynon) when receiving antibacterial treatment with a fluoroquinolone (ofloxacin). *Advances in Contraception*, 1996, 12:101-9.
249. de Groot AC, Eshuis H, Stricker BH. [Inefficacy of oral contraception during use of minocycline]. *Nederlands Tijdschrift voor Geneeskunde*, 1990, 134:1227-9.
250. DeSano EA, Hurley SC. Possible interactions of antihistamines and antibiotics with oral contraceptive effectiveness. *Fertility & Sterility*, 1982, 37:853-4.
251. Devenport MH et al. Metabolic effects of low-dose fluconazole in healthy female users and non-users of oral contraceptives. *British Journal of Clinical Pharmacology*, 1989, 27:851-9.
252. Donley TG, Smith RF, Roy B. Reduced oral contraceptive effectiveness with concurrent antibiotic use: a protocol for prescribing antibiotics to women of childbearing age. *Compendium*, 1990, 11:392-6.
253. Dosseter J. Drug interaction with oral contraceptives. *BMJ*, 1984, 4:467-8.
254. Driver J. Use of antibiotics during pregnancy. *British Dental Journal*, 1982, 153:48.
255. el-Raghy I et al. Contraceptive steroid concentrations in women with early active schistosomiasis: lack of effect of antischistosomal drugs. *Contraception*, 1986, 33:373-7.
256. Friedman CI et al. The effect of ampicillin on oral contraceptive effectiveness. *Obstetrics & Gynecology*, 1980, 55:33-7.
257. Grimmer SF et al. The effect of cotrimoxazole on oral contraceptive steroids in women. *Contraception*, 1983, 28:53-9.
258. Helms SE et al. Oral contraceptive failure rates and oral antibiotics. *Journal of the American Academy of Dermatology*, 1997, 36:705-10.
259. Hempel E et al. [Enzyme induction by drugs and hormonal contraception]. [German]. *Zentralblatt für Gynäkologie*, 1973, 95:1451-7.
260. Hempel E, Zorn C, Graf K. [Effect of chemotherapy agents and antibiotics on hormonal contraception]. [German]. *Zeitschrift für Ärztliche Fortbildung (Jena)*, 1978, 72:924-6.
261. Hetenyi G. Possible interactions between antibiotics and oral contraceptives. *Therapia Hungarica*, 1989, 37:86-9.
262. Hilbert J et al. Evaluation of interaction between fluconazole and an oral contraceptive in healthy women. *Obstetrics & Gynecology*, 2001, 98:218-23.
263. Hughes BR, Cunliffe WJ. Interactions between the oral contraceptive pill and antibiotics. *British Journal of Dermatology*, 1990, 122:717-8.
264. Joshi JV et al. A study of interaction of low-dose combination oral contraceptive with ampicillin and metronidazole. *Contraception*, 1980, 22:643-52.

265. Kakouris H, Kovacs GT. How common are predisposing factors to pill failure among pill users? *British Journal of Family Planning*, 1994, 20:33-5.
266. Kakouris H, Kovacs GT. Pill failure and nonuse of secondary precautions. *British Journal of Family Planning*, 1992, 18:41-4.
267. Kovacs GT et al. Inadvertent pregnancies in oral contraceptive users. *Medical Journal of Australia*, 1989, 150:549-51.
268. Kovacs I, Somos P, Hamori M. Examination of the potential interaction between ketoconazole (Nizoral) and oral contraceptives with special regard to products of low hormone content (Rigevidon, Anteovin). *Therapia Hungarica*, 1986, 34:167-70.
269. Lequeux A. [Pregnancy under oral contraceptives after treatment with tetracycline]. *Louvain Médical*, 1980, 99:413-4.
270. London BM, Lookingbill DP. Frequency of pregnancy in acne patients taking oral antibiotics and oral contraceptives. *Archives of Dermatology*, 1994, 130:392-3.
271. Lunell NO et al. Evaluation of the possible interaction of the antifungal triazole SCH 39304 with oral contraceptives in normal healthy women. *Gynecologic & Obstetric Investigation*, 1991, 32:91-7.
272. Maggiolo F et al. The effects of ciprofloxacin on oral contraceptive steroid treatments. *Drugs under Experimental and Clinical Research*, 1991, 17:451-4.
273. McDaniel PA, Caldrony RD. Oral contraceptives and griseofulvin interactions. *Drug Intelligence & Clinical Pharmacy*, 1986, 20:384.
274. Meyboom RH et al. Disturbance of withdrawal bleeding during concomitant use of itraconazole and oral contraceptives. *New Zealand Medical Journal*, 1997, 110:300.
275. Murphy AA et al. The effect of tetracycline on levels of oral contraceptives. *American Journal of Obstetrics and Gynecology*, 1991, 164:28-33.
276. Neely JL et al. The effect of doxycycline on serum levels of ethinyl estradiol, norethindrone, and endogenous progesterone. *Obstetrics & Gynecology*, 1991, 77:416-20.
277. Organon. Nuvaring prescribing information. 2001.
278. Pillans PI, Sparrow MJ. Pregnancy associated with a combined oral contraceptive and itraconazole. *New Zealand Medical Journal*, 1993, 106:436.
279. Rieth H, Sauerbrey N. [Interaction studies with fluconazole, a new triazole antifungal drug]. [German]. *Wiener Medizinische Wochenschrift*, 1989, 139:370-4.
280. Scholten PC et al. No interaction between ciprofloxacin and an oral contraceptive. *Antimicrobial Agents & Chemotherapy*, 1998, 42:3266-8.
281. Silber TJ. Apparent oral contraceptive failure associated with antibiotic administration. *Journal of Adolescent Health Care*, 1983, 4:287-9.
282. Sinofsky FE, Pasquale SA. The effect of fluconazole on circulating ethinyl estradiol levels in women taking oral contraceptives. *American Journal of Obstetrics & Gynecology*, 1998, 178:300-4.
283. Sparrow MJ. Pill method failures. *New Zealand Medical Journal*, 1987, 100:102-5.
284. Sparrow MJ. Pregnancies in reliable pill takers. *New Zealand Medical Journal*, 1989, 102:575-7.
285. Sparrow MJ. Pill method failures in women seeking abortion: fourteen years experience. *New Zealand Medical Journal*, 1998, 111:386-8.
286. van Dijke CP, Weber JC. Interaction between oral contraceptives and griseofulvin. *British Medical Journal Clinical Research Edition*, 1984, 288:1125-6.
287. van Puijenbroek EP, Feenstra J, Meyboom RH. [Pill cycle disturbance in simultaneous use of itraconazole and oral contraceptives]. [Dutch]. *Nederlands Tijdschrift voor Geneeskunde*, 1998, 142:146-9.
288. van Puijenbroek EP et al. Signalling possible drug-drug interactions in a spontaneous reporting system: delay of withdrawal bleeding during concomitant use of oral contraceptives and itraconazole. *British Journal of Clinical Pharmacology*, 1999, 47:689-93.
289. Wermeling DP et al. Dirithromycin increases ethinyl estradiol clearance without allowing ovulation. *Obstetrics & Gynecology*, 1995, 86:78-84.
290. Young LK et al. The contraceptive practices of women seeking termination of pregnancy in an Auckland clinic. *New Zealand Medical Journal*, 1994, 107:189-92.

Table of contents Combined injectable contraceptives, patch & ring

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	3
Pregnancy.....	3
Age	3
Parity.....	3
Breastfeeding.....	3
Postpartum	3
Post-abortion.....	3
Past ectopic pregnancy.....	4
History of pelvic surgery.....	4
Smoking.....	4
Obesity.....	4
Blood pressure measurement unavailable	4
CARDIOVASCULAR DISEASE.....	4
Multiple risk factors for arterial cardiovascular disease.....	4
Hypertension.....	5
History of high blood pressure during pregnancy	5
Deep venous thrombosis (DVT)/Pulmonary embolism (PE)	6
Known thrombogenic mutations.....	6
Superficial venous thrombosis	6
Current and history of ischaemic heart disease	6
Stroke	6
Known hyperlipidaemias	7
Valvular heart disease	7
NEUROLOGIC CONDITIONS.....	7
Headaches.....	7
Epilepsy	7
DEPRESSIVE DISORDERS	8
Depressive disorders	8
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS.....	8
Vaginal bleeding patterns.....	8
Unexplained vaginal bleeding	8
Endometriosis	8
Benign ovarian tumours	8
Severe dysmenorrhoea.....	8
Trophoblast disease.....	8
Cervical ectropion	8
Cervical intraepithelial neoplasia (CIN)	9
Cervical cancer	9
Breast disease	9
Endometrial cancer	9
Ovarian cancer.....	9
Uterine fibroids.....	9
Pelvic inflammatory disease (PID)	9
STIs	10
HIV/AIDS	10
High risk of HIV	10
HIV-infected	10
AIDS	10

OTHER INFECTIONS	10
Schistosomiasis	10
Tuberculosis.....	10
Malaria	10
ENDOCRINE CONDITIONS.....	11
Diabetes.....	11
Thyroid disorders	11
GASTROINTESTINAL CONDITIONS.....	11
Gall-bladder disease	11
History of cholestasis.....	11
Viral hepatitis	12
Cirrhosis.....	12
Liver tumours	12
ANAEMIAS.....	12
Thalassaemia.....	12
Sickle cell disease.....	12
Iron-deficiency anaemia	12
DRUG INTERACTIONS	12
Drugs which affect liver enzymes.....	12
Antibiotics	12
Antiretroviral therapy.....	13
Additional comments	14
References for combined injectable contraceptives, patch, and ring	17

COMBINED INJECTABLE CONTRACEPTIVES (CICs)

Combined injectable contraceptives (CICs) provide for the release of a natural estrogen plus a progestogen and act through the inhibition of ovulation.¹⁻⁵ Two CIC formulations, both given at four-week intervals, are considered here:

- 1) **Cyclofem** = Medroxyprogesterone acetate 25mg plus estradiol cypionate 5mg
- 2) **Mesigyna** = Norethisterone enantate 50mg plus estradiol valerate 5mg

Because the estrogens in CICs may be more physiologic and may be less potent compared with the synthetic estrogens of combined oral contraceptives (COCs), the type and magnitude of estrogen-related side-effects associated with CICs may be different from those experienced by COC users. In fact, short-term studies of CICs have shown little effect on blood pressure, haemostasis and coagulation, lipid metabolism, and liver function in comparison with COCs.⁶⁻⁸ In addition, the parenteral administration of CICs eliminates the first-pass effect of the hormones on the liver.

However, CICs are a relatively new contraceptive method, and there are few epidemiological data on their long-term effects. There is also the concern that, while the effect of the hormonal exposure associated with use of COCs and progestogen-only pills (POPs) can be reduced immediately by discontinuing their use, this is not the case with injectables, for which the effect continues for some time after the last injection.

Pending further evidence, the Working Group concluded that the evidence available for COCs applies to CICs in many but not all instances. Therefore, the Working Group assigned categories for CICs somewhere between the categories for COCs and POPs. However, for severe pathologies (e.g., ischaemic heart disease), the classification of conditions was the same as for COCs. The assigned categories should, therefore, be considered a preliminary, best judgement, which will be re-evaluated as new data become available.

COMBINED CONTRACEPTIVE PATCH (P)

The combined contraceptive patch uses a square 20 cm², three-layer system applied to the buttocks, torso, abdomen, or upper arm, to release ethinylestradiol and a progestogen (norelgestromin) transdermally. The contraceptive effect of the combined patch is achieved through inhibition of ovulation.⁹ The combined contraceptive patch currently available for consideration was:

Evra = 17-deacetyl norgestimate (norelgestromin) 150µg plus ethinylestradiol 20µg (both dosages are approximate daily release rates).

The combined contraceptive patch is a new contraceptive method. Relatively limited information is available on the safety of the combined contraceptive patch among healthy women and even less information is available for women with specific medical conditions. Moreover, epidemiological data on the long-term effects of the combined contraceptive patch were not available for the Working Group to review and all available studies received support from the patch manufacturer.

According to available evidence, the combined contraceptive patch provides a comparable safety and pharmacokinetic profile to COCs with similar hormone formulations.⁹⁻¹⁸ Reports of transient, short-term breast discomfort and skin site reactions occurred among less than 25% of combined contraceptive patch users.¹⁰⁻¹³ Limited evidence suggests the effectiveness of the patch may decline for women weighing 90kg or more.¹⁰⁻¹¹ To date, no studies have examined whether the avoidance of the first-pass effect of hormones on the liver with patch use lessens concerns about drug interactions or use of the patch among women with liver conditions.

Pending further evidence, the Working Group concluded that the evidence available for COCs applies to the patch. Therefore, the patch should have the same categories as COCs. The assigned categories should, therefore, be considered a preliminary, best judgement, which will be re-evaluated as new data become available.

COMBINED VAGINAL RING (R)

The combined contraceptive vaginal ring releases ethinylestradiol and a progestogen (etonogestrel) from a 54mm ethylene vinyl acetate copolymer ring. The contraceptive effect of the combined vaginal ring is achieved through inhibition of ovulation.¹⁹⁻²⁰ The vaginal ring formulation currently available for consideration was: **NuvaRing** = etonogestrel 120 µg plus ethinylestradiol 15 µg (both dosages are approximate daily release rates).

The combined contraceptive vaginal ring is a new contraceptive method. Relatively limited information is available on the safety of the combined contraceptive ring among healthy women and even less information is available for women with specific medical conditions. Moreover, epidemiological data on the long-term effects of the combined contraceptive ring were not available for the Working Group to review and all available studies received support from the ring manufacturer.

According to available evidence, the combined contraceptive vaginal ring provides a comparable safety and pharmacokinetic profile to COCs with similar hormone formulations.²⁰⁻²⁵ Evidence among healthy women suggests the vaginal ring does not alter vaginal flora,²³⁻²⁴ and limited evidence on women with low-grade squamous intraepithelial lesions found use of the vaginal ring did not worsen the condition.²³ To date, no studies have examined whether the avoidance of the first-pass effect of hormones on the liver with vaginal ring use lessens concerns about drug interactions or use of the ring among women with liver conditions.

Pending further evidence, the Working Group concluded that the evidence available for COCs applies to the ring. Therefore, the ring should have the same categories as COCs. The assigned categories should, therefore, be considered a preliminary, best judgement, which will be re-evaluated as new data become available.

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY				
PREGNANCY	NA	NA	NA	Clarification: Use of CICs, P, or R is not required. There is no known harm to the woman, the course of her pregnancy, or the fetus if CICs, P, or R are accidentally used during pregnancy.
AGE*				
a) Menarche to < 40 years	1	1	1	
b) ≥ 40 years	2	2	2	
PARITY				
a) Nulliparous	1	1	1	
b) Parous	1	1	1	
BREASTFEEDING*				
a) < 6 weeks postpartum	4	4	4	
b) ≥ 6 weeks to < 6 months postpartum (primarily breastfeeding)	3	3	3	
c) ≥ 6 months postpartum	2	2	2	
POSTPARTUM* (in non-breastfeeding women)				
a) < 21 days	3	3	3	
b) ≥ 21 days	1	1	1	
POST-ABORTION				
a) First trimester	1	1	1	Clarification: CICs, P, or R may be started immediately post-abortion.
b) Second trimester	1	1	1	
c) Immediate post-septic abortion	1	1	1	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
PAST ECTOPIC PREGNANCY*	1	1	1	
HISTORY OF PELVIC SURGERY	1	1	1	
SMOKING				
a) Age < 35 years	2	2	2	
b) Age ≥ 35 years				
(i) <15 cigarettes/day	2	3	3	
(ii) ≥15 cigarettes/day	3	4	4	
OBESITY ≥ 30 kg/m ² body mass index (BMI)	2	2	2	Evidence: Limited evidence suggests the effectiveness of the patch may decline for women weighing 90kg or more. ¹⁰⁻¹¹
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA	NA	NA	Clarification: It is desirable to have blood pressure measurements taken before initiation of CIC, P, or R use. However, in some settings blood pressure measurements are unavailable. In many of these settings, pregnancy morbidity and mortality risks are high, and CICs, P, or R may be one of the few methods available. In such settings, women should not be denied use of CICs, P, or R simply because their blood pressure cannot be measured.
CARDIOVASCULAR DISEASE				
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	3/4	3/4	3/4	Clarification: When a woman has multiple major risk factors, any of which alone would substantially increase the risk of cardiovascular disease, use of CICs, P, or R may increase her risk to an unacceptable level. However, a simple addition of categories for multiple risk factors is not intended; for example, a combination of two risk factors assigned a category 2 may not necessarily warrant a higher category.

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
HYPERTENSION*				
For all categories of hypertension, classifications are based on the assumption that no other risk factors for cardiovascular disease exist. When multiple risk factors do exist, risk of cardiovascular disease may increase substantially. A single reading of blood pressure level is not sufficient to classify a woman as hypertensive.				
a) History of hypertension, where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	3	3	3	Clarification: Evaluation of cause and level of hypertension is recommended as soon as feasible. Clarification: Women adequately treated for hypertension are at reduced risk of acute myocardial infarction and stroke as compared with untreated women. Although there are no data, CIC, P, or R users with adequately controlled and monitored hypertension should be at reduced risk of acute myocardial infarction and stroke compared with untreated hypertensive CIC, P, or R users.
b) Adequately controlled hypertension, where blood pressure CAN be evaluated	3	3	3	
c) Elevated blood pressure levels (properly taken measurements)				
(i) systolic 140-159 or diastolic 90-99	3	3	3	
(ii) systolic \geq 160 or diastolic \geq 100	4	4	4	
d) Vascular disease	4	4	4	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	2	2	2	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
DEEP VENOUS THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)* a) History of DVT/PE b) Current DVT/PE c) Family history of DVT/PE (first-degree relatives) d) Major surgery (i) with prolonged immobilization (ii) without prolonged immobilization e) Minor surgery without immobilization	 4 4 2 4 2 1	 4 4 2 4 2 1	 4 4 2 4 2 1	
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	4	4	4	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
SUPERFICIAL VENOUS THROMBOSIS* a) Varicose veins b) Superficial thrombophlebitis	 1 2	 1 2	 1 2	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE*	4	4	4	
STROKE* (history of cerebrovascular accident)	4	4	4	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.					
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE		
	CICs	P	R			
KNOWN HYPERLIPIDAEMIAS	2/3	2/3	2/3	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening. While some types of hyperlipidaemias are risk factors for vascular disease, the category should be assessed according to the type, its severity, and the presence of other cardiovascular risk factors.		
VALVULAR HEART DISEASE*						
a) Uncomplicated	2	2	2			
b) Complicated (pulmonary hypertension, risk of atrial fibrillation, history of subacute bacterial endocarditis)	4	4	4			
NEUROLOGIC CONDITIONS						
HEADACHES*	I	C	I	C	I	C
a) Non-migrainous (mild or severe)	1	2	1	2	1	2
b) Migraine						
(i) without aura						
Age < 35	2	3	2	3	2	3
Age ≥ 35	3	4	3	4	3	4
(ii) with aura, at any age	4	4	4	4	4	4
	Clarification: Classification depends on accurate diagnosis of those severe headaches that are migrainous and those that are not. Any new headaches or marked changes in headaches should be evaluated. Classification is for women without any other risk factors for stroke. Risk of stroke increases with age, hypertension and smoking.					
EPILEPSY	1	1	1	Clarification: If a woman is taking anticonvulsants, refer to the section on drug interactions. Certain anticonvulsants lower COC effectiveness. The extent to which CIC, P or R use are similar to COC use in this regard remains unclear.		

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
DEPRESSIVE DISORDERS				
DEPRESSIVE DISORDERS	1	1	1	Clarification: The classification is based on data for women with selected depressive disorders. No data on bipolar disorder or postpartum depression were available. There is a potential for drug interactions between certain antidepressant medications and hormonal contraceptives.
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS				
VAGINAL BLEEDING PATTERNS*				
a) Irregular pattern <i>without</i> heavy bleeding	1	1	1	
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	1	1	1	Clarification: Unusually heavy bleeding should raise the suspicion of a serious underlying condition.
UNEXPLAINED VAGINAL BLEEDING* (suspicious for serious condition)				
Before evaluation	2	2	2	Clarification: If pregnancy or an underlying pathological condition (such as pelvic malignancy) is suspected, it must be evaluated and the category adjusted after evaluation.
ENDOMETRIOSIS*	1	1	1	
BENIGN OVARIAN TUMOURS (including cysts)	1	1	1	
SEVERE DYSMENORRHOEA	1	1	1	
TROPHOBLAST DISEASE				
a) Benign gestational trophoblastic disease	1	1	1	
b) Malignant gestational trophoblastic disease	1	1	1	
CERVICAL ECTROPION*	1	1	1	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	2	2	2	Evidence: Limited evidence on women with low-grade squamous intraepithelial lesions found use of the vaginal ring did not worsen the condition. ²³
CERVICAL CANCER* (awaiting treatment)	2	2	2	
BREAST DISEASE* a) Undiagnosed mass b) Benign breast disease c) Family history of cancer d) Breast cancer (i) current (ii) past and no evidence of current disease for 5 years	2	2	2	Clarification: Evaluation should be pursued as early as possible.
ENDOMETRIAL CANCER*	1	1	1	
OVARIAN CANCER*	1	1	1	
UTERINE FIBROIDS* a) Without distortion of the uterine cavity b) With distortion of the uterine cavity	1	1	1	
PELVIC INFLAMMATORY DISEASE (PID)* a) Past PID (assuming no current risk factors for STIs) i) with subsequent pregnancy ii) without subsequent pregnancy b) PID - current	1	1	1	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
STIs*				
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	1	1	
b) Other STIs (excluding HIV and hepatitis)	1	1	1	
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	1	1	
d) Increased risk of STIs	1	1	1	
HIV/AIDS				
HIGH RISK OF HIV*	1	1	1	
HIV-INFECTED	1	1	1	
AIDS	1	1	1	
On ARV therapy	2	2	2	Clarification: If a woman is taking antiretroviral (ARV) therapy, refer to the section on drug interactions. Because there may be drug interactions between hormonal contraceptives and ARVs, AIDS with ARV therapy is classified as Category 2.
OTHER INFECTIONS				
SCHISTOSOMIASIS				
a) Uncomplicated	1	1	1	
b) Fibrosis of liver (if severe, see cirrhosis)	1	1	1	
TUBERCULOSIS				
a) Non-pelvic	1	1	1	Clarification: If a woman is taking rifampicin, refer to the section on drug interactions. Rifampicin is likely to decrease COC effectiveness. The extent to which CIC, P, or R use is similar to COC use in this regard remains unclear.
b) Known pelvic	1	1	1	
MALARIA	1	1	1	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
ENDOCRINE CONDITIONS				
DIABETES*				
a) History of gestational disease	1	1	1	
b) Non-vascular disease				
(i) non-insulin dependent	2	2	2	
(ii) insulin dependent	2	2	2	
c) Nephropathy/ retinopathy/ neuropathy	3/4	3/4	3/4	Clarification: The category should be assessed according to the severity of the condition.
d) Other vascular disease or diabetes of >20 years' duration	3/4	3/4	3/4	Clarification: The category should be assessed according to the severity of the condition.
THYROID DISORDERS				
a) Simple goitre	1	1	1	
b) Hyperthyroid	1	1	1	
c) Hypothyroid	1	1	1	
GASTROINTESTINAL CONDITIONS				
GALL-BLADDER DISEASE*				
a) Symptomatic				
(i) treated by cholecystectomy	2	2	2	
(ii) medically treated	2	3	3	
(iii) current	2	3	3	
b) Asymptomatic	2	2	2	
HISTORY OF CHOLESTASIS*				
a) Pregnancy-related	2	2	2	
b) Past COC or CIC related	2	3	3	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
VIRAL HEPATITIS*				
a) Active	3/4	4	4	Clarification: The category should be assessed according to the severity of the condition. Clarification: In women with symptomatic viral hepatitis, CICs, P and R should be withheld until liver function returns to normal or 3 months after the woman becomes asymptomatic, whichever occurs earlier.
b) Carrier	1	1	1	
CIRRHOSIS*				
a) Mild (compensated)	2	3	3	
b) Severe (decompensated)	3	4	4	
LIVER TUMOURS*				
a) Benign (adenoma)	3	4	4	
b) Malignant (hepatoma)	3/4	4	4	
ANAEMIAS				
THALASSAEMIA	1	1	1	
SICKLE CELL DISEASE	2	2	2	
IRON-DEFICIENCY ANAEMIA*	1	1	1	
DRUG INTERACTIONS				
DRUGS WHICH AFFECT LIVER ENZYMES				
a) Rifampicin	2	3	3	Clarification: Although the interaction of rifampicin or certain anticonvulsants with P or R use is not harmful to women, it is likely to reduce P or R effectiveness. Use of other contraceptives should be encouraged for women who are long-term users of any of these drugs.
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	2	3	3	
ANTIBIOTICS (excluding rifampicin)				
a) Griseofulvin	1	2	2	
b) Other antibiotics	1	1	1	

* See also additional comments at end of table

COMBINED INJECTABLE CONTRACEPTIVES (CICs), PATCH (P), RING (R)	CICs, patch, and ring do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation C=Continuation			CLARIFICATIONS/EVIDENCE
	CICs	P	R	
ANTIRETROVIRAL THERAPY	2	2	2	Clarification: It is important to note that antiretroviral drugs (ARV) have the potential to either decrease or increase the bioavailability of steroid hormones in hormonal contraceptives. The limited data available (outlined in Annex 1) suggest that potential drug interactions between many ARVs (particularly some non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs)) and hormonal contraceptives may alter safety and effectiveness of both the hormonal contraceptives and the ARVs. It is not known whether the contraceptive effectiveness of progestogen-only injectable contraceptives (such as depot medroxyprogesterone acetate and norethisterone enantate) would be compromised, as these methods provide higher blood hormone levels than other progestogen-only hormonal contraceptives, as well as than combined oral contraceptives. Studies are underway to evaluate potential interactions between depot medroxyprogesterone acetate and selected PI and NNRTI drugs. Thus, if a woman on ARV treatment decides to initiate or continue hormonal contraceptive use, the consistent use of condoms is recommended for preventing HIV transmission and may also compensate for any possible reduction in the effectiveness of the hormonal contraceptive.

* See also additional comments at end of table

Additional comments

AGE

Menarche to < 40 years: Theoretical concerns about the use of combined hormonal contraceptives among young adolescents have not been substantiated.

≥ 40 years: The risk of cardiovascular disease increases with age and may also increase with combined hormonal contraceptive use. In the absence of other adverse clinical conditions, combined hormonal contraceptives can be used until menopause.

BREASTFEEDING

< 6 weeks postpartum: There is some theoretical concern that the neonate may be at risk due to exposure to steroid hormones during the first 6 weeks postpartum.

≥ 6 weeks to < 6 months (primarily breastfeeding): Use of combined hormonal contraceptives during breastfeeding diminishes the quantity of breast milk, decreases the duration of lactation, and may thereby adversely affect the growth of the infant.

POSTPARTUM

< 21 days: There is some theoretical concern regarding the association between combined hormonal contraceptive use up to 3 weeks postpartum and risk of thrombosis in the mother. Blood coagulation and fibrinolysis are essentially normalized by 3 weeks postpartum.

PAST ECTOPIC PREGNANCY

The risk of future ectopic pregnancy is increased among women who have had an ectopic pregnancy in the past. Combined hormonal contraceptives provide protection against pregnancy in general, including ectopic gestation.

HYPERTENSION

Vascular disease: Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

DEEP VEIN THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)

Family history of DVT/PE (first-degree relatives): Some conditions which increase the risk of DVT/PE are heritable.

Major surgery: The degree of risk of DVT/PE associated with major surgery varies depending on the length of time that a woman is immobilized. There is no need to stop combined hormonal contraceptives prior to female surgical sterilization.

SUPERFICIAL VEIN THROMBOSIS

Varicose veins: Varicose veins are not risk factors for DVT/PE.

CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE

Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

STROKE

Among women with underlying vascular disease, the increased risk of arterial thrombosis associated with combined hormonal contraceptive use should be avoided.

VALVULAR HEART DISEASE

Among women with valvular heart disease, combined hormonal contraceptive use may further increase the risk of arterial thrombosis; women with complicated valvular heart disease are at greatest risk.

HEADACHES

Aura is a specific focal neurologic symptom. For more information on this and other diagnostic criteria, see: Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders, 2nd Edition. Cephalalgia. 2004; 24 (Suppl 1): 1- 150.

http://216.25.100.131/ihsccommon/guidelines/pdfs/ihc_II_main_no_print.pdf

VAGINAL BLEEDING PATTERNS

Irregular menstrual bleeding patterns are common among healthy women.

UNEXPLAINED VAGINAL BLEEDING

There are no conditions that cause vaginal bleeding that will be worsened in the short term by use of combined hormonal contraceptives.

ENDOMETRIOSIS

Combined hormonal contraceptives do not worsen, and may alleviate, the symptoms of endometriosis.

CERVICAL ECTROPION

Cervical ectropion is not a risk factor for cervical cancer, and there is no need for restriction of combined hormonal contraceptive use.

CERVICAL CANCER (awaiting treatment)

There is some theoretical concern that combined hormonal contraceptive use may affect prognosis of the existing disease. While awaiting treatment, women may use combined hormonal contraceptives. In general, treatment of this condition renders a woman sterile.

BREAST DISEASE

Family history of cancer: Women with BRCA1 or BRCA2 mutations have a much higher baseline risk of breast cancer than women who do not have these mutations. Most women with a family history of breast cancer do not have these mutations.

Breast cancer: Breast cancer is a hormonally sensitive tumour, and the prognosis of women with current or recent breast cancer may worsen with combined hormonal contraceptive use.

ENDOMETRIAL CANCER

It is not known whether CIC, P, or R use reduces the risk of developing endometrial cancer, as is the case with COCs. While awaiting treatment, women may use CICs, P or R. In general, treatment of this condition renders a woman sterile.

OVARIAN CANCER

It is not known whether CIC, P, or R use reduces the risk of developing ovarian cancer, as is the case with COCs. While awaiting treatment, women may use CICs, P, or R. In general, treatment of this condition renders a woman sterile.

UTERINE FIBROIDS

COCs do not appear to cause growth of uterine fibroids and CICs, P, or R are not expected to do either.

PELVIC INFLAMMATORY DISEASE (PID)

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs. Whether CICs, P or R reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STIs.

STIs

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs. Whether CICs, P or R reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STIs.

HIGH RISK OF HIV

COCs may reduce the risk of PID among women with STIs, but do not protect against HIV or lower genital tract STIs. Whether CICs, P or R reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STIs.

DIABETES

Although carbohydrate tolerance may change with combined hormonal contraceptive use, the major concerns are vascular disease due to diabetes and additional risk of arterial thrombosis due to combined hormonal contraceptive use.

GALL-BLADDER DISEASE

P or R, like COCs, may cause a small increased risk of gall-bladder disease. There is also concern that they may worsen existing gall-bladder disease. However, unlike COCs, CICs have been shown to have a minimal effect on liver function in healthy women, and have no first-pass effect on the liver.

HISTORY OF CHOLESTASIS

Pregnancy-related: History of pregnancy-related cholestasis may predict an increased risk of developing cholestasis associated with combined hormone therapy.

Past COC or CIC related: History of COC- or CIC-related cholestasis predicts an increased risk associated with P or R use. Unlike COCs, CICs have been shown to have a minimal effect on liver function in healthy women and have no first-pass effect on the liver.

VIRAL HEPATITIS

Active: Unlike COCs, CICs have been shown to have a minimal effect on liver function in healthy women and have no first-pass effect on the liver. However, because CICs are metabolized by the liver, they could, in theory, lead to adverse effects in women whose liver function is already compromised.

CIRRHOSIS

Unlike COCs, CICs have been shown to have a minimal effect on liver function in healthy women and have no first-pass effect on the liver. However, because CICs are metabolized by the liver, they could, in theory, lead to adverse effects in women whose liver function is already compromised

LIVER TUMOURS

Unlike COCs, CICs have been shown to have a minimal effect on liver function in healthy women and have no first-pass effect on the liver. However, because CICs are metabolized by the liver, they could, in theory, lead to adverse effects in women whose liver function is already compromised.

IRON-DEFICIENCY ANAEMIA

Combined hormonal contraceptive use may decrease menstrual blood loss.

References for combined injectable contraceptives, patch, and ring

1. Aedo AR et al. Pharmacokinetic and pharmacodynamic investigations with monthly injectable contraceptive preparations. *Contraception*, 1985, 31:453-69.
2. Garza-Flores J. Pharmacokinetics of once-a-month injectable contraceptives. *Contraception*, 1994, 49:347-59.
3. Fotherby K et al. A preliminary pharmacological trial of the monthly injectable contraceptive Cycloprovera. *Contraception*, 1982, 25:261-72.
4. Garza-Flores J et al. A multicentered pharmacokinetic, pharmacodynamic study of once-a-month injectable contraceptives. I. Different doses of HRP112 and of Depoprovera. *Contraception*, 1987, 36:441-57.
5. Said S et al. A multicentered phase III comparative study of two hormonal contraceptive preparations given once-a-month by intramuscular injection: I. Contraceptive efficacy and side effects. *Contraception*, 1988, 37:1-20.
6. Meng Y-X et al. Hemostatic changes in women using a monthly injectable contraceptive for one year. *Contraception*, 1990, 455-66.
7. Kesserü EV et al. A multicentered, two-year, phase III clinical trial of norethisterone enanthate 50 mg plus estradiol valerate 5 mg as a monthly injectable contraceptive. *Contraception*, 1991, 44:589-98.
8. Haiba NA et al. Clinical evaluation of two monthly injectable contraceptives and their effects on some metabolic parameters. *Contraception*, 1989, 39:619-32.
9. Pierson RA et al. Ortho Evra/Evra versus oral contraceptives: follicular development and ovulation in normal cycles and after an intentional dosing error. *Fertility & Sterility*, 2003, 80:34-42.
10. Audet M-C et al. Evaluation of contraceptive efficacy and cycle control of a transdermal contraceptive patch vs. an oral contraceptive: a randomized controlled trial. *JAMA*, 2001, 285:2347-54.
11. Smallwood GH et al. Efficacy and safety of a transdermal contraceptive system. *Obstetrics and Gynecology*, 2001, 98:799-805.
12. Helmerhorst FM et al. Comparison of efficacy, cycle control, compliance and safety in users of a contraceptive patch vs. an oral contraceptive. XVI FIGO World Congress of Gynecology and Obstetrics [FC 2.30.06], Washington DC, USA, 3-8 September 2000.
13. Dittrich R et al. Transdermal contraception: evaluation of three transdermal norelgestromin/ethinylestradiol doses in a randomized, multicenter, dose-response study. *American Journal of Obstetrics and Gynecology*, 2002, 186:15-20.
14. Abrams LS et al. Pharmacokinetics of a contraceptive patch (Evra/Ortho Evra) containing norelgestromin and ethinylestradiol at four application sites. *British Journal of Clinical Pharmacology*, 2002, 53:141-6.
15. Abrams LS et al. Multiple-dose pharmacokinetics of a contraceptive patch in healthy women participants. *Contraception*, 2001, 64:287-94.
16. Abrams LS et al. Pharmacokinetics of norelgestromin and ethinylestradiol delivered by a contraceptive patch (Ortho Evra/Evra) under conditions of heat, humidity, and exercise. *Journal of Clinical Pharmacology*, 2001, 41:1301-9.
17. Ziemann M. The introduction of a transdermal hormonal contraceptive (Ortho Evra/Evra). *Fertility & Sterility*, 2002, 77(2): s1-s2.
18. Burkman, RT. The transdermal contraceptive patch: a new approach to hormonal contraception. *International Journal of Fertility*, 2002, 47:69-76.
19. Stone SC. Desogestrel. *Clinical Obstetrics and Gynecology*, 1995, 38:821-8.
20. Mulders TMT, Dieben TOM. Use of the novel combined contraceptive vaginal ring NuvaRing for ovulation inhibition. *Fertility & Sterility*, 2001, 75:865-70.
21. Timmer CJ, Mulders TMT. Pharmacokinetics of etonogestrel and ethinylestradiol released from a combined contraceptive vaginal ring. *Clinical Pharmacokinetics*, 2000, 39:233-42.
22. Bjarnadottir RI, Tuppurainen M, Killick SR. Comparison of cycle control with a combined contraceptive vaginal ring and oral levonorgestrel/ethinyl estradiol. *American Journal of Obstetrics and Gynecology*, 2002, 186:389-95.
23. Dieben TOM, Roumen FJME, Apter D. Efficacy, cycle control, and user acceptability of a novel combined contraceptive vaginal ring. *Obstetrics and Gynecology*, 2002, 100:585-93.
24. Roumen FJME et al. The cervico-vaginal epithelium during 20 cycles of use of a combined contraceptive vaginal ring. *Human Reproduction*, 1996, 11:2443-8.
25. Davies GC, Feng LX, Newton JR. The effects of a combined contraceptive vaginal ring releasing ethinylestradiol and 3-ketodesogestrel on vaginal flora. *Contraception*, 1992, 45:511-8.

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	1
Pregnancy.....	1
Age.....	1
Parity.....	2
Breastfeeding.....	2
Postpartum.....	2
Post-abortion.....	2
Past ectopic pregnancy.....	3
History of pelvic surgery.....	3
Smoking.....	3
Obesity.....	3
Blood pressure measurement unavailable.....	3
CARDIOVASCULAR DISEASE	3
Multiple risk factors for arterial cardiovascular disease.....	3
Hypertension.....	4
History of high blood pressure during pregnancy.....	4
Deep venous thrombosis (DVT)/Pulmonary embolism (PE).....	5
Known thrombogenic mutations.....	5
Superficial venous thrombosis.....	5
Current and history of ischaemic heart disease.....	5
Stroke.....	5
Known hyperlipidaemias.....	6
Valvular heart disease.....	6
NEUROLOGIC CONDITIONS	6
Headaches.....	6
Epilepsy.....	6
DEPRESSIVE DISORDERS	7
Depressive disorders.....	7
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS	7
Vaginal bleeding patterns.....	7
Unexplained vaginal bleeding.....	7
Endometriosis.....	7
Benign ovarian tumours.....	7
Severe dysmenorrhoea.....	7
Trophoblast disease.....	8
Cervical ectropion.....	8
Cervical intraepithelial neoplasia (CIN).....	8
Cervical cancer.....	8
Breast disease.....	8
Endometrial cancer.....	8
Ovarian cancer.....	8
Uterine fibroids.....	8
Pelvic inflammatory disease (PID).....	9
STIs.....	9
HIV/AIDS	9
High risk of HIV.....	9
HIV-infected.....	9
AIDS.....	10

OTHER INFECTIONS	10
Schistosomiasis	10
Tuberculosis.....	10
Malaria	10
ENDOCRINE CONDITIONS.....	10
Diabetes.....	10
Thyroid disorders	11
GASTROINTESTINAL CONDITIONS.....	11
Gall-bladder disease	11
History of cholestasis	11
Viral hepatitis	11
Cirrhosis.....	11
Liver tumours	12
ANAEMIAS.....	12
Thalassaemia.....	12
Sickle cell disease.....	12
Iron-deficiency anaemia	12
DRUG INTERACTIONS	12
Drugs which affect liver enzymes.....	12
Antibiotics	12
Antiretroviral therapy.....	13
Additional comments	14
References for progestogen-only contraception.....	16

PROGESTOGEN-ONLY CONTRACEPTIVES

POP = Progestogen-only pill (POP)
 D/NE = Depot medroxyprogesterone acetate (DMPA)/norethisterone enantate (NET-EN)
 LNG/ETG = Levonorgestrel implants (Norplant and Jadelle) and etonogestrel implants (Implanon)

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY				
PREGNANCY	NA			Clarification: Use of POCs is not required. There is no known harm to the woman, the course of her pregnancy, or the fetus if POCs are accidentally used during pregnancy. However, the relationship between DMPA use during pregnancy and its effects on the fetus remains unclear.
AGE*				
a) Menarche to < 18 years	1	2	1	Evidence: Limited evidence shows decreased bone mineral density over time among adolescent DMPA users, but not among levonorgestrel implant users. No studies have examined whether DMPA use among adolescents affects peak bone mass levels. ¹⁻⁵
b) 18 to 45 years	1	1	1	Evidence: In general, current DMPA users had decreased bone mineral density compared with non-users; this decrease was usually within one standard deviation of normal values. ⁶ Results for current Norplant users were mixed. ⁶ One study of Implanon users showed no change in bone mineral density over two years. ⁷
c) > 45 years	1	2	1	Evidence: Older DMPA users had decreased bone mineral density compared with non-users. However, limited evidence found that women gained bone mass following discontinuation of DMPA prior to menopause. Further, among postmenopausal women, there was no difference in bone mineral density between former DMPA users and never users. ⁸⁻¹³

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
PARITY				
a) Nulliparous	1	1	1	
b) Parous	1	1	1	
BREASTFEEDING				
a) < 6 weeks postpartum	3	3	3	<p>Clarification: There is concern that the neonate may be at risk of exposure to steroid hormones during the first 6 weeks postpartum. However, in many settings pregnancy morbidity and mortality risks are high, and access to services is limited. In such settings, POCs may be one of the few types of methods widely available and accessible to breastfeeding women immediately postpartum.</p> <p>Evidence: Studies have shown that among breast-feeding women less than 6 weeks postpartum, progestogen-only contraceptives did not affect breast-feeding performance and infant health and growth. However, there are no data evaluating the effects of progestogen exposure via breast milk on brain and liver development.¹⁴⁻³⁸</p>
b) ≥ 6 weeks to < 6 months postpartum (primarily breastfeeding)	1	1	1	
c) ≥ 6 months postpartum	1	1	1	
POSTPARTUM* (in non-breastfeeding women)				
a) < 21 days	1	1	1	
b) ≥ 21 days	1	1	1	
POST-ABORTION				
a) First trimester	1	1	1	<p>Clarification: POCs may be started immediately post-abortion.</p> <p>Evidence: Limited evidence suggests that there are no adverse side effects when Norplant or NET-EN are initiated after a first trimester abortion.³⁹⁻⁴²</p>
b) Second trimester	1	1	1	
c) Immediate post-septic abortion	1	1	1	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
PAST ECTOPIC PREGNANCY*	2	1	1	
HISTORY OF PELVIC SURGERY	1	1	1	
SMOKING				
a) Age < 35 years	1	1	1	
b) Age ≥ 35 years				
i) <15 cigarettes/day	1	1	1	
ii) ≥15 cigarettes/day	1	1	1	
OBESITY ≥ 30 kg/m ² body mass index (BMI)	1	1	1	Evidence: Studies provide conflicting evidence regarding whether obese women are at increased risk of weight gain and bleeding problems with DMPA use relative to non-obese women with DMPA use. ⁴³⁻⁴⁵ Studies show that obese women do not experience decreased effectiveness when using Norplant soft capsules or Jadelle. ⁴⁶⁻⁴⁸
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA	NA	NA	Clarification: It is desirable to have blood pressure measurements taken before initiation of POC use. However, in some settings blood pressure measurements are unavailable. In many of these settings, pregnancy morbidity and mortality risks are high, and POCs are one of the few types of methods widely available. In such settings, women should not be denied use of POCs simply because their blood pressure cannot be measured.
CARDIOVASCULAR DISEASE				
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	2	3	2	Clarification: When multiple major risk factors exist, risk of cardiovascular disease may increase substantially. Some POCs may increase the risk of thrombosis, although this increase is substantially less than with COCs. The effects of DMPA and NET-EN may persist for some time after discontinuation.

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
HYPERTENSION*				
For all categories of hypertension, classifications are based on the assumption that no other risk factors for cardiovascular disease exist. When multiple risk factors do exist, risk of cardiovascular disease may increase substantially. A single reading of blood pressure level is not sufficient to classify a woman as hypertensive.				
a) History of hypertension where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	2	2	2	Clarification: It is desirable to have blood pressure measurements taken before initiation of POC use. However, in some settings blood pressure measurements are unavailable. In many of these settings pregnancy morbidity and mortality risks are high, and POCs are one of the few types of methods widely available. In such settings, women should not be denied use of POCs simply because their blood pressure cannot be measured. Clarification: Women adequately treated for hypertension are at reduced risk of acute myocardial infarction and stroke as compared with untreated women. Although there are no data, POC users with adequately controlled and monitored hypertension should be at reduced risk of acute myocardial infarction and stroke compared with untreated hypertensive POC users. Evidence: Limited evidence suggests that among women with hypertension, those who used POPs or progestogen-only injectables had a small increased risk of cardiovascular events compared with women who did not use these methods. ⁴⁹
b) Adequately controlled hypertension where blood pressure CAN be evaluated	1	2	1	
c) Elevated blood pressure levels (properly taken measurements)				
i) systolic 140-159 or diastolic 90-99	1	2	1	
ii) systolic \geq 160 or diastolic \geq 100	2	3	2	
d) Vascular disease	2	3	2	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	1	1	1	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
DEEP VENOUS THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)*				
a) History of DVT/PE	2	2	2	
b) Current DVT/PE	3	3	3	
c) Family history of DVT/PE (first-degree relatives)	1	1	1	
d) Major surgery				
i) with prolonged immobilization	2	2	2	
ii) without prolonged immobilization	1	1	1	
e) Minor surgery without immobilization	1	1	1	
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	2	2	2	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
SUPERFICIAL VENOUS THROMBOSIS				
a) Varicose veins	1	1	1	
b) Superficial thrombophlebitis	1	1	1	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE*	I C 2 3	3	I C 2 3	
STROKE* (history of cerebrovascular accident)	I C 2 3	3	I C 2 3	

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.						
CONDITION	CATEGORY I=Initiation, C=Continuation						CLARIFICATIONS/EVIDENCE
	POP	D/NE		LNG/ETG			
KNOWN HYPERLIPIDAEMIAS	2	2		2		Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening. Some types of hyperlipidaemias are risk factors for vascular disease.	
VALVULAR HEART DISEASE							
a) Uncomplicated	1	1		1			
b) Complicated (pulmonary hypertension, risk of atrial fibrillation, history of subacute bacterial endocarditis)	1	1		1			
NEUROLOGIC CONDITIONS							
HEADACHES*	I	C	I	C	I	C	
a) Non-migrainous (mild or severe)	1	1	1	1	1	1	Clarification: Classification depends on accurate diagnosis of those severe headaches that are migrainous and those that are not. Any new headaches or marked changes in headaches should be evaluated. Classification is for women without any other risk factors for stroke. Risk of stroke increases with age, hypertension, and smoking.
b) Migraine							
i) without aura							
Age < 35	1	2	2	2	2	2	
Age ≥ 35	1	2	2	2	2	2	
ii) with aura, at any age	2	3	2	3	2	3	
EPILEPSY	1	1		1		Clarification: If a woman is taking anticonvulsants, refer to the section on drug interactions. Certain anticonvulsants lower POC effectiveness.	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
DEPRESSIVE DISORDERS				
DEPRESSIVE DISORDERS	1	1	1	Clarification: The classification is based on data for women with selected depressive disorders. No data on bipolar disorder or postpartum depression were available. There is a potential for drug interactions between certain antidepressant medications and hormonal contraceptives. Evidence: POCs did not increase depressive symptoms in women with depression compared to baseline. ⁵⁰⁻⁵³
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS				
VAGINAL BLEEDING PATTERNS*				
a) Irregular pattern <i>without</i> heavy bleeding	2	2	2	
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	2	2	2	Clarification: Unusually heavy bleeding should raise the suspicion of a serious underlying condition.
UNEXPLAINED VAGINAL BLEEDING* (suspicious for serious underlying condition)				
Before evaluation	2	3	3	Clarification: If pregnancy or an underlying pathological condition (such as pelvic malignancy) is suspected, it must be evaluated and the category adjusted after evaluation.
ENDOMETRIOSIS	1	1	1	
BENIGN OVARIAN TUMOURS (including cysts)	1	1	1	
SEVERE DYSMENORRHOEA	1	1	1	

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
TROPHOBLAST DISEASE				
a) Benign gestational trophoblastic disease	1	1	1	
b) Malignant gestational trophoblastic disease	1	1	1	
CERVICAL ECTROPION	1	1	1	
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	1	2	2	Evidence: Among women with persistent HPV infection, long-term DMPA use (≥ 5 years) may increase the risk of carcinoma in situ and invasive carcinoma. ⁵⁴
CERVICAL CANCER (awaiting treatment)*	1	2	2	
BREAST DISEASE*				
a) Undiagnosed mass	2	2	2	Clarification: Evaluation should be pursued as early as possible.
b) Benign breast disease	1	1	1	
c) Family history of cancer	1	1	1	
d) Breast cancer				
(i) current	4	4	4	
(ii) past and no evidence of current disease for 5 years	3	3	3	
ENDOMETRIAL CANCER*	1	1	1	
OVARIAN CANCER*	1	1	1	
UTERINE FIBROIDS*				
a) Without distortion of the uterine cavity	1	1	1	
b) With distortion of the uterine cavity	1	1	1	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
PELVIC INFLAMMATORY DISEASE (PID)*				
a) Past PID (assuming no current risk factors for STIs)				
(i) with subsequent pregnancy	1	1	1	
(ii) without subsequent pregnancy	1	1	1	
b) PID - current	1	1	1	
STIs*				
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	1	1	Evidence: Limited evidence suggests that there may be an increased risk of chlamydial cervicitis among DMPA users at high risk of STIs. For other STIs, there is either evidence of no association between DMPA use and STI acquisition or too limited evidence to draw any conclusions. There is no evidence for other POCs. ⁵⁵⁻⁶¹
b) Other STIs (excluding HIV and hepatitis)	1	1	1	
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	1	1	
d) Increased risk of STIs	1	1	1	
HIV/AIDS				
HIGH RISK OF HIV*	1	1	1	Evidence: Overall, evidence is inconsistent regarding whether there is any increased risk of HIV acquisition among POC users compared with non-users. ⁶²⁻⁷⁸
HIV-INFECTED	1	1	1	Evidence: Studies are conflicting regarding whether there is an increased risk of HIV and herpes simplex virus (HSV) shedding among HIV-infected women using DMPA. ⁷⁹⁻⁸¹

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
AIDS On ARV therapy	1 2	1 2	1 2	Clarification: If a woman is taking antiretroviral (ARV) therapy, refer to the section on drug interactions. Because there may be drug interactions between hormonal contraceptives and ARVs, AIDS with ARV therapy is classified as Category 2.
OTHER INFECTIONS				
SCHISTOSOMIASIS				
a) Uncomplicated	1	1	1	Evidence: Among women with uncomplicated schistosomiasis, limited evidence showed that DMPA use had no adverse effects on liver function. ⁸²
b) Fibrosis of liver (if severe, see cirrhosis)	1	1	1	
TUBERCULOSIS				
a) Non-pelvic	1	1	1	Clarification: If a woman is taking rifampicin, refer to the section on drug interactions. Rifampicin is likely to decrease POC effectiveness.
b) Known pelvic	1	1	1	
MALARIA	1	1	1	
ENDOCRINE CONDITIONS				
DIABETES*				
a) History of gestational disease	1	1	1	
b) Non-vascular disease				
(i) non-insulin dependent	2	2	2	
(ii) insulin dependent	2	2	2	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
DIABETES (Cont'd)				
c) Nephropathy/ retinopathy/ neuropathy	2	3	2	
d) Other vascular disease or diabetes of >20 years' duration	2	3	2	
THYROID DISORDERS				
a) Simple goitre	1	1	1	
b) Hyperthyroid	1	1	1	
c) Hypothyroid	1	1	1	
GASTROINTESTINAL CONDITIONS				
GALL-BLADDER DISEASE				
a) Symptomatic				
(i) treated by cholecystectomy	2	2	2	
(ii) medically treated	2	2	2	
(iii) current	2	2	2	
b) Asymptomatic	2	2	2	
HISTORY OF CHOLESTASIS*				
a) Pregnancy-related	1	1	1	
b) Past COC-related	2	2	2	
VIRAL HEPATITIS*				
a) Active	3	3	3	
b) Carrier	1	1	1	
CIRRHOSIS*				
a) Mild (compensated)	2	2	2	
b) Severe (decompensated)	3	3	3	

* See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
LIVER TUMOURS*				
a) Benign (adenoma)	3	3	3	
b) Malignant (hepatoma)	3	3	3	
ANAEMIAS				
THALASSAEMIA	1	1	1	
SICKLE CELL DISEASE	1	1	1	Evidence: Among women with sickle cell disease, POC use did not have adverse effects on haematological parameters and, in some studies, was beneficial with respect to clinical symptoms. ⁸³⁻⁹⁰
IRON-DEFICIENCY ANAEMIA*	1	1	1	
DRUG INTERACTIONS				
DRUGS WHICH AFFECT LIVER ENZYMES				
a) Rifampicin	3	2	3	Clarification: Although the interaction of rifampicin or certain anticonvulsants with POPs and LNG/ETG implants is not harmful to women, it is likely to reduce the effectiveness of POPs and LNG/ETG implants. Use of other contraceptives should be encouraged for women who are long-term users of any of these drugs. Whether increasing the hormone dose of POPs alleviates this concern remains unclear. Evidence: Use of certain anticonvulsants decreased the contraceptive effectiveness of POCs. ⁹¹⁻⁹³
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	3	2	3	
ANTIBIOTICS (excluding rifampicin)				
a) Griseofulvin	2	1	2	
b) Other antibiotics	1	1	1	

See also additional comments at end of table

PROGESTOGEN-ONLY CONTRACEPTIVES (POCs)	POCs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation			CLARIFICATIONS/EVIDENCE
	POP	D/NE	LNG/ETG	
ANTIRETROVIRAL THERAPY	2	2	2	Clarification: It is important to note that antiretroviral drugs (ARV) have the potential to either decrease or increase the bioavailability of steroid hormones in hormonal contraceptives. The limited data available (outlined in Annex 1) suggest that potential drug interactions between many ARVs (particularly some non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs)) and hormonal contraceptives may alter safety and effectiveness of both the hormonal contraceptives and the ARVs. It is not known whether the contraceptive effectiveness of progestogen-only injectable contraceptives (such as depot medroxyprogesterone acetate and norethisterone enantate) would be compromised, as these methods provide higher blood hormone levels than other progestogen-only hormonal contraceptives, as well as than combined oral contraceptives. Studies are underway to evaluate potential interactions between depot medroxyprogesterone acetate and selected PI and NNRTI drugs. Thus, if a woman on ARV treatment decides to initiate or continue hormonal contraceptive use, the consistent use of condoms is recommended for preventing HIV transmission and may also compensate for any possible reduction in the effectiveness of the hormonal contraceptive.

* See also additional comments at end of table

Additional comments

AGE

Menarche to < 18 years: For women under 18 years of age, there are theoretical concerns regarding the hypo-estrogenic effects of DMPA use, including whether these women will achieve their appropriate peak bone mass.

> 45 years: For women greater than age 45, there are theoretical concerns regarding hypo-estrogenic effects of DMPA use, including whether these women will regain all lost bone mass after discontinuation of DMPA.

POSTPARTUM

< 21 days: POCs may be safely used by non-breastfeeding women immediately postpartum.

PAST ECTOPIC PREGNANCY

POPs have a higher absolute rate of ectopic pregnancy compared with other POCs, but still less than using no method.

HYPERTENSION

Vascular disease: There is concern regarding hypo-estrogenic effects and reduced HDL levels, particularly among users of DMPA and NET-EN. However, there is little concern about these effects with regard to POPs or LNG/ETG implants. The effects of DMPA and NET-EN may persist for some time after discontinuation.

DEEP VEIN THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)

Some POCs may increase the risk of venous thrombosis, although this increase is substantially less than with COCs.

CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE

There is concern regarding hypo-estrogenic effects and reduced HDL levels, particularly among users of DMPA and NET-EN. However, there is little concern about these effects with regard to POPs or LNG/ETG implants. The effects of DMPA and NET-EN may persist for some time after discontinuation.

STROKE

There is concern regarding hypo-estrogenic effects and reduced HDL levels, particularly among users of DMPA and NET-EN. However, there is little concern about these effects with regard to POPs or LNG/ETG implants. The effects of DMPA and NET-EN may persist for some time after discontinuation.

HEADACHES

Aura is a specific focal neurologic symptom. For more information on this and other diagnostic criteria, see: Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders, 2nd Edition. Cephalalgia. 2004; 24 (Suppl 1): 1- 150.

http://216.25.100.131/ihscommon/guidelines/pdfs/ihc_II_main_no_print.pdf

There is concern that severe headaches may increase with use of NET-EN, DMPA and implants. The effects of NET-EN and DMPA may persist for some time after discontinuation.

VAGINAL BLEEDING PATTERNS

Irregular menstrual bleeding patterns are common among healthy women. POC use frequently induces an irregular bleeding pattern. Implant use may induce irregular bleeding patterns, especially during the first 3-6 months, but these patterns may persist longer. ETG users are more likely than LNG users to develop amenorrhoea.

UNEXPLAINED VAGINAL BLEEDING

POCs may cause irregular bleeding patterns which may mask symptoms of underlying pathology. The effects of DMPA and NET-EN may persist for some time after discontinuation.

CERVICAL CANCER (awaiting treatment)

There is some theoretical concern that POC use may affect prognosis of the existing disease. While awaiting treatment, women may use POCs. In general, treatment of this condition renders a woman sterile.

BREAST DISEASE

Breast cancer: Breast cancer is a hormonally sensitive tumour, and the prognosis of women with current or recent breast cancer may worsen with POC use.

ENDOMETRIAL CANCER

While awaiting treatment, women may use POCs. In general, the treatment of this condition renders a woman sterile.

OVARIAN CANCER

While awaiting treatment, women may use POCs. In general, the treatment of this condition renders a woman sterile.

UTERINE FIBROIDS

POCs do not appear to cause growth of uterine fibroids.

PELVIC INFLAMMATORY DISEASE (PID)

Whether POCs, like COCs, reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STI.

STIs

Whether POCs, like COCs, reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STI.

HIGH RISK OF HIV

Whether POCs, like COCs, reduce the risk of PID among women with STIs is unknown, but they do not protect against HIV or lower genital tract STI.

DIABETES

Non-vascular disease: POCs may alter carbohydrate metabolism.

Nephropathy, retinopathy, neuropathy: There is concern regarding hypo-estrogenic effects and reduced HDL levels, particularly among users of DMPA and NET-EN. The effects of DMPA and NET-EN may persist for some time after discontinuation. Some POCs may increase the risk of thrombosis, although this increase is substantially less than with COCs.

Other vascular disease or diabetes of > 20 years' duration: There is concern regarding hypo-estrogenic effects and reduced HDL levels, particularly among users of DMPA and NET-EN. The effects of DMPA and NET-EN may persist for some time after discontinuation. Some POCs may increase the risk of thrombosis, although this increase is substantially less than with COCs.

HISTORY OF CHOLESTASIS

Theoretically, a history of COC-related cholestasis may predict subsequent cholestasis with POC use. However, this has not been documented.

VIRAL HEPATITIS

Active: POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. This concern is similar to, but less than, that with COCs.

CIRRHOSIS

POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. This concern is similar to, but less than, that with COCs.

LIVER TUMOURS

POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. In addition, POC use may enhance the growth of tumours. This concern is similar to, but less than, that with COCs.

IRON-DEFICIENCY ANAEMIA

Changes in the menstrual pattern associated with POC use have little effect on haemoglobin levels.

References for progestogen-only contraception

1. Busen NH, Britt RB, Rianon N. Bone mineral density in a cohort of adolescent women using depot medroxyprogesterone acetate for one to two years. *Journal of Adolescent Health*, 2003, 32:257-9.
2. Cromer BA et al. A prospective comparison of bone density in adolescent girls receiving depot medroxyprogesterone acetate (Depo-Provera), levonorgestrel (Norplant), or oral contraceptives. *Journal of Pediatrics*, 1996, 129:671-6.
3. Cundy T et al. Spinal bone density in women using depot medroxyprogesterone contraception. *Obstetrics & Gynecology*, 1998, 92:569-73.
4. Lappe JM, Stegman MR, Recker RR. The impact of lifestyle factors on stress fractures in female army recruits. *Osteoporosis International*, 2001, 12:35-42.
5. Tharnprisan W, Taneepanichskul S. Bone mineral density in adolescent and young Thai girls receiving oral contraceptives compared with depot medroxyprogesterone acetate: a cross-sectional study in young Thai women. *Contraception*, 2002, 66:101-3.
6. Banks E, Berrington A, Casabonne D. Overview of the relationship between use of progestogen-only contraceptives and bone mineral density. *British Journal of Obstetrics and Gynaecology*, 2001, 108:1214-21.
7. Beerthuis R et al. Bone mineral density during long-term use of the progestagen contraceptive implant Implanon compared to a non-hormonal method of contraception. *Human Reproduction*, 2000, 15:118-22.
8. Cundy T et al. Recovery of bone density in women who stop using medroxyprogesterone acetate. *BMJ*, 1994, 308:247-8.
9. Cundy T, Reid I. Depot medroxyprogesterone and bone density. *BMJ*, 1994, 308:1567-8.
10. Cundy T et al. Menopausal bone loss in long-term users of depot medroxyprogesterone acetate contraception. *American Journal of Obstetrics & Gynecology*, 2002, 186:978-83.
11. Gbolade B et al. Bone density in long term users of depot medroxyprogesterone acetate. *British Journal of Obstetrics & Gynaecology*, 1998, 105:790-4.
12. Orr-Walker BJ et al. The effect of past use of the injectable contraceptive depot medroxyprogesterone acetate on bone mineral density in normal post-menopausal women. *Clinical Endocrinology*, 1998, 49:615-8.
13. Tang OS et al. Further evaluation on long-term depot-medroxyprogesterone acetate use and bone mineral density: a longitudinal cohort study. *Contraception*, 2000, 62:161-4.
14. Abdel-Aleem H et al. The use of nomegestrol acetate subdermal contraceptive implant, Uniplant, during lactation. *Contraception*, 1996, 54:281-6.
15. Abdulla KA et al. Effect of early postpartum use of the contraceptive implants, Norplant, on the serum levels of immunoglobulins of the mothers and their breastfed infants. *Contraception*, 1985, 32:261-6.
16. Bjarnadottir RI et al. Comparative study of the effects of a progestogen-only pill containing desogestrel and an intrauterine contraceptive device in lactating women. *British Journal of Obstetrics and Gynecology*, 2001, 108:1174-80.
17. Croxatto HB et al. Fertility regulation in nursing women. Comparative performance of progesterone implants versus placebo and copper T. *American Journal of Obstetrics & Gynecology*, 1982, 144:201-8.
18. Diaz S et al. Fertility regulation in nursing women. VI. Contraceptive effectiveness of a subdermal progesterone implant. *Contraception*, 1984, 30:311-25.
19. Halderman LD, Nelson AL. Impact of early postpartum administration of progestin-only hormonal contraceptives compared with nonhormonal contraceptives on short-term breast-feeding patterns. *American Journal of Obstetrics & Gynecology*, 2002, 186:1250-6.
20. Hannon PR et al. The influence of medroxyprogesterone on the duration of breast-feeding in mothers in an urban community. *Archives of Pediatrics & Adolescent Medicine*, 1997, 151:490-6.
21. Jimenez J et al. Long-term follow-up of children breast-fed by mothers receiving depot-medroxyprogesterone acetate. *Contraception*, 1984, 30:523-33.
22. Kamal I et al. Clinical, biochemical, and experimental studies on lactation: clinical effects of steroids on the initiation of lactation. *American Journal of Obstetrics & Gynecology*, 1970, 108:655-8.
23. Karim M et al. Injected progestogen and lactation. *British Medical Journal*, 1971, 1:200-3.
24. Massai R et al. Preregistration study on the safety and contraceptive efficacy of a progesterone-releasing vaginal ring in Chilean nursing women. *Contraception*, 1999, 60:9-14.
25. McCann MF et al. The effects of a progestin-only oral contraceptive (levonorgestrel 0.03 mg) on breast-feeding. *Contraception*, 1989, 40:635-48.

26. McEwan JA et al. Early experience in contraception with a new progestogen. *Contraception*, 1977, 16:339-50.
27. Melis GB et al. Norethisterone enanthate as an injectable contraceptive in puerperal and non-puerperal women. *Contraception*, 1981, 23:77-88.
28. Moggia AV et al. A comparative study of a progestin-only oral contraceptive versus non-hormonal methods in lactating women in Buenos Aires, Argentina.[Erratum appears in *Contraception*, 1991 Sep; 44(3):339]. *Contraception*, 1991, 44:31-43.
29. Narducci U, Piatti N. [Use of Depo-Provera as a contraceptive during the puerperium]. *Minerva Ginecologica*, 1973, 107-11.
30. Seth U et al. Effect of a subdermal silastic implant containing norethindrone acetate on human lactation. *Contraception*, 1977, 16:383-98.
31. Shaaban MM, Salem HT, Abdullah KA. Influence of levonorgestrel contraceptive implants, Norplant, initiated early postpartum upon lactation and infant growth. *Contraception*, 1985, 32:623-35.
32. Shaaban MM. Contraception with progestogens and progesterone during lactation. *Journal of Steroid Biochemistry & Molecular Biology*, 1991, 40:705-10.
33. Shikary ZK et al. Pharmacodynamic effects of levonorgestrel (LNG) administered either orally or subdermally to early postpartum lactating mothers on the urinary levels of follicle stimulating hormone (FSH), luteinizing hormone (LH) and testosterone (T) in their breast-fed male infants. *Contraception*, 1986, 34:403-12.
34. Sivin I et al. Contraceptives for lactating women: a comparative trial of a progesterone-releasing vaginal ring and the copper t 380a iud. *Contraception*, 1997, 55:225-32.
35. Velasquez J et al. [Effect of daily oral administration of 0.350 mg of norethindrone on lactation and chemical composition of milk]. *Ginecologia y Obstetricia de Mexico*, 1976, 40(237):31-9.
36. West CP. The acceptability of a progestagen-only contraceptive during breast-feeding. *Contraception*, 1983, 27:563-9.
37. Zanartu J, Aguilera E, Munoz-Pinto G. Maintenance of lactation by means of continuous low-dose progestogen given post-partum as a contraceptive. *Contraception*, 1976, 13:313-8.
38. Reinprayoon D et al. Effects of the etonogestrel-releasing contraceptive implant (Implanon) on parameters of breastfeeding compared to those of an intrauterine device. *Contraception*, 2000, 62:239-46.
39. Kurunmaki H. Contraception with levonorgestrel-releasing subdermal capsules, Norplant, after pregnancy termination. *Contraception*, 1983, 27:473-82.
40. Kurunmaki H et al. Immediate postabortal contraception with Norplant: levonorgestrel, gonadotropin, estradiol, and progesterone levels over two postabortal months and return of fertility after removal of Norplant capsules. *Contraception*, 1984, 30:431-42.
41. Lahteenmaki P, Toivonen J, Lahteenmaki PL. Postabortal contraception with norethisterone enanthate injections. *Contraception*, 1983, 27:553-62.
42. Ortayli N et al. Immediate postabortal contraception with the levonorgestrel intrauterine device, Norplant, and traditional methods. *Contraception*, 2001, 63:309-14.
43. Connor PD et al. Determining risk between Depo-Provera use and increased uterine bleeding in obese and overweight women. *Journal of the American Board of Family Practice*, 2002, 15:7-10.
44. Leiman G. Depo-medroxyprogesterone acetate as a contraceptive agent: its effect on weight and blood pressure. *American Journal of Obstetrics & Gynecology*, 1972, 114:97-102.
45. Mangan SA, Larsen PG, Hudson S. Overweight teens at increased risk for weight gain while using depot medroxyprogesterone acetate. *Journal of Pediatric & Adolescent Gynecology*, 2002, 15:79-82.
46. Sivin I et al. Contraception with two levonorgestrel rod implants. A 5-year study in the United States and Dominican Republic. *Contraception*, 1998, 58:275-82.
47. Sivin I et al. Prolonged effectiveness of Norplant(R) capsule implants: a 7-year study. *Contraception*, 2000, 61:187-94.
48. Sivin I et al. Levonorgestrel concentrations during 7 years of continuous use of Jadelle contraceptive implants. *Contraception*, 2001, 64:43-9.
49. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Cardiovascular disease and use of oral and injectable progestogen-only contraceptives and combined injectable contraceptives. Results of an international, multicenter, case-control study. *Contraception*, 1998, 57:315-24.
50. Cromer BA et al. A prospective study of adolescents who choose among levonorgestrel implant (Norplant), medroxyprogesterone acetate (Depo-Provera), or the combined oral contraceptive pill as contraception. *Pediatrics*, 1994, 94:687-94.

51. Gupta N et al. Mood changes in adolescents using depot-medroxyprogesterone acetate for contraception: a prospective study. *Journal of Pediatric & Adolescent Gynecology*, 2001, 14:71-6.
52. Westhoff C et al. Depressive symptoms and Norplant contraceptive implants. *Contraception*, 1998, 57:241-5.
53. Westhoff C et al. Depressive symptoms and Depo-Provera. *Contraception*, 1998, 57:237-40.
54. Smith JS et al. Cervical cancer and use of hormonal contraceptives: a systematic review. *Lancet*, 2003, 361:1159-67.
55. Baeten JM et al. Hormonal contraception and risk of sexually transmitted disease acquisition: results from a prospective study. *American Journal of Obstetrics & Gynecology*, 2001, 185:380-5.
56. Giuliano AR et al. Human papillomavirus infection at the United States-Mexico border: implications for cervical cancer prevention and control. *Cancer Epidemiology, Biomarkers & Prevention*, 2001, 10:1129-36.
57. Jacobson DL et al. Relationship of hormonal contraception and cervical ectopy as measured by computerized planimetry to chlamydial infection in adolescents. *Sexually Transmitted Diseases*, 2000, 27:313-9.
58. Lavreys L et al. Human herpesvirus 8: seroprevalence and correlates in prostitutes in Mombasa, Kenya. *Journal of Infectious Diseases*, 2003, 187:359-63.
59. Moscicki AB et al. Risks for incident human papillomavirus infection and low-grade squamous intraepithelial lesion development in young females. *JAMA*, 2001, 285:2995-3002.
60. Nsofor BI, Bello CS, Ekwempu CC. Sexually transmitted disease among women attending a family planning clinic in Zaria, Nigeria. *International Journal of Gynaecology & Obstetrics*, 1989, 28:365-7.
61. Ruijs GJ et al. Direct immunofluorescence for Chlamydia trachomatis on urogenital smears for epidemiological purposes. *European Journal of Obstetrics, Gynecology & Reproductive Biology*, 1988, 27:289-97.
62. Aklilu M et al. Factors associated with HIV-1 infection among sex workers of Addis Ababa, Ethiopia. *AIDS*, 2001, 15:87-96.
63. Allen S et al. Pregnancy and contraception use among urban Rwandan women after HIV testing and counseling. *American Journal of Public Health*, 1993, 83:705-10.
64. Carael M et al. Human immunodeficiency virus transmission among heterosexual couples in Central Africa. *AIDS*, 1988, 2:201-5.
65. Cohen E, Navaline H, Metzger D. High-risk behaviors for HIV: a comparison between crack-abusing and opioid-abusing African-American women. *Journal of Psychoactive Drugs*, 1994, 26:233-41.
66. Kapiga SH et al. Risk factors for HIV infection among women in Dar-es-Salaam, Tanzania. *Journal of Acquired Immune Deficiency Syndromes*, 1994, 7:301-9.
67. Kapiga SH et al. The incidence of HIV infection among women using family planning methods in Dar es Salaam, Tanzania. *AIDS*, 1998, 12:75-84.
68. Kiddugavu M et al. Hormonal contraceptive use and HIV-1 infection in a population-based cohort in Rakai, Uganda. *AIDS*, 2003, 17:233-40.
69. Limpakarnjanarat K et al. HIV-1 and other sexually transmitted infections in a cohort of female sex workers in Chiang Rai, Thailand. *Sexually Transmitted Infections*, 1999, 75:30-5.
70. Martin HL, Jr. et al. Hormonal contraception, sexually transmitted diseases, and risk of heterosexual transmission of human immunodeficiency virus type 1. *Journal of Infectious Diseases*, 1998, 178:1053-9.
71. Mati JK et al. Contraceptive use and the risk of HIV infection in Nairobi, Kenya. *International Journal of Gynaecology & Obstetrics*, 1995, 48:61-7.
72. Nagachinta T et al. Risk factors for HIV-1 transmission from HIV-seropositive male blood donors to their regular female partners in northern Thailand. *AIDS*, 1997, 11:1765-72.
73. Nzila N et al. HIV and other sexually transmitted diseases among female prostitutes in Kinshasa. *AIDS*, 1991, 5:715-21.
74. Plourde PJ et al. Human immunodeficiency virus type 1 infection in women attending a sexually transmitted diseases clinic in Kenya. *Journal of Infectious Diseases*, 1992, 166:86-92.
75. Rehle T et al. Risk factors of HIV-1 infection among female prostitutes in Khon Kaen, Northeast Thailand. *Infection*, 1992, 20:328-31.
76. Siraprasiri T et al. Risk factors for HIV among prostitutes in Chiangmai, Thailand. *AIDS*, 1991, 5:579-82.
77. Taneepanichskul S, Phuapradit W, Chaturachinda K. Association of contraceptives and HIV-1 infection in Thai female commercial sex workers. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 1997, 37:86-8.

78. Ungchusak K et al. Determinants of HIV infection among female commercial sex workers in northeastern Thailand: results from a longitudinal study.[erratum appears in *J Acquir Immune Defic Syndr Hum Retrovirol* 1998 Jun 1;18(2):192]. *Journal of Acquired Immune Deficiency Syndromes & Human Retrovirology*, 1996, 12:500-7.
79. McClelland RS et al. A prospective study of hormonal contraceptive use and cervical shedding of herpes simplex virus in human immunodeficiency virus type 1-seropositive women. *Journal of Infectious Diseases*, 2002, 185:1822-5.
80. Mostad SB et al. Hormonal contraception, vitamin A deficiency, and other risk factors for shedding of HIV-1 infected cells from the cervix and vagina. *Lancet*, 1997, 350:922-7.
81. Mostad SB et al. Cervical shedding of herpes simplex virus in human immunodeficiency virus-infected women: effects of hormonal contraception, pregnancy, and vitamin A deficiency. *Journal of Infectious Diseases*, 2000, 181:58-63.
82. Tagy AH et al. The effect of low-dose combined oral contraceptive pills versus injectable contraceptive (Depot Provera) on liver function tests of women with compensated bilharzial liver fibrosis. *Contraception*, 2001, 64:173-6.
83. Adadevoh BK, Isaacs WA. The effect of megestrol acetate on sickling. *American Journal of the Medical Sciences*, 1973, 265:367-70.
84. Barbosa IC et al. Carbohydrate Metabolism in Sickle Cell Patients Using a Subdermal Implant Containing Nomegestrol Acetate (Uniplant). *Contraception*, 2001, 63:263-5.
85. de Abood M et al. Effects of Depo-Provera or Microgynon on the Painful Crises of Sickle Cell Anemia Patients. *Contraception*, 1997, 56:313-6.
86. De Ceulaer K et al. Medroxyprogesterone acetate and homozygous sickle-cell disease. *Lancet*, 1982, 2:229-31.
87. Howard RJ, Lillis C, Tuck SM. Contraceptives, Counselling, and Pregnancy in Women with Sickle Cell Disease. *BMJ*, 1993, 306:1735-7.
88. Ladipo OA et al. Norplant use by women with sickle cell disease. *International Journal of Gynaecology and Obstetrics*, 1993, 41:85-7.
89. Nascimento ML, Ladipo OA, Coutinho E. Nomegestrol acetate contraceptive implant use by women with sickle cell disease. *Clinical Pharmacology and Therapeutics*, 1998, 64:433-8.
90. Yoong WC, Tuck SM, Yardumian A. Red cell deformability in oral contraceptive pill users with sickle cell anaemia. *British Journal of Haematology*, 1999, 104:868-70.
91. Haukkamaa M. Contraception by Norplant subdermal capsules is not reliable in epileptic patients on anticonvulsant treatment. *Contraception*, 1986, 33:559-65.
92. Odland V, Olsson SE. Enhanced metabolism of levonorgestrel during phenytoin treatment in a woman with Norplant implants. *Contraception*, 1986, 33:257-61.
93. Shane-McWhorter L et al. Enhanced metabolism of levonorgestrel during phenobarbital treatment and resultant pregnancy. *Pharmacotherapy*, 1998, 18:1360-4.

EMERGENCY CONTRACEPTIVE PILLS

EMERGENCY CONTRACEPTION PILLS (ECPs) (including levonorgestrel contraceptive pills and combined oral contraceptive pills)	ECPs do not protect against STI/HIV. If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
PREGNANCY	NA	Clarification: Although this method is not indicated for a woman with a known or suspected pregnancy, there is no known harm to the woman, the course of her pregnancy, or the fetus if ECPs are accidentally used.
BREASTFEEDING	1	
HISTORY OF ECTOPIC PREGNANCY	1	
HISTORY OF SEVERE CARDIOVASCULAR COMPLICATIONS* (ischaemic heart disease, cerebrovascular attack, or other thromboembolic conditions)	2	
ANGINA PECTORIS*	2	
MIGRAINE*	2	
SEVERE LIVER DISEASE (including jaundice)*	2	
REPEATED ECP USE	1	Clarification: Recurrent ECP use is an indication that the woman requires further counselling on other contraceptive options. Frequently repeated ECP use may be harmful for women with conditions classified as 2, 3 or 4 for COC, CIC or POC use.
RAPE*	1	

* Additional comments

HISTORY OF SEVERE CARDIOVASCULAR COMPLICATIONS

The duration of use of ECPs is less than that of regular use of COCs or POPs and thus would be expected to have less clinical impact.

ANGINA PECTORIS

The duration of use of ECPs is less than that of regular use of COCs or POPs and thus would be expected to have less clinical impact.

MIGRAINE

The duration of use of ECPs is less than that of regular use of COCs or POPs and thus would be expected to have less clinical impact.

SEVERE LIVER DISEASE (including jaundice)

The duration of use of ECPs is less than that of regular use of COCs or POPs and thus would be expected to have less clinical impact.

RAPE

There are no restrictions for use of ECPs in cases of rape.

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	1
Pregnancy.....	1
Age	1
Parity.....	1
Postpartum	1
Post-abortion.....	2
Past ectopic pregnancy.....	2
History of pelvic surgery.....	2
Smoking.....	2
Obesity.....	2
Blood pressure measurement unavailable	2
CARDIOVASCULAR DISEASE.....	3
Multiple risk factors for arterial cardiovascular disease.....	3
Hypertension.....	3
History of high blood pressure during pregnancy.....	3
Deep venous thromboembolism (DVT)/pulmonary embolism (PE).....	4
Known thrombogenic mutations.....	4
Superficial venous thrombosis	4
Current and history of ischaemic heart disease	4
Stroke	4
Known hyperlipidaemias	5
Valvular heart disease	5
NEUROLOGIC CONDITIONS.....	5
Headaches.....	5
Epilepsy	5
DEPRESSIVE DISORDERS	5
Depressive disorders	5
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS.....	6
Vaginal bleeding patterns.....	6
Unexplained vaginal bleeding	6
Endometriosis	6
Benign ovarian tumours	6
Severe dysmenorrhoea.....	6
Trophoblast disease.....	6
Cervical ectropion	6
Cervical intraepithelial neoplasia (CIN)	6
Cervical cancer	6
Breast disease	7
Endometrial cancer	7
Ovarian cancer.....	7
Uterine fibroids.....	7
Anatomical abnormalities.....	7
Pelvic inflammatory disease (PID)	8
STIs	8
HIV/AIDS	9
High risk of HIV	9
HIV-infected	9
AIDS	9

OTHER INFECTIONS	9
Schistosomiasis	9
Tuberculosis.....	9
Malaria	10
ENDOCRINE CONDITIONS.....	10
Diabetes.....	10
Thyroid disorders	10
GASTROINTESTINAL CONDITIONS.....	10
Gall-bladder disease	10
History of cholestasis	10
Viral hepatitis	11
Cirrhosis.....	11
Liver tumours	11
ANAEMIAS.....	11
Thalassaemia.....	11
Sickle cell disease.....	11
Iron-deficiency anaemia	11
DRUG INTERACTIONS	11
Drugs which affect liver enzymes.....	11
Antibiotics	11
Antiretroviral therapy.....	12
Additional comments	13
References for intrauterine devices.....	15

INTRAUTERINE DEVICES

Cu-IUD = Copper-bearing IUD

LNG-IUD = Levonorgestrel-releasing IUD (20 g/24hours)

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY			
PREGNANCY	4	4	Clarification: The IUD is not indicated during pregnancy and should not be used because of the risk of serious pelvic infection and septic spontaneous abortion.
AGE*			
a) Menarche to < 20 years	2	2	
b) ≥ 20 years	1	1	
PARITY*			
a) Nulliparous	2	2	Evidence: There are conflicting data regarding whether IUD use is associated with infertility among nulliparous women, although recent, well-conducted studies suggest no increased risk. ¹⁻⁹
b) Parous	1	1	
POSTPARTUM* (breastfeeding or non-breastfeeding, including post-caesarean section)			
a) < 48 hours	2	3	Evidence: There was some increase in expulsion rates with delayed postpartum insertion compared to immediate insertion and with immediate postpartum insertion compared to interval insertion. ¹⁰⁻¹⁶
b) 48 hours to < 4 weeks	3	3	
c) ≥ 4 weeks	1	1	
d) Puerperal sepsis	4	4	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
POST-ABORTION*			
a) First trimester	1	1	<p>Clarification: IUDs can be inserted immediately after first-trimester, spontaneous or induced abortion.</p> <p>Evidence: There was no difference in risk of complications for immediate versus delayed insertion of an IUD after abortion. Expulsion was greater when an IUD was inserted following a second-trimester abortion versus following a first-trimester abortion. There were no differences in safety or expulsions for post-abortion insertion of an LNG-IUD compared with Cu-IUD.¹⁷⁻³⁰</p>
b) Second trimester	2	2	
c) Immediate post-septic abortion	4	4	
PAST ECTOPIC PREGNANCY*	1	1	
HISTORY OF PELVIC SURGERY (see postpartum, including caesarean section)	1	1	
SMOKING			
a) Age < 35 years	1	1	
b) Age ≥ 35 years			
(i) < 15 cigarettes/day	1	1	
(ii) ≥ 15 cigarettes/day	1	1	
OBESITY ≥ 30 kg/m ² body mass index (BMI)	1	1	
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA	NA	Clarification: While a blood pressure measurement may be appropriate for good preventative health care, it is not materially related to safe and effective IUD use. Women should not be denied use of IUDs simply because their blood pressure cannot be measured.

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
CARDIOVASCULAR DISEASE			
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	1	2	
HYPERTENSION*			
For all categories of hypertension, classifications are based on the assumption that no other risk factors for cardiovascular disease exist. When multiple risk factors do exist, risk of cardiovascular disease may increase substantially. A single reading of blood pressure level is not sufficient to classify a woman as hypertensive.			
a) History of hypertension where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	1	2	
b) Adequately controlled hypertension where blood pressure CAN be evaluated	1	1	
c) Elevated blood pressure levels (properly taken measurements)			
(i) systolic 140-159 or diastolic 90-99	1	1	
(ii) systolic \geq 160 or diastolic \geq 100	1	2	
d) Vascular disease	1	2	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	1	1	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
DEEP VENOUS THROMBOEMBOLISM (DVT)/PULMONARY EMBOLISM (PE)*			
a) History of DVT/PE	1	2	
b) Current DVT/PE	1	3	
c) Family history of DVT/PE (first-degree relatives)	1	1	
d) Major surgery			
(i) with prolonged immobilization	1	2	
(ii) without prolonged immobilization	1	1	
e) Minor surgery without immobilization	1	1	
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	1	2	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
SUPERFICIAL VENOUS THROMBOSIS			
a) Varicose veins	1	1	
b) Superficial thrombophlebitis	1	1	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE*		I C	
	1	2 3	
STROKE* (history of cerebrovascular accident)	1	2	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
KNOWN HYPERLIPIDAEMIAS	1	2	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
VALVULAR HEART DISEASE a) Uncomplicated b) Complicated (pulmonary hypertension, risk of atrial fibrillation, history of subacute bacterial endocarditis)	1 2	1 2	Clarification: Prophylactic antibiotics to prevent endocarditis are advised for insertion.
NEUROLOGIC CONDITIONS			
HEADACHES*		I C	
a) Non-migrainous (mild or severe)	1	1 1	Clarification: Any new headaches or marked changes in headaches should be evaluated.
b) Migraine			
(i) without aura			
Age < 35	1	2 2	
Age ≥ 35	1	2 2	
(ii) with aura, at any age	1	2 3	
EPILEPSY	1	1	
DEPRESSIVE DISORDERS			
DEPRESSIVE DISORDERS	1	1	Clarification: The classification is based on data for women with selected depressive disorders. No data on bipolar disorder or postpartum depression were available. There is a potential for drug interactions between certain antidepressant medications and hormonal contraceptives.

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE	
	Cu-IUD	LNG-IUD		
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS				
VAGINAL BLEEDING PATTERNS*		I	C	
a) Irregular pattern <i>without</i> heavy bleeding	1	1	1	
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	2	1	2	Clarification: Unusually heavy bleeding should raise the suspicion of a serious underlying condition. Evidence: Among women with heavy or prolonged bleeding, LNG-IUDs were beneficial in treating menorrhagia. ³¹⁻³⁵
UNEXPLAINED VAGINAL BLEEDING (suspicion for serious condition)	I	C	I	C
Before evaluation	4	2	4	2
ENDOMETRIOSIS*	2	1		
BENIGN OVARIAN TUMOURS (including cysts)	1	1		
SEVERE DYSMENORRHOEA*	2	1		
TROPHOBLAST DISEASE*				
a) Benign gestational trophoblastic disease	3	3		
b) Malignant gestational trophoblastic disease	4	4		
CERVICAL ECTROPION	1	1		
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)*	1	2		
CERVICAL CANCER* (awaiting treatment)	I	C	I	C
	4	2	4	2

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE	
	Cu-IUD	LNG-IUD		
BREAST DISEASE* a) Undiagnosed mass b) Benign breast disease c) Family history of cancer d) Breast cancer: (i) current (ii) past and no evidence of current disease for 5 years	1 1 1 1 1	2 1 1 4 3		
ENDOMETRIAL CANCER*	I C 4 2	I C 4 2		
OVARIAN CANCER*	3 2	3 2		
UTERINE FIBROIDS* a) Without distortion of the uterine cavity b) With distortion of the uterine cavity	1 4	1 4	Evidence: Among women with fibroids, there were no adverse health events with LNG-IUD use and there was a decrease in symptoms and size of fibroids for some women. ³⁸⁻⁴⁴	
ANATOMICAL ABNORMALITIES* a) Distorted uterine cavity (any congenital or acquired uterine abnormality distorting the uterine cavity in a manner that is incompatible with IUD insertion) b) Other abnormalities (including cervical stenosis or cervical lacerations) not distorting the uterine cavity or interfering with IUD insertion	4 2	4 2		

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.																			
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE																	
	Cu-IUD	LNG-IUD																		
PELVIC INFLAMMATORY DISEASE (PID)* a) Past PID (assuming no known current risk factors for STIs) (i) with subsequent pregnancy (ii) without subsequent pregnancy b) PID - current	<table border="1"> <tr> <td>I</td> <td>C</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>2</td> <td>2</td> </tr> <tr> <td>4</td> <td>2</td> </tr> </table>	I	C	1	1	2	2	4	2	<table border="1"> <tr> <td>I</td> <td>C</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>2</td> <td>2</td> </tr> <tr> <td>4</td> <td>2</td> </tr> </table>	I	C	1	1	2	2	4	2	<p>Clarification for continuation: Treat the PID using appropriate antibiotics. There is usually no need for removal of the IUD if the client wishes to continue its use. (See <i>Selected Practice Recommendations for Contraceptive Use</i>. WHO: Geneva, 2002). Continued use of an IUD depends on the woman's informed choice and her current risk factors for STIs and PID.</p> <p>Evidence: Among IUD users treated for PID, there was no difference in clinical course if the IUD was removed or left in place.⁴⁵⁻⁴⁷</p>	
I	C																			
1	1																			
2	2																			
4	2																			
I	C																			
1	1																			
2	2																			
4	2																			
STIs* a) Current purulent cervicitis or chlamydial infection or gonorrhoea b) Other STIs (excluding HIV and hepatitis) c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	<table border="1"> <tr> <td>I</td> <td>C</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>2</td> <td>2</td> </tr> <tr> <td>2</td> <td>2</td> </tr> </table>	I	C	4	2	2	2	2	2	<table border="1"> <tr> <td>I</td> <td>C</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>2</td> <td>2</td> </tr> <tr> <td>2</td> <td>2</td> </tr> </table>	I	C	4	2	2	2	2	2	<p>Clarification for continuation: Treat the STI using appropriate antibiotics. There is usually no need for removal of the IUD if the client wishes to continue its use. Continued use of an IUD depends on the woman's informed choice and her current risk factors for STIs and PID.</p> <p>Evidence: There is no evidence regarding whether IUD insertion among women with STIs increases the risk of PID compared with no IUD insertion. Among women who have an IUD inserted, the absolute risk of subsequent PID was low among women with STI at the time of insertion but greater than among women with no STI at the time of IUD insertion.⁴⁸⁻⁵⁴</p>	
I	C																			
4	2																			
2	2																			
2	2																			
I	C																			
4	2																			
2	2																			
2	2																			

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.				
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE		
	Cu-IUD	LNG-IUD			
STIs (Cont'd)					
d) Increased risk of STIs	2/3	2	2/3	2	<p>Clarification for initiation: If a woman has a very high individual likelihood of exposure to gonorrhoea or chlamydial infection, the condition is a Category 3.</p> <p>Evidence: Using an algorithm to classify STI risk status among IUD users, one study reported that 11% of high STI-risk women experienced IUD-related complications compared with 5% of those not classified as high risk.⁵⁰</p>
HIV/AIDS					
HIGH RISK OF HIV*	I	C	I	C	
	2	2	2	2	Evidence: Among women at risk of HIV, copper IUD use did not increase risk of HIV acquisition. ⁵⁵⁻⁶⁵
HIV-INFECTED	2	2	2	2	Evidence: Among IUD users, there is limited evidence showing no increased risk of overall complications or infection-related complications when comparing HIV-infected women with non-infected women. Furthermore, IUD use among HIV-infected women was not associated with increased risk of transmission to sexual partners. ^{55, 66-69}
AIDS	3	2	3	2	Clarification for continuation: IUD users with AIDS should be closely monitored for pelvic infection.
Clinically well on ARV therapy	2	2	2	2	
OTHER INFECTIONS					
SCHISTOSOMIASIS					
a) Uncomplicated	1		1		
b) Fibrosis of the liver (if severe, see cirrhosis)	1		1		
TUBERCULOSIS*					
	I	C	I	C	
a) Non-pelvic	1	1	1	1	
b) Known pelvic	4	3	4	3	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
MALARIA	1	1	
ENDOCRINE CONDITIONS			
DIABETES*			
a) History of gestational disease	1	1	
b) Non-vascular disease			
(i) non-insulin dependent	1	2	
(ii) insulin dependent	1	2	
c) Nephropathy/ retinopathy/ neuropathy	1	2	
d) Other vascular disease or diabetes of >20 years' duration	1	2	
THYROID DISORDERS			
a) Simple goitre	1	1	
b) Hyperthyroid	1	1	
c) Hypothyroid	1	1	
GASTROINTESTINAL CONDITIONS			
GALL-BLADDER DISEASE			
a) Symptomatic			
(i) treated by cholecystectomy	1	2	
(ii) medically treated	1	2	
(iii) current	1	2	
b) Asymptomatic	1	2	
HISTORY OF CHOLESTASIS*			
a) Pregnancy-related	1	1	
b) Past COC-related	1	2	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE
	Cu-IUD	LNG-IUD	
VIRAL HEPATITIS*			
a) Active	1	3	
b) Carrier	1	1	
CIRRHOSIS*			
a) Mild (compensated)	1	2	
b) Severe (decompensated)	1	3	
LIVER TUMOURS*			
a) Benign (adenoma)	1	3	
b) Malignant (hepatoma)	1	3	
ANAEMIAS			
THALASSAEMIA*	2	1	
SICKLE CELL DISEASE*	2	1	
IRON-DEFICIENCY ANAEMIA*	2	1	
DRUG INTERACTIONS			
DRUGS WHICH AFFECT LIVER ENZYMES			
a) Rifampicin	1	1	Evidence: One study found that rifabutin, which is in the same class of drugs as rifampicin, has no impact on the effectiveness of LNG-IUD. ⁷⁰
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	1	1	
ANTIBIOTICS (excluding rifampicin)	1	1	
a) Griseofulvin	1	1	
b) Other antibiotics	1	1	

* See also additional comments at end of table

INTRAUTERINE DEVICES (IUDs)	IUDs do not protect against STI/HIV. If there is risk of STI/HIV (including the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.				
CONDITION	CATEGORY I=Initiation, C=Continuation		CLARIFICATIONS/EVIDENCE		
	Cu-IUD	LNG-IUD			
ANTIRETROVIRAL THERAPY	I	C	I	C	
	2/3	2	2/3	2	Clarification: There is no known drug interaction between ARV therapy and IUD use. However, AIDS as a condition is classified as Category 3 for insertion and Category 2 for continuation unless the woman is clinically well on ARV therapy in which case, both insertion and continuation are classified as Category 2. (See AIDS condition above.)

* See also additional comments at end of table

Additional comments

AGE

Menarche to < 20 years: There is concern both about the risk of expulsion due to nulliparity and risk of STIs due to sexual behaviour in younger age groups.

PARITY

Nulliparous: Nulliparity is related to an increased risk of expulsion.

POSTPARTUM

< 48 hours, 48 hours to < 4 weeks, ≥ 4 weeks: Concern that the neonate may be at risk due to exposure to steroid hormones with LNG-IUD use during the first 6 weeks postpartum is the same as for other POCs.

Puerperal sepsis: Insertion of an IUD may substantially worsen the condition.

POST-ABORTION

Immediate post-septic abortion: Insertion of an IUD may substantially worsen the condition.

PAST ECTOPIC PREGNANCY

The absolute risk of ectopic pregnancy is extremely low due to the high effectiveness of IUDs. However, when a woman becomes pregnant during IUD use, the relative likelihood of ectopic pregnancy is greatly increased.

HYPERTENSION

There is theoretical concern about the effect of LNG on lipids. There is no restriction for copper IUDs.

DEEP VEIN THROMBOEMBOLISM (DVT)/ PULMONARY EMBOLISM (PE)

Some progestogens may increase the risk of venous thrombosis, although this increase is substantially less than for COCs.

CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE

There is theoretical concern about the effect of LNG on lipids. There is no restriction for copper IUDs.

STROKE

There is theoretical concern about the effect of LNG on lipids. There is no restriction for copper IUDs.

HEADACHES

Aura is a specific focal neurologic symptom. For more information on this and other diagnostic criteria, see: Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders, 2nd Edition. Cephalalgia. 2004; 24 (Suppl 1): 1- 150.

http://216.25.100.131/ihscommon/guidelines/pdfs/ihc_II_main_no_print.pdf

VAGINAL BLEEDING PATTERNS

LNG-IUD use frequently causes changes in menstrual bleeding patterns. Over time, LNG-IUD users are more likely than non-users to become amenorrhoeic, thus LNG-IUDs are sometimes used as a treatment to correct heavy bleeding.

ENDOMETRIOSIS

Copper IUD use may worsen dysmenorrhoea associated with the condition.

SEVERE DYSMENORRHOEA

Dysmenorrhoea may intensify with copper IUD use. LNG-IUD use has been associated with reduction of dysmenorrhoea.

TROPHOBLAST DISEASE

There is an increased risk of perforation since the treatment for the condition may require multiple uterine curettages.

CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)

There is some theoretical concern that LNG-IUDs may enhance progression of CIN.

CERVICAL CANCER (awaiting treatment)

There is concern about the increased risk of infection and bleeding at insertion. The IUD will likely need to be removed at the time of treatment but, until then, the woman is at risk of pregnancy.

BREAST DISEASE

Breast cancer: Breast cancer is a hormonally sensitive tumour. Concerns about progression of the disease may be less with LNG-IUDs than with COCs or higher-dose POCs.

ENDOMETRIAL CANCER

There is concern about the increased risk of infection, perforation and bleeding at insertion. The IUD will likely need to be removed at the time of treatment but, until then, the woman is at risk of pregnancy.

OVARIAN CANCER

The IUD will likely need to be removed at the time of treatment but, until then, the woman is at risk of pregnancy.

UTERINE FIBROIDS

Without distortion of the uterine cavity: Women with heavy or prolonged bleeding should be assigned the category for that condition.

With distortion of the uterine cavity: Pre-existing uterine fibroids that distort the uterine cavity may be incompatible with insertion and proper placement of the IUD.

ANATOMICAL ABNORMALITIES

Distorted uterine cavity: In the presence of an anatomic abnormality that distorts the uterine cavity, proper IUD placement may not be possible.

PELVIC INFLAMMATORY DISEASE (PID)

IUDs do not protect against STI/HIV/PID. In women at low risk of STIs, IUD insertion poses little risk of PID. Current risk of STIs and desire for future pregnancy are relevant considerations.

STIs

IUDs do not protect against STI/HIV/PID. Among women with chlamydial infection or gonorrhoea, the potential increased risk of PID with IUD insertions should be avoided. The concern is less for other STIs.

HIGH RISK OF HIV

IUDs do not protect against STI/HIV/PID.

TUBERCULOSIS

Known pelvic: Insertion of an IUD may substantially worsen known the condition.

DIABETES

Whether the amount of LNG released by the IUD may slightly influence carbohydrate and lipid metabolism is unclear. Some progestogens may increase the risk of thrombosis, although this increase is substantially less than for COCs.

HISTORY OF CHOLESTASIS

There is concern that a history of COC-related cholestasis may predict subsequent cholestasis with LNG use. Whether there is any risk with use of an LNG-IUD is unclear.

VIRAL HEPATITIS

Active: POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. This concern is similar to, but less than, that with COCs.

CIRRHOSIS

POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. This concern is similar to, but less than, that with COCs.

LIVER TUMOURS

POCs are metabolized by the liver and their use may adversely affect women whose liver function is compromised. In addition, POC use may enhance the growth of tumours. This concern is similar to, but less than, that with COCs.

THALASSAEMIA

There is concern about an increased risk of blood loss with copper IUDs.

SICKLE CELL DISEASE

There is concern about an increased risk of blood loss with copper IUDs.

IRON-DEFICIENCY ANAEMIA

There is concern about an increased risk of blood loss with copper IUDs.

References for intrauterine devices

1. Cramer DW et al. Tubal infertility and the intrauterine device. *New England Journal of Medicine*, 1985, 312:941-7.
2. Daling JR et al. Primary tubal infertility in relation to the use of an intrauterine device. *New England Journal of Medicine*, 1985, 312:937-41.
3. Daling JR et al. The intrauterine device and primary tubal infertility. *New England Journal of Medicine*, 1992, 326:203-4.
4. Delborge W et al. Return to fertility in nulliparous and parous women after removal of the GyneFix intrauterine contraceptive system. *European Journal of Contraception & Reproductive Health Care*, 2002, 7:24-30.
5. Doll H, Vessey M, Painter R. Return of fertility in nulliparous women after discontinuation of the intrauterine device: comparison with women discontinuing other methods of contraception. *BJOG: an International Journal of Obstetrics & Gynaecology*, 2001, 108:304-14.
6. Hubacher D et al. Use of copper intrauterine devices and the risk of tubal infertility among nulligravid women. *New England Journal of Medicine*, 2001, 345:561-7.
7. Skjeldestad FE, Bratt H. Return of fertility after use of IUDs (Nova-T, MLCu250 and MLCu375). *Advances in Contraception*, 1987, 3:139-45.
8. Urbach DR et al. Association of perforation of the appendix with female tubal infertility. *American Journal of Epidemiology*, 2001, 153:566-71.
9. Wilson JC. A prospective New Zealand study of fertility after removal of copper intrauterine contraceptive devices for conception and because of complications: a four-year study. *American Journal of Obstetrics & Gynecology*, 1989, 160:391-6.
10. Brenner PF. A clinical trial of the Delta-T intrauterine device: immediate postpartum insertion. *Contraception*, 1983, 28:135-47.
11. Chi IC, Wilkens L, Rogers S. Expulsions in immediate postpartum insertions of Lippes Loop D and Copper T IUDs and their counterpart Delta devices--an epidemiological analysis. *Contraception*, 1985, 32:119-34.
12. El-Shafei M. Postpartum and postabortion intrauterine device insertion unmet needs of safe reproductive health: three year experience of Mansoura University Hospital. *European Journal of Contraception & Reproductive Health Care*, 2000, 26:253-62.
13. Grimes D et al. Immediate post-partum insertion of intrauterine devices.[update of *Cochrane Database Systematic Reviews*, 2001, CD003036; PMID: 11406064]. [Review] [30 refs]. *Cochrane Database of Systematic Reviews*, 2003, CD003036.
14. Morrison C et al. Clinical outcomes of two early postpartum IUD insertion programs in Africa. *Contraception*, 1996, 53:17-21.
15. Thiery M et al. The ML Cu250; clinical experience in Belgium and The Netherlands. *British Journal of Obstetrics & Gynaecology*, 1982, 89:51-3.
16. Thiery M, Delbeke L, Van Kets H. Comparative performance of two copper-wired IUDs (ML Cu 250 and T Cu 200): immediate postpartum and interval insertion. *Advances in Contraceptive Delivery Systems*, 1980, 1:27-35.
17. The World Health Organization's Special Programme of Research, Development and Research Training in Human Reproduction. Task Force on Intrauterine Devices for Fertility Regulation. The Alza T IPCS 52, a longer acting progesterone IUD: safety and efficacy compared to the TCu220C and multiload 250 in two randomized multicentre trials. *Clinical Reproduction & Fertility*, 1983, 2:113-28.
18. The World Health Organization's Special Programme of Research, Development and Research Training in Human Reproduction. Task Force on Intrauterine Devices for Fertility Regulation. IUD insertion following termination of pregnancy: a clinical trial of the TCu 220C, Lippes loop D, and copper 7. *Studies in Family Planning*, 1983, 14:99-108.
19. The World Health Organization's Special Programme of Research, Development and Research Training in Human Reproduction. Task Force on Intrauterine Devices for Fertility Regulation. IUD insertion following spontaneous abortion: a clinical trial of the TCu 220C, Lippes loop D, and copper 7. *Studies in Family Planning*, 1983, 14:109-14.
20. El Tagy A et al. Safety and acceptability of post-abortion IUD insertion and the importance of counseling. *Contraception*, 2003, 67:229-34.
21. Gillett PG et al. A comparison of the efficacy and acceptability of the Copper-7 intrauterine device following immediate or delayed insertion after first-trimester therapeutic abortion. *Fertility & Sterility*, 1980, 34:121-4.

22. Grimes D, Schulz K, Stanwood N. Immediate postabortal insertion of intrauterine devices.[update of Cochrane Database Syst Rev. 2000;(2):CD001777; PMID: 10796820]. [Review] [28 refs]. *Cochrane Database of Systematic Reviews*, 2002, CD001777.
23. Gupta I, Devi PK. Studies on immediate post-abortion copper "T" device. *Indian Journal of Medical Research*, 1975, 63:736-9.
24. Moussa A. Evaluation of postabortion IUD insertion in Egyptian women. *Contraception*, 2001, 63:315-7.
25. Pakarinen P, Toivonen J, Luukkainen T. Randomized comparison of levonorgestrel- and copper-releasing intrauterine systems immediately after abortion, with 5 years' follow-up. *Contraception*, 2003, 68:31-4.
26. Stanwood NL, Grimes DA, Schulz KF. Insertion of an intrauterine contraceptive device after induced or spontaneous abortion: a review of the evidence. *BJOG: an International Journal of Obstetrics & Gynaecology*, 2001, 108:1168-73.
27. Suvisaari J, Lahteenmaki P. Detailed analysis of menstrual bleeding patterns after postmenstrual and postabortal insertion of a copper IUD or a levonorgestrel-releasing intrauterine system. *Contraception*, 1996, 54:201-8.
28. Timonen H, Luukkainen T. Immediate postabortion insertion of the copper-T (TCu-200) with eighteen months follow-up. *Contraception*, 1974, 9:153-60.
29. Tuveng JM, Skjeldestad FE, Iversen T. Postabortal insertion of IUD. *Advances in Contraception*, 1986, 2:387-92.
30. Zhang PZ. Five years experience with the copper T 200 in Shanghai - 856 cases. *Contraception*, 1980, 22:561-71.
31. Barrington JW, Arunkalaivanan AS, Abdel-Fattah. Comparison between the levonorgestrel intrauterine system (LNG-IUS) and thermal balloon ablation in the treatment of menorrhagia. *European Journal of Obstetrics, Gynecology & Reproductive Biology*, 2003, 108:72-4.
32. Hurskainen R et al. Quality of life and cost-effectiveness of levonorgestrel-releasing intrauterine system versus hysterectomy for treatment of menorrhagia: a randomised trial. *Lancet*, 2001, 357:273-7.
33. Istre O, Trolle B. Treatment of menorrhagia with the levonorgestrel intrauterine system versus endometrial resection. *Fertility & Sterility*, 2001, 76:304-9.
34. Lethaby AE, Cooke I, Rees M. Progesterone/progestogen releasing intrauterine systems versus either placebo or any other medication for heavy menstrual bleeding. *Cochrane Database of Systematic Reviews*, 2000, CD002126.
35. Stewart A et al. The effectiveness of the levonorgestrel-releasing intrauterine system in menorrhagia: a systematic review. *BJOG: an International Journal of Obstetrics & Gynaecology*, 2001, 108:74-86.
36. Fedele L et al. Use of a levonorgestrel-releasing intrauterine device in the treatment of rectovaginal endometriosis. *Fertility & Sterility*, 2001, 75:485-8.
37. Vercellini P et al. A levonorgestrel-releasing intrauterine system for the treatment of dysmenorrhea associated with endometriosis: a pilot study. *Fertility & Sterility*, 1999, 72:505-8.
38. Fedele L et al. Treatment of adenomyosis-associated menorrhagia with a levonorgestrel-releasing intrauterine device. *Fertility & Sterility*, 1997, 68:426-9.
39. Grigorieva V et al. Use of a levonorgestrel-releasing intrauterine system to treat bleeding related to uterine leiomyomas. *Fertility & Sterility*, 2003, 79:1194-8.
40. Mercorio F et al. The effect of a levonorgestrel-releasing intrauterine device in the treatment of myoma-related menorrhagia. *Contraception*, 2003, 67:277-80.
41. Wildemeersch D, Schacht E, Wildemeersch P. Treatment of primary and secondary dysmenorrhea with a novel 'frameless' intrauterine levonorgestrel-releasing drug delivery system: a pilot study. *European Journal of Contraception & Reproductive Health Care*, 2001, 6:192-8.
42. Wildemeersch D, Schacht E. The effect on menstrual blood loss in women with uterine fibroids of a novel "frameless" intrauterine levonorgestrel-releasing drug delivery system: a pilot study. *European Journal of Obstetrics, Gynecology & Reproductive Biology*, 2002, 102:74-9.
43. Wildemeersch D, Schacht E, Wildemeersch P. Contraception and treatment in the perimenopause with a novel "frameless" intrauterine levonorgestrel-releasing drug delivery system: an extended pilot study. *Contraception*, 2002, 66:93-99.
44. Wildemeersch D, Schacht E, Wildemeersch P. Performance and acceptability of intrauterine release of levonorgestrel with a miniature delivery system for hormonal substitution therapy, contraception and treatment in peri and postmenopausal women. *Maturitas*, 2003, 44:237-45.

45. Larsson B, Wennergren M. Investigation of a copper-intrauterine device (Cu-IUD) for possible effect on frequency and healing of pelvic inflammatory disease. *Contraception*, 1977, 15:143-9.
46. Soderberg G, Lindgren S. Influence of an intrauterine device on the course of an acute salpingitis. *Contraception*, 1981, 24:137-43.
47. Teisala K. Removal of an intrauterine device and the treatment of acute pelvic inflammatory disease. *Annals of Medicine*, 1989, 21:63-5.
48. Faundes A et al. The risk of inadvertent intrauterine device insertion in women carriers of endocervical Chlamydia trachomatis. *Contraception*, 1998, 58:105-9.
49. Ferraz do Lago R et al. Follow-up of users of intrauterine device with and without bacterial vaginosis and other cervicovaginal infections. *Contraception*, 2003, 68:105-9.
50. Morrison CS et al. Use of sexually transmitted disease risk assessment algorithms for selection of intrauterine device candidates. *Contraception*, 1999, 59:97-106.
51. Pap-Akeson M et al. Genital tract infections associated with the intrauterine contraceptive device can be reduced by inserting the threads into the uterine cavity. *British Journal of Obstetrics & Gynaecology*, 1992, 99:676-9.
52. Sinei SK et al. Preventing IUCD-related pelvic infection: the efficacy of prophylactic doxycycline at insertion. *British Journal of Obstetrics & Gynaecology*, 1990, 97:412-9.
53. Skjeldestad FE et al. IUD users in Norway are at low risk for genital C. trachomatis infection. *Contraception*, 1996, 54:209-12.
54. Walsh TL et al. IUD Study Group. Effect of prophylactic antibiotics on morbidity associated with IUD insertion: results of a pilot randomized controlled trial. *Contraception*, 1994, 50:319-27.
55. European Study Group on Heterosexual Transmission of HIV. Comparison of female to male and male to female transmission of HIV in 563 stable couples. *BMJ*, 1992, 304:809-13.
56. Carael M et al. Human immunodeficiency virus transmission among heterosexual couples in Central Africa. *AIDS*, 1988, 2:201-5.
57. Kapiga SH et al. Risk factors for HIV infection among women in Dar-es-Salaam, Tanzania. *Journal of Acquired Immune Deficiency Syndromes*, 1994, 7:301-9.
58. Kapiga SH et al. The incidence of HIV infection among women using family planning methods in Dar es Salaam, Tanzania. *AIDS*, 1998, 12:75-84.
59. Mann JM et al. HIV infection and associated risk factors in female prostitutes in Kinshasa, Zaire. *AIDS*, 1988, 2:249-54.
60. Martin HL, Jr. et al. Hormonal contraception, sexually transmitted diseases, and risk of heterosexual transmission of human immunodeficiency virus type 1. *Journal of Infectious Diseases*, 1998, 178:1053-9.
61. Mati JK et al. Contraceptive use and the risk of HIV infection in Nairobi, Kenya. *International Journal of Gynaecology & Obstetrics*, 1995, 48:61-7.
62. Nicolosi A et al. Italian Study Group on HIV Heterosexual Transmission. The efficiency of male-to-female and female-to-male sexual transmission of the human immunodeficiency virus: a study of 730 stable couples. *Epidemiology*, 1994, 5:570-5.
63. Plourde PJ et al. Human immunodeficiency virus type 1 infection in women attending a sexually transmitted diseases clinic in Kenya. *Journal of Infectious Diseases*, 1992, 166:86-92.
64. Sinei SK et al. Contraceptive use and HIV infection in Kenyan family planning clinic attenders. *International Journal of STD & AIDS*, 1996, 7:65-70.
65. Spence MR et al. Seroprevalence of human immunodeficiency virus type I (HIV-1) antibodies in a family-planning population. *Sexually Transmitted Diseases*, 1991, 18:143-5.
66. Morrison CS et al. Is the intrauterine device appropriate contraception for HIV-1-infected women? *BJOG: an International Journal of Obstetrics & Gynaecology*, 2001, 108:784-90.
67. Sinei SK et al. Complications of use of intrauterine devices among HIV-1-infected women. *Lancet*, 1998, 351:1238-41.
68. Mostad SB et al. Hormonal contraception, vitamin A deficiency, and other risk factors for shedding of HIV-1 infected cells from the cervix and vagina. *Lancet*, 1997, 350:922-7.
69. Richardson BA et al. Effect of intrauterine device use on cervical shedding of HIV-1 DNA. *AIDS*, 1999, 13:2091-7.
70. Bounds W, Guillebaud J. Observational series on women using the contraceptive Mirena concurrently with anti-epileptic and other enzyme-inducing drugs. *Journal of Family Planning and Reproductive Health Care*, 2002, 28:78-80.

COPPER IUD FOR EMERGENCY CONTRACEPTION

This method is highly effective for preventing pregnancy. A copper-releasing IUD (Cu-IUD) can be used within 5 days of unprotected intercourse as an emergency contraceptive. However, when the time of ovulation can be estimated, the Cu-IUD can be inserted beyond 5 days after intercourse, if necessary, as long as the insertion does not occur more than 5 days after ovulation.

The eligibility criteria for interval Cu-IUD insertion also apply for the insertion of Cu-IUDs as emergency contraception.

COPPER IUD FOR EMERGENCY CONTRACEPTION	IUDs for emergency contraception do not protect against STI/HIV. If there is risk of STI/HIV, the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
PREGNANCY	4	Clarification: The IUD is not indicated during pregnancy and should not be used because of the risk of serious pelvic infection and septic spontaneous abortion.
RAPE*		
a) High risk of STI	3	
b) Low risk of STI	1	

*** Additional comments**

RAPE

IUDs do not protect against STI/HIV/PID. Among women with chlamydial infection or gonorrhoea, the potential increased risk of PID with IUD insertion should be avoided. The concern is less for other STIs.

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	1
Pregnancy.....	1
Age	1
Parity.....	1
Postpartum	1
Post-abortion.....	1
Past ectopic pregnancy.....	1
History of pelvic surgery.....	1
Smoking.....	2
Obesity.....	2
Blood pressure measurement unavailable	2
CARDIOVASCULAR DISEASE.....	2
Multiple risk factors for arterial cardiovascular disease.....	2
Hypertension.....	2
History of high blood pressure during pregnancy.....	3
Deep venous thrombosis (DVT)/pulmonary embolism (PE).....	3
Known thrombogenic mutations.....	3
Superficial venous thrombosis	3
Current and history of ischaemic heart disease	3
Stroke	3
Known hyperlipidaemias	4
Valvular heart disease	4
NEUROLOGIC CONDITIONS.....	4
Headaches.....	4
Epilepsy	4
DEPRESSIVE DISORDERS	4
Depressive disorders	4
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS.....	4
Unexplained vaginal bleeding	4
Endometriosis	4
Benign ovarian tumours	5
Severe dysmenorrhoea.....	5
Trophoblast disease.....	5
Cervical ectropion	5
Cervical intraepithelial neoplasia (CIN)	5
Cervical cancer	5
Breast disease	5
Endometrial cancer	5
Ovarian cancer.....	5
Uterine fibroids.....	5
Anatomical abnormalities.....	6
Pelvic inflammatory disease (PID)	6
STIs	6
HIV/AIDS	6
High risk of HIV	6
HIV-infected	6
AIDS	6

OTHER INFECTIONS	6
Schistosomiasis	6
Tuberculosis.....	6
Malaria	7
History of toxic shock syndrome	7
Urinary tract infection	7
ENDOCRINE CONDITIONS.....	7
Diabetes.....	7
Thyroid disorders	7
GASTROINTESTINAL CONDITIONS.....	7
Gall-bladder disease	7
History of cholestasis	7
Viral hepatitis	8
Cirrhosis.....	8
Liver tumours	8
ANAEMIAS.....	8
Thalassaemia.....	8
Sickle cell disease.....	8
Iron-deficiency anaemia.....	8
DRUG INTERACTIONS	8
Drugs which affect liver enzymes.....	8
Antibiotics	8
Antiretroviral therapy.....	8
Allergy to latex	9
Additional comments	10
References for barrier methods	11

BARRIER METHODS

C = Male latex condoms, male polyurethane condoms, female condoms

S = Spermicide (film, tablets, foam, gel)

D = Diaphragm (with spermicide), cervical cap

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY				
PREGNANCY	NA	NA	NA	Clarification: None of these methods are relevant for contraception during known pregnancy. However, for women who continue to be at risk of STI/HIV during pregnancy, the correct and consistent use of condoms is recommended.
AGE				
a) Menarche to < 40 years	1	1	1	
b) ≥ 40 years	1	1	1	
PARITY				
a) Nulliparous	1	1	1	
b) Parous	1	1	2	Clarification: There is a higher risk of cervical cap failure in parous women than in nulliparous women.
POSTPARTUM				
a) < 6 weeks postpartum	1	1	NA	Clarification: Diaphragm and cap are unsuitable until uterine involution is complete.
b) ≥ 6 weeks postpartum	1	1	1	
POST-ABORTION				
a) First trimester	1	1	1	
b) Second trimester	1	1	1	Clarification: Diaphragm and cap are unsuitable until 6 weeks after second-trimester abortion.
c) Immediate post-septic abortion	1	1	1	
PAST ECTOPIC PREGNANCY	1	1	1	
HISTORY OF PELVIC SURGERY	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
SMOKING				
a) Age < 35	1	1	1	
b) Age > 35				
(i) <15 cigarettes/day	1	1	1	
(ii) ≥15 cigarettes/day	1	1	1	
OBESITY* ≥ 30 kg/m ² body mass index (BMI)	1	1	1	
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA	NA	NA	Clarification: While a blood pressure measurement may be appropriate for good preventative health care, it is not required for safe and effective barrier method use. Women should not be denied use of barrier methods simply because their blood pressure cannot be measured.
CARDIOVASCULAR DISEASE				
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	1	1	1	
HYPERTENSION				
a) History of hypertension where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	1	1	1	
b) Adequately controlled hypertension, where blood pressure CAN be evaluated	1	1	1	
c) Elevated blood pressure levels (properly taken measurements)	1	1	1	
(i) systolic 140-159 or diastolic 90-99	1	1	1	
(ii) systolic ≥160 or diastolic ≥100	1	1	1	
d) Vascular disease	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	1	1	1	
DEEP VENOUS THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)				
a) History of DVT/PE	1	1	1	
b) Current DVT/PE	1	1	1	
c) Family history of DVT/PE (first degree relatives)	1	1	1	
d) Major surgery				
(i) with prolonged immobilization	1	1	1	
(ii) without prolonged immobilization	1	1	1	
e) Minor surgery without immobilization	1	1	1	
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	1	1	1	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
SUPERFICIAL VENOUS THROMBOSIS				
a) Varicose veins	1	1	1	
b) Superficial thrombophlebitis	1	1	1	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE	1	1	1	
STROKE (history of cerebrovascular accident)	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
KNOWN HYPERLIPIDAEMIAS	1	1	1	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
VALVULAR HEART DISEASE*				
a) Uncomplicated	1	1	1	
b) Complicated (pulmonary hypertension, atrial fibrillation, history of subacute bacterial endocarditis)	1	1	2	
NEUROLOGIC CONDITIONS				
HEADACHES				
a) Non-migrainous (mild or severe)	1	1	1	
b) Migraine				
(i) without aura				
Age < 35	1	1	1	
Age ≥ 35	1	1	1	
(ii) With aura, at any age	1	1	1	
EPILEPSY	1	1	1	
DEPRESSIVE DISORDERS				
DEPRESSIVE DISORDERS	1	1	1	
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS				
UNEXPLAINED VAGINAL BLEEDING (suspicious for serious condition)				
Before evaluation	1	1	1	Clarification: If pregnancy or an underlying pathological condition (such as pelvic malignancy) is suspected, it must be evaluated and the category adjusted after evaluation.
ENDOMETRIOSIS	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
BENIGN OVARIAN TUMOURS (including cysts)	1	1	1	
SEVERE DYSMENORRHOEA	1	1	1	
TROPHOBLAST DISEASE				
a) Benign gestational trophoblastic disease	1	1	1	
b) Malignant gestational trophoblastic disease	1	1	1	
CERVICAL ECTROPION	1	1	1	
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	1	1	1	Clarification: The cap should not be used. There is no restriction for diaphragm use.
CERVICAL CANCER* (awaiting treatment)	1	2	1	Clarification: The cap should not be used. There is no restriction for diaphragm use.
BREAST DISEASE				
a) Undiagnosed mass	1	1	1	
b) Benign breast disease	1	1	1	
c) Family history of cancer	1	1	1	
d) Breast cancer				
(i) current	1	1	1	
(ii) past and no evidence of current disease for 5 years	1	1	1	
ENDOMETRIAL CANCER	1	1	1	
OVARIAN CANCER	1	1	1	
UTERINE FIBROIDS				
a) Without distortion of the uterine cavity	1	1	1	
b) With distortion of the uterine cavity	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
ANATOMICAL ABNORMALITIES	1	1	NA	Clarification: The diaphragm cannot be used in certain cases of prolapse. Cap use is not appropriate for a client with a markedly distorted cervical anatomy.
PELVIC INFLAMMATORY DISEASE (PID)				
a) Past PID (assuming no current risk factors of STIs)				
(i) with subsequent pregnancy	1	1	1	
(ii) without subsequent pregnancy	1	1	1	
b) PID current	1	1	1	
STIS				
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	1	1	
b) Other STIs (excluding HIV and hepatitis)	1	1	1	
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	1	1	
d) Increased risk of STIs	1	1	1	
HIV/AIDS				
HIGH RISK OF HIV*	1	4	3	Evidence: Repeated and high-dose use of the spermicide nonoxynol-9 was associated with increased risk of genital lesions, which may increase the risk of acquiring HIV infection. ¹
HIV-INFECTED	1	4	3	
AIDS	1	4	3	
OTHER INFECTIONS				
SCHISTOSOMIASIS				
a) Uncomplicated	1	1	1	
b) Fibrosis of liver	1	1	1	
TUBERCULOSIS				

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
a) Non-pelvic	1	1	1	
b) Known pelvic	1	1	1	
MALARIA	1	1	1	
HISTORY OF TOXIC SHOCK SYNDROME*	1	1	3	
URINARY TRACT INFECTION*	1	1	2	
ENDOCRINE CONDITIONS				
DIABETES				
a) History of gestational disease	1	1	1	
b) Non-vascular disease				
(i) non-insulin dependent	1	1	1	
(ii) insulin dependent	1	1	1	
c) Nephropathy/retinopathy/neuropathy	1	1	1	
d) Other vascular disease or diabetes of > 20 years' duration	1	1	1	
THYROID DISORDERS				
a) Simple goitre	1	1	1	
b) Hyperthyroid	1	1	1	
c) Hypothyroid	1	1	1	
GASTROINTESTINAL CONDITIONS				
GALL-BLADDER DISEASE				
a) Symptomatic				
(i) treated by cholecystectomy	1	1	1	
(ii) medically treated	1	1	1	
(iii) current	1	1	1	
b) Asymptomatic	1	1	1	
HISTORY OF CHOLESTASIS				

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
a) Pregnancy-related	1	1	1	
b) Past COC-related	1	1	1	
VIRAL HEPATITIS				
a) Active	1	1	1	
b) Carrier	1	1	1	
CIRRHOSIS				
a) Mild (compensated)	1	1	1	
b) Severe (decompensated)	1	1	1	
LIVER TUMOURS				
a) Benign (adenoma)	1	1	1	
b) Malignant (hepatoma)	1	1	1	
ANAEMIAS				
THALASSAEMIA	1	1	1	
SICKLE CELL DISEASE	1	1	1	
IRON-DEFICIENCY ANAEMIA	1	1	1	
DRUG INTERACTIONS				
DRUGS WHICH AFFECT LIVER ENZYMES				
a) Rifampicin	1	1	1	
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	1	1	1	
ANTIBIOTICS (excluding rifampicin)				
a) Griseofulvin	1	1	1	
b) Other antibiotics	1	1	1	
ANTIRETROVIRAL THERAPY	1	1	1	

* See also additional comments at end of table

BARRIER METHODS	If there is risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.			
Women with conditions which make pregnancy an unacceptable risk should be advised that barrier methods for pregnancy prevention may not be appropriate for those who cannot use them consistently and correctly because of their relatively-higher typical-use failure rates.				
CONDITION	CATEGORY			CLARIFICATIONS/EVIDENCE
	C	S	D	
ALLERGY TO LATEX	3	1	3	Clarification: This does not apply to plastic condoms/diaphragms.

* See also additional comments at end of table

Additional comments

OBESITY

Severe obesity may make diaphragm and cap placement difficult.

VALVULAR HEART DISEASE

Risk of urinary tract infection with the diaphragm may increase risk in a client with sub-acute bacterial endocarditis.

CERVICAL CANCER (awaiting treatment)

Repeated and high-dose use of nonoxynol-9 can cause vaginal and cervical irritation or abrasions.

HIGH RISK OF HIV

Category 3 for diaphragm use is assigned due to concerns about the spermicide, not the diaphragm.

HISTORY OF TOXIC SHOCK SYNDROME

Toxic shock syndrome has been reported in association with contraceptive sponge and diaphragm use.

URINARY TRACT INFECTION

There is a potential increase of urinary tract infection with diaphragms and spermicides.

References for barrier methods

1. Wilkinson D et al. Nonoxynol-9 for preventing vaginal acquisition of HIV infection by women from men. Cochrane Database of Systematic Reviews, 2002, 4:CD003936.

FERTILITY AWARENESS-BASED METHODS

Fertility awareness-based (FAB) methods of family planning involve identification of the fertile days of the menstrual cycle, whether by observing fertility signs such as cervical secretions and basal body temperature, or by monitoring cycle days. FAB methods can be used in combination with abstinence or barrier methods during the fertile time. If barrier methods are used, refer to the section on barrier methods (BARR).

There are no medical conditions which become worse because of use of FAB methods. In general, these methods can be provided without concern for health effects to people who choose them. However, there are a number of conditions that make their use more complex. The existence of these conditions suggests that (1) use of these methods should be delayed until the condition is corrected or resolved or (2) they will require special counselling, and a more highly trained provider is generally necessary to ensure correct use.

Definitions

SYM	Symptoms-based methods	FAB methods based on observation of fertility signs (e.g., cervical secretions, basal body temperature) such as the Cervical Mucus Method, the Symptothermal Method, and the Two Day Method.
CAL	Calendar-based methods	FAB methods based on calendar calculations such as the Calendar Rhythm Method and the Standard Days Method.
A	Accept	There is no medical reason to deny the particular FAB method to a woman in this circumstance.
C	Caution	The method is normally provided in a routine setting, but with extra preparation and precautions. For FAB methods, this usually means that special counselling may be needed to ensure correct use of the method by a woman in this circumstance.
D	Delay	Use of this method should be delayed until the condition is evaluated or corrected. Alternative temporary methods of contraception should be offered.
NA	Not applicable	

FERTILITY AWARENESS-BASED METHODS	Fertility awareness-based methods do not protect against STI/HIV. If there is a risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.		
Women with conditions which make pregnancy an unacceptable risk should be advised that fertility awareness-based methods may not be appropriate for them because of their relatively-higher typical-use failure rates.			
CONDITION	CATEGORY		CLARIFICATIONS/EVIDENCE
	SYM	CAL	
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY			
PREGNANCY	NA		Comments: FAB methods are not relevant during pregnancy.
LIFE STAGE			
a) Post-menarche	C	C	Clarification: Menstrual irregularities are common in post-menarche and peri-menopause and may complicate the use of FAB methods.
b) Peri-menopause	C	C	
BREASTFEEDING*			
a) < 6 weeks postpartum	D	D	
b) ≥ 6 weeks	C	D	
c) After menses begin	C	C	
POSTPARTUM* (in non-breastfeeding women)			
a) < 4 weeks	D	D	
b) ≥ 4 weeks	A	D	
POST-ABORTION*	C	D	
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS			
IRREGULAR VAGINAL BLEEDING*	D	D	
VAGINAL DISCHARGE*	D	A	
OTHER			
USE OF DRUGS WHICH AFFECT CYCLE REGULARITY, HORMONES AND/OR FERTILITY SIGNS*	C/D	C/D	
DISEASES WHICH ELEVATE BODY TEMPERATURE*			
a) Chronic diseases	C	A	
b) Acute diseases	D	A	

* See also additional comments at end of table

Additional comments

BREASTFEEDING

FAB methods during breastfeeding may be less effective than when not breastfeeding.

< 6 weeks postpartum: Women who are primarily breastfeeding and are amenorrhoeic are unlikely to have sufficient ovarian function to produce detectable fertility signs and hormonal changes during the first 6 months postpartum. However, the likelihood of resumption of fertility increases with time postpartum and with substitution of breast milk by other foods.

After menses begin: When the woman notices fertility signs (particularly cervical secretions), she can use a symptoms-based method. When she has had 3 postpartum menses, she can use a calendar-based method. Prior to that time, a barrier method should be offered if the woman plans to use a FAB method later.

POSTPARTUM

< 4 weeks: Non-breastfeeding women are not likely to have sufficient ovarian function to either require a FAB method or to have detectable fertility signs or hormonal changes prior to 4 weeks postpartum. Although the risk of pregnancy is low, a method appropriate for the postpartum period should be offered.

≥ 4 weeks: Non-breastfeeding women are likely to have sufficient ovarian function to produce detectable fertility signs and/or hormonal changes at this time; likelihood increases rapidly with time postpartum. Women can use calendar-based methods as soon as they have completed 3 postpartum menses. Methods appropriate for the postpartum period should be offered prior to that time.

POST-ABORTION

Post-abortion women are likely to have sufficient ovarian function to produce detectable fertility signs and/or hormonal changes; likelihood increases with time post-abortion. Women can start using calendar based methods after they have had at least one post-abortion menses (e.g. women who before this pregnancy had most cycles between 26 and 32 days can use the Standard Days Method then). Methods appropriate for the post-abortion period should be offered prior to that time.

IRREGULAR VAGINAL BLEEDING

Presence of this condition makes FAB methods unreliable. Therefore, barrier methods should be recommended until the bleeding pattern is compatible with proper method use. The condition should be evaluated and treated as necessary.

VAGINAL DISCHARGE

Because vaginal discharge makes recognition of cervical secretions difficult, the condition should be evaluated and treated if needed prior to providing methods based on cervical secretions.

USE OF DRUGS WHICH AFFECT CYCLE REGULARITY, HORMONES AND/OR FERTILITY SIGNS

Use of certain mood-altering drugs such as lithium, tricyclic antidepressants, and anti-anxiety therapies, as well as certain antibiotics and anti-inflammatory drugs, may alter cycle regularity or affect fertility signs. The condition should be carefully evaluated and a barrier method offered until the degree of effect has been determined or the drug is no longer being used.

DISEASES WHICH ELEVATE BODY TEMPERATURE

Elevated temperature levels may make basal body temperature difficult to interpret, but there is no effect on cervical secretions. Thus the use of a method that relies on temperature should be delayed until the acute disease abates. Temperature-based methods are not appropriate for women with chronically-elevated temperatures. In addition, some chronic diseases interfere with cycle regularity, making calendar-based methods difficult to interpret.

LACTATIONAL AMENORRHOEA METHOD

The lactational amenorrhoea method does not protect against STI/HIV. If there is a risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.

Women with conditions which make pregnancy an unacceptable risk should be advised that the lactational amenorrhoea method may not be appropriate for them because of its relatively-higher typical-use failure rates.

The Bellagio Consensus provided the scientific basis for defining the conditions under which breastfeeding can be used safely and effectively for birth-spacing purposes, and programmatic guidelines were developed for the use of lactational amenorrhoea in family planning. These guidelines include the following three criteria, all of which must be met to ensure adequate protection from an unplanned pregnancy: **1) Amenorrhoea; 2) Fully or nearly fully breastfeeding; and 3) Less than six months postpartum.**

The main indications for breastfeeding remain the need to provide an ideal food for the infant and to protect it against disease. There are no medical conditions in which the use of lactational amenorrhoea is restricted and there is no documented evidence of its negative impact on maternal health. However, certain conditions or obstacles which affect breastfeeding may also affect the duration of amenorrhoea, making this a less useful choice for family planning purposes. These include:

HIV infection

Breastfeeding should be promoted, protected, and supported in all populations, for all women who are HIV-negative or of unknown HIV status. When replacement feeding is acceptable, feasible, affordable, sustainable and safe, avoidance of all breastfeeding by HIV-infected mothers is recommended. Otherwise, exclusive breastfeeding is recommended during the first months of life, and should then be discontinued as soon as it is feasible. Women who are HIV-positive should receive counselling that includes information about both the risks and benefits of various infant feeding options based on local assessments, guidance in selecting the most suitable option for their situation, and be supported in their choice. They should also have access to follow-up care and support, including family planning and nutritional support.

Medication used during breastfeeding

In order to protect infant health, breastfeeding is not recommended for women using such drugs as: anti-metabolites, bromocriptine, certain anticoagulants, corticosteroids (high doses), cyclosporin, ergotamine, lithium, mood-altering drugs, radioactive drugs, and reserpine.

Conditions affecting the newborn

Congenital deformities of the mouth, jaw or palate; newborns who are small-for-date or premature and needing intensive neonatal care; and certain metabolic disorders of the infant all can make breastfeeding difficult.

COITUS INTERRUPTUS

Coitus interruptus does not protect against STI/HIV. If there is a risk of STI/HIV (including during pregnancy or postpartum), the correct and consistent use of condoms should be recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.

Women with conditions that make pregnancy an unacceptable risk should be advised that coitus interruptus may not be appropriate for them because of its relatively-higher typical-use failure rates.

Coitus interruptus (CI), also known as withdrawal, is a traditional family planning method in which the man completely removes his penis from the vagina, and away from the external genitalia of the female partner, before he ejaculates. CI prevents sperm from entering the woman's vagina, thereby preventing contact between spermatozoa and the ovum.

This method may be appropriate for couples:

- who are highly motivated and able to use this method effectively;
- with religious or philosophical reasons for not using other methods of contraception;
- who need contraception immediately and have entered into a sexual act without alternative methods available;
- who need a temporary method while awaiting the start of another method;
- who have intercourse infrequently.

Some benefits of CI are that the method, if used correctly, does not affect breastfeeding and is always available for primary use or use as a back-up method. In addition, CI involves no economic cost or use of chemicals. There are no health risks associated directly with CI. Men and women who are at high risk of STI/HIV infection should use a condom with each act of intercourse.

CI is unforgiving of incorrect use, and its effectiveness depends on the willingness and ability of the couple to use withdrawal with every act of intercourse.

A. Female surgical sterilization	2
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	2
Pregnancy.....	2
Young age.....	2
Parity.....	2
Breastfeeding.....	2
Postpartum	2
Post-abortion.....	3
Past ectopic pregnancy.....	3
Smoking.....	3
Obesity.....	3
CARDIOVASCULAR DISEASE.....	4
Multiple risk factors for arterial cardiovascular disease*.....	4
Hypertension.....	4
History of high blood pressure during pregnancy.....	4
Deep venous thrombosis (DVT)/pulmonary embolism (PE).....	4
Known thrombogenic mutations.....	5
Superficial venous thrombosis.....	5
Current and history of ischaemic heart disease*.....	5
Stroke.....	5
Known hyperlipidaemias.....	5
Valvular heart disease.....	5
NEUROLOGIC CONDITIONS.....	6
Headaches.....	6
Epilepsy.....	6
DEPRESSIVE DISORDERS	6
Depressive disorders.....	6
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS.....	6
Vaginal bleeding patterns.....	6
Unexplained vaginal bleeding.....	6
Endometriosis.....	6
Benign ovarian tumours.....	6
Severe dysmenorrhoea.....	6
Trophoblast disease.....	7
Cervical ectropion.....	7
Cervical intraepithelial neoplasia (CIN).....	7
Cervical cancer.....	7
Breast disease.....	7
Endometrial cancer.....	7
Ovarian cancer.....	7
Uterine fibroids.....	7
Pelvic inflammatory disease (PID).....	7
STIs.....	8
HIV/AIDS	8
High risk of HIV.....	8
HIV-infected.....	8
AIDS.....	8
OTHER INFECTIONS	8
Schistosomiasis.....	8
Tuberculosis.....	8
Malaria.....	8

ENDOCRINE CONDITIONS.....	8
Diabetes.....	8
Thyroid disorders	9
GASTROINTESTINAL CONDITIONS.....	9
Gall-bladder disease	9
History of cholestasis	9
Viral hepatitis	9
Cirrhosis.....	9
Liver tumours	9
ANAEMIAS.....	9
Thalassaemia.....	9
Sickle-cell disease	9
Iron-deficiency anaemia.....	10
OTHER CONDITIONS RELEVANT ONLY FOR FEMALE SURGICAL STERILIZATION.....	10
Local infection	10
Coagulation disorders	10
Respiratory diseases	10
Systemic infection or gastroenteritis	10
Fixed uterus due to previous surgery or infection	10
Abdominal wall or umbilical hernia.....	10
Diaphragmatic hernia.....	10
Kidney disease.....	10
Severe nutritional deficiencies	10
Previous abdominal or pelvic surgery	10
Sterilization concurrent with abdominal surgery.....	11
Sterilization concurrent with Caesarean section	11
B. Male surgical sterilization.....	13
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY.....	13
Young age.....	13
DEPRESSIVE DISORDERS	13
Depressive disorders	13
HIV/AIDS	13
High risk of HIV	13
HIV-infected	13
AIDS	13
ENDOCRINE CONDITIONS.....	13
Diabetes.....	13
ANAEMIAS.....	13
Sickle-cell disease	13
OTHER CONDITIONS RELEVANT ONLY FOR MALE SURGICAL STERILIZATION	13
Local infections	13
Coagulation disorders	13
Previous scrotal injury.....	13
Systemic infection or gastroenteritis	13
Large varicocele.....	14
Large hydrocele*.....	14
Filariasis; elephantiasis	14
Intrascrotal mass.....	14
Cryptorchidism	14
Inguinal hernia	14
Additional comments	15
References for sterilization procedures	17

SURGICAL STERILIZATION PROCEDURES

Given that sterilization is a surgical procedure that is intended to be permanent, special care must be taken to assure that every client makes a voluntary informed choice of the method. Particular attention must be given in the case of young people, nulliparous women, men who have not yet been fathers, and clients with mental health problems, including depressive conditions. All clients should be carefully counselled about the intended permanence of sterilization and the availability of alternative, long-term, highly effective methods. This is of extra concern for young people. The national laws and existing norms for the delivery of sterilization procedures must be considered in the decision process.

Transcervical methods of female sterilization are not addressed in these recommendations.

There is no medical condition that would absolutely restrict a person's eligibility for sterilization, although some conditions and circumstances will require that certain precautions are taken, including those where the recommendation is C (Caution), D (Delay), or S (Special). For some of these conditions and circumstances, the theoretical or proven risks may outweigh the advantages of undergoing sterilization, particularly female sterilization. Where the risks of sterilization outweigh the benefits, long-term, highly effective contraceptive methods are a preferable alternative. Decisions in this regard will have to be made on an individual basis, considering the risks and benefits of sterilization versus the risks of pregnancy, and the availability and acceptability of highly effective, alternative methods.

The following classification of conditions into the four different categories is based on an in-depth review of the epidemiological and clinical evidence relevant to medical eligibility. Sterilization procedures should only be performed by well-trained providers in appropriate clinical settings using proper equipment and supplies. Appropriate service delivery guidelines, including infection prevention protocols, should be followed to maximize client safety.

DEFINITIONS

- | | | |
|----------|----------------|---|
| A | Accept | There is no medical reason to deny sterilization to a person with this condition. |
| C | Caution | The procedure is normally conducted in a routine setting, but with extra preparation and precautions. |
| D | Delay | The procedure is delayed until the condition is evaluated and/or corrected. Alternative temporary methods of contraception should be provided. |
| S | Special | The procedure should be undertaken in a setting with an experienced surgeon and staff, equipment needed to provide general anaesthesia, and other back-up medical support. For these conditions, the capacity to decide on the most appropriate procedure and anaesthesia regimen is also needed. Alternative temporary methods of contraception should be provided, if referral is required or there is otherwise any delay. |

A. Female surgical sterilization

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY		
PREGNANCY	D	
YOUNG AGE*	C	Clarification: Young women, like all women, should be counselled about the permanency of sterilization and the availability of alternative, long-term, highly effective methods. Evidence: Studies show that up to 20% of women sterilized at a young age later regret this decision, and that young age is one of the strongest predictors of regret (including request for reversal information and obtaining reversal) that can be identified before sterilization. ¹⁻¹⁹
PARITY*		
a) Nulliparous	A	
b) Parous	A	
BREASTFEEDING	A	
POSTPARTUM*		
a) < 7 days	A	
7 to < 42 days	D	
≥ 42 days	A	
b) Pre-eclampsia/ eclampsia		
(i) mild pre-eclampsia	A	
(ii) severe pre-eclampsia/ eclampsia	D	
c) Prolonged rupture of membranes: 24 hours or more	D	
d) Puerperal sepsis, intrapartum or puerperal fever	D	
e) Severe antepartum or postpartum haemorrhage	D	
f) Severe trauma to the genital tract: cervical or vaginal tear at time of delivery	D	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
POSTPARTUM (Cont'd)		
g) Uterine rupture or perforation	S	Clarification: If exploratory surgery or laparoscopy is conducted and the patient is stable, repair of the problem and tubal sterilization may be performed concurrently if no additional risk is involved.
POST-ABORTION*		
a) Uncomplicated	A	
b) Post-abortal sepsis or fever	D	
c) Severe post-abortal haemorrhage	D	
d) Severe trauma to the genital tract: cervical or vaginal tear at time of abortion	D	
e) Uterine perforation	S	Clarification: If exploratory surgery or laparoscopy is conducted, repair of the problem and tubal sterilization may be performed concurrently if no additional risk is involved.
f) Acute haematometra	D	
PAST ECTOPIC PREGNANCY	A	
SMOKING		
a) Age < 35 years	A	
b) Age ≥ 35 years		
(i) <15 cigarettes/day	A	
(ii) ≥15 cigarettes/day	A	
OBESITY ≥ 30 kg/m ² body mass index (BMI)	C	Clarification: The procedure may be more difficult. There is an increased risk of wound infection and disruption. Obese women may have limited respiratory function and may be more likely to require general anaesthesia. Evidence: Women who were obese were more likely to have complications when undergoing sterilization. ²⁰⁻²³

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
CARDIOVASCULAR DISEASE		
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE* (such as older age, smoking, diabetes and hypertension)	S	
HYPERTENSION		
For all categories of hypertension, classifications are based on the assumption that no other risk factors for cardiovascular disease exist. When multiple risk factors do exist, risk of cardiovascular disease may increase substantially. A single reading of blood pressure level is not sufficient to classify a woman as hypertensive.		
a) Hypertension, adequately controlled	C	Clarification: Elevated blood pressure should be controlled before surgery. There are increased anaesthesia-related risks and an increased risk of cardiac arrhythmia with uncontrolled hypertension. Careful monitoring of blood pressure intraoperatively is particularly necessary in this situation.
b) Elevated blood pressure levels (properly taken measurements)		
(i) systolic 140-159 or diastolic 90-99	C	
(ii) systolic ≥ 160 or diastolic ≥ 100	S	
c) Vascular disease	S	
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	A	
DEEP VEIN THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)		Clarification: To reduce the risk of DVT/PE, early ambulation is recommended.
a) History of DVT/PE	A	
b) Current DVT/PE	D	
c) Family history of DVT/PE (first-degree relatives)	A	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
DEEP VEIN THROMBOSIS (DVT)/PULMONARY EMBOLISM (PE) (Cont'd)		
d) Major surgery (i) with prolonged immobilization (ii) without prolonged immobilization e) Minor surgery without immobilization	D A A	
KNOWN THROMBOGENIC MUTATIONS (e.g., Factor V Leiden; Prothrombin mutation; Protein S, Protein C, and Antithrombin deficiencies)	A	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
SUPERFICIAL VEIN THROMBOSIS a) Varicose veins b) Superficial thrombophlebitis	A A	
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE* a) Current ischaemic heart disease b) History of ischaemic heart disease	D C	
STROKE (history of cerebrovascular accident)	C	
KNOWN HYPERLIPIDAEMIAS	A	Clarification: Routine screening is not appropriate because of the rarity of the conditions and the high cost of screening.
VALVULAR HEART DISEASE a) Uncomplicated b) Complicated (pulmonary hypertension, atrial fibrillation, history of subacute bacterial endocarditis)	C S	Clarification: The woman requires prophylactic antibiotics. Clarification: The woman is at high risk for complications associated with anaesthesia and surgery. If the woman has atrial fibrillation that has not been successfully managed or current subacute bacterial endocarditis, the procedure should be delayed.

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
NEUROLOGIC CONDITIONS		
HEADACHES		
a) Non-migrainous (mild or severe)	A	
b) Migraine		
(i) without aura		
<i>Age < 35</i>	A	
<i>Age ≥ 35</i>	A	
(ii) with aura (at any age)	A	
EPILEPSY	C	
DEPRESSIVE DISORDERS		
DEPRESSIVE DISORDERS	C	
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS		
VAGINAL BLEEDING PATTERNS		
a) Irregular pattern <i>without</i> heavy bleeding	A	
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	A	
UNEXPLAINED VAGINAL BLEEDING (suspicious for serious condition)		
Before evaluation	D	Clarification: The condition must be evaluated before the procedure is performed.
ENDOMETRIOSIS	S	
BENIGN OVARIAN TUMOURS (including cysts)	A	
SEVERE DYSMENORRHOEA	A	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
TROPHOBLAST DISEASE		
a) Benign gestational trophoblastic disease	A	
b) Malignant gestational trophoblastic disease	D	
CERVICAL ECTROPION	A	
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	A	
CERVICAL CANCER* (awaiting treatment)	D	
BREAST DISEASE		
a) Undiagnosed mass	A	
b) Benign breast disease	A	
c) Family history of cancer	A	
d) Breast cancer		
(i) current	C	
(ii) past and no evidence of current disease for 5 years	A	
ENDOMETRIAL CANCER*	D	
OVARIAN CANCER*	D	
UTERINE FIBROIDS*		
a) Without distortion of the uterine cavity	C	
b) With distortion of the uterine cavity	C	
PELVIC INFLAMMATORY DISEASE (PID)*		
a) Past PID (assuming no current risk factors for STIs)		Clarification: A careful pelvic examination must be performed to rule out recurrent or persistent infection and to determine the mobility of the uterus.
(i) with subsequent pregnancy	A	
(ii) without subsequent pregnancy	C	
b) PID - current	D	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
STIs*		
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	D	Clarification: If no symptoms persist following treatment, sterilization may be performed.
b) Other STIs (excluding HIV and hepatitis)	A	
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	A	
d) Increased risk of STIs	A	
HIV/AIDS		
HIGH RISK OF HIV	A	Clarification: No routine screening is needed. Appropriate infection prevention procedures, including universal precautions, must be carefully observed with all surgical procedures. The use of condoms is recommended following sterilization.
HIV-INFECTED	A	
AIDS	S	Clarification: The presence of an AIDS-related illness may require that the procedure be delayed.
OTHER INFECTIONS		
SCHISTOSOMIASIS		
a) Uncomplicated	A	Clarification: Liver function may need to be evaluated.
b) Fibrosis of liver	C	
TUBERCULOSIS		
a) Non-pelvic	A	
b) Known pelvic	S	
MALARIA	A	
ENDOCRINE CONDITIONS		
DIABETES*		
a) History of gestational disease	A	Clarification: If blood glucose is not well controlled, referral to a higher-level facility is recommended. Clarification: There is a possible decrease in healing and an increased risk of wound infection. Use of prophylactic antibiotics is recommended. Evidence: Diabetic women were more likely to have complications when undergoing sterilization. ²²
b) Non-vascular disease:		
(i) non-insulin dependent	C	
(ii) insulin dependent	C	
c) Nephropathy/retinopathy/neuropathy	S	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
DIABETES (Cont'd)		
d) Other vascular disease or diabetes of > 20 years' duration	S	
THYROID DISORDERS*		
a) Simple goitre	A	
b) Hyperthyroid	S	
c) Hypothyroid	C	
GASTROINTESTINAL CONDITIONS		
GALL-BLADDER DISEASE		
a) Symptomatic		
(i) treated by cholecystectomy	A	
(ii) medically treated	A	
(iii) current	D	
b) Asymptomatic	A	
HISTORY OF CHOLESTASIS		
a) Pregnancy-related	A	
b) Past COC-related	A	
VIRAL HEPATITIS*		
a) Active	D	Clarification: Appropriate infection prevention procedures, including universal precautions, must be carefully observed with all surgical procedures.
b) Carrier	A	
CIRRHOSIS		
a) Mild (compensated)	C	Clarification: Liver function and clotting might be altered. Liver function should be evaluated.
b) Severe (decompensated)	S	
LIVER TUMOURS		
a) Benign (adenoma)	C	Clarification: Liver function and clotting might be altered. Liver function should be evaluated.
b) Malignant (hepatoma)	C	
ANAEMIAS		
THALASSAEMIA	C	
SICKLE-CELL DISEASE*	C	

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
IRON-DEFICIENCY ANAEMIA a) Hb < 7g/dl b) Hb ≥ 7 to < 10g/dl	D C	Clarification: The underlying disease should be identified. Both preoperative Hb level and operative blood loss are important factors in women with anaemia. If peripheral perfusion is inadequate, this may decrease wound healing.
OTHER CONDITIONS RELEVANT ONLY FOR FEMALE SURGICAL STERILIZATION		
LOCAL INFECTION Abdominal skin infection	D	Clarification: There is an increased risk of postoperative infection.
COAGULATION DISORDERS*	S	
RESPIRATORY DISEASES* a) Acute (bronchitis, pneumonia) b) Chronic (i) asthma (ii) bronchitis (iii) emphysema (iv) lung infection	D S S S S	Clarification: The procedure should be delayed until the condition is corrected. There are increases in anaesthesia-related and other perioperative risks.
SYSTEMIC INFECTION OR GASTROENTERITIS*	D	
FIXED UTERUS DUE TO PREVIOUS SURGERY OR INFECTION*	S	
ABDOMINAL WALL OR UMBILICAL HERNIA	S	Clarification: Hernia repair and tubal sterilization should be performed concurrently, if possible.
DIAPHRAGMATIC HERNIA*	C	
KIDNEY DISEASE*	C	
SEVERE NUTRITIONAL DEFICIENCIES*	C	
PREVIOUS ABDOMINAL OR PELVIC SURGERY	C	Evidence: Women with previous abdominal or pelvic surgery were more likely to have complications when undergoing sterilization. ^{21, 22, 24-26}

* See also additional comments at end of table

FEMALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV (including during the postpartum period), the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
STERILIZATION CONCURRENT WITH ABDOMINAL SURGERY		
a) Elective	C	
b) Emergency (without previous counselling)	D	
c) Infectious condition	D	
STERILIZATION CONCURRENT WITH CAESAREAN SECTION*	A	

* See also additional comments at end of table

B. Male surgical sterilization

MALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV, the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY		
YOUNG AGE	C	Clarification: Young men, like all men, should be counselled about the permanency of sterilization and the availability of alternative, long-term, highly effective methods. Evidence: Men who underwent vasectomy at young ages were more likely to have the procedure reversed than those who underwent vasectomy at older ages. ¹⁸
DEPRESSIVE DISORDERS		
DEPRESSIVE DISORDERS	C	
HIV/AIDS		
HIGH RISK OF HIV HIV-INFECTED	A A	Clarification: No routine screening is needed. Appropriate infection prevention procedures, including universal precautions, must be carefully observed with all surgical procedures. The use of condoms is recommended following sterilization.
AIDS	S	Clarification: The presence of an AIDS-related illness may require a delay in the procedure.
ENDOCRINE CONDITIONS		
DIABETES*	C	
ANAEMIAS		
SICKLE-CELL DISEASE	A	
OTHER CONDITIONS RELEVANT ONLY FOR MALE SURGICAL STERILIZATION		
LOCAL INFECTIONS* a) scrotal skin infection b) active STI c) balanitis d) epididymitis or orchitis	D D D D	
COAGULATION DISORDERS*	S	
PREVIOUS SCROTAL INJURY	C	
SYSTEMIC INFECTION OR GASTROENTERITIS*	D	

* See also additional comments at end of table

MALE SURGICAL STERILIZATION	Sterilization does not protect against STI/HIV. If there is risk of STI/HIV, the correct and consistent use of condoms is recommended, either alone or with another contraceptive method. Male latex condoms are proven to protect against STI/HIV.	
CONDITION	CATEGORY	CLARIFICATIONS/EVIDENCE
LARGE VARICOCELE*	C	
LARGE HYDROCELE*	C	
FILARIASIS; ELEPHANTIASIS*	D	
INTRASCROTAL MASS*	D	
CRYPTORCHIDISM	C	Clarification: If cryptorchidism is bilateral, and fertility has been demonstrated, this will require extensive surgery to locate the vas, and this becomes category S. If the cryptorchidism is unilateral, and fertility has been demonstrated, vasectomy may be performed on the normal side and semen analysis performed, as per routine. If the man continues to have a persistent presence of sperm, more extensive surgery may be required to locate the other vas, and this becomes category S.
INGUINAL HERNIA*	S	

* See also additional comments at end of table

Additional comments

A. Female surgical sterilization

PARITY

Nulliparous: Nulliparous women, like all women, should be counselled about the permanency of sterilization and the availability of alternative, long-term, highly effective methods.

POSTPARTUM

< 7 days postpartum: Sterilization can be safely performed immediately postpartum.

7 to < 42 days: There is an increased risk of complications when the uterus has not fully involuted.

Pre-eclampsia/eclampsia: There are increased anaesthesia-related risks.

Prolonged rupture of membranes, 24 hours or more: There is an increased risk of postoperative infection.

Puerperal sepsis, intrapartum or puerperal fever: There is an increased risk of postoperative infection.

Severe antepartum or postpartum haemorrhage: The woman may be anaemic and unable to tolerate further blood loss.

Severe trauma to the genital tract: cervical or vaginal tear at time of delivery: There may have been significant blood loss and anaemia. The procedure may be more painful.

Uterine rupture or perforation: There may have been significant blood loss or damage to abdominal contents.

POST-ABORTION

Post-abortal sepsis or fever: There is an increased risk of postoperative infection.

Severe post-abortal haemorrhage: The woman may be anaemic and unable to tolerate further blood loss.

Severe trauma to the genital tract: cervical or vaginal tear at time of abortion: The woman may be anaemic and unable to tolerate further blood loss. The procedure may be more painful.

Uterine perforation: There may have been significant blood loss or damage to abdominal contents.

Acute haematometra: The woman may be anaemic and unable to tolerate further blood loss.

MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE

When multiple risk factors are present concurrently, the woman may be at high risk for complications associated with anaesthesia and surgery.

CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE

The woman is at high risk for complications associated with anaesthesia and surgery.

CERVICAL CANCER (awaiting treatment)

In general, the treatment renders a woman sterile.

ENDOMETRIAL CANCER

In general, the treatment renders a woman sterile.

OVARIAN CANCER

In general, the treatment renders a woman sterile.

UTERINE FIBROIDS

Depending on the size and location of the fibroids, it might be difficult to localize the tubes and mobilize the uterus.

PELVIC INFLAMMATORY DISEASE (PID)

PID can lead to an increased risk of post-sterilization infection or adhesions.

STIs

There is an increased risk of postoperative infection.

DIABETES

There is a risk of hypoglycaemia or ketoacidosis.

THYROID DISORDERS

The woman is at high risk for complications associated with anaesthesia and surgery.

VIRAL HEPATITIS

The woman is at high risk for complications associated with anaesthesia and surgery.

SICKLE-CELL DISEASE

There is an increased risk of pulmonary, cardiac or neurologic complications and possible increased risk of wound infection.

COAGULATION DISORDERS

Women with coagulation disorders are at increased risk of haematologic complications of surgery.

RESPIRATORY DISEASES

For laparoscopy, the woman may experience acute cardiorespiratory complications induced by pneumoperitoneum or the Trendelenburg position.

SYSTEMIC INFECTION OR GASTROENTERITIS

There are increased risks of postoperative infection, complications from dehydration, and anaesthesia-related complications.

FIXED UTERUS DUE TO PREVIOUS SURGERY OR INFECTION

Decreased mobility of the uterus, fallopian tubes and bowel may make laparoscopy and minilaparotomy difficult and increase the risk of complications.

DIAPHRAGMATIC HERNIA

For laparoscopy, the woman may experience acute cardiorespiratory complications induced by pneumoperitoneum or the Trendelenburg position.

KIDNEY DISEASE

Blood clotting may be impaired. There may be an increased risk of infection and hypovolemic shock. Condition may cause baseline anaemia, electrolyte disturbances and abnormalities in drug metabolism and excretion.

SEVERE NUTRITIONAL DEFICIENCIES

There may be an increased risk of wound infection and impaired healing.

STERILIZATION CONCURRENT WITH CAESAREAN SECTION

Concurrent sterilization does not increase the risk of complications in a surgically stable client.

B. Male surgical sterilization**COAGULATION DISORDERS**

Bleeding disorders lead to an increased risk of postoperative haematoma formation which, in turn, leads to an increased risk of infection.

DIABETES

Diabetics are more likely to get postoperative wound infections. If signs of infection appear, treatment with antibiotics needs to be given.

LOCAL INFECTIONS

There is an increased risk of postoperative infection.

SYSTEMIC INFECTION OR GASTROENTERITIS

There is an increased risk of postoperative infection.

LARGE VARICOCELE

The vas may be difficult or impossible to locate; a single procedure to repair varicocele and perform a vasectomy decreases the risk of complications.

LARGE HYDROCELE

The vas may be difficult or impossible to locate; a single procedure to repair hydrocele and perform a vasectomy decreases the risk of complications.

FILARIASIS; ELEPHANTIASIS

If elephantiasis involves the scrotum, it may be impossible to palpate the spermatic cord and testis.

INTRASCROTAL MASS

This may indicate an underlying disease.

INGUINAL HERNIA

Vasectomy can be performed concurrent with hernia repair.

References for sterilization procedures

1. Abraham S et al. The characteristics, perceptions and personalities of women seeking a reversal of their tubal sterilization. *Medical Journal of Australia*, 1986, 145:4-7.
2. Allyn DP et al. Presterilization counseling and women's regret about having been sterilized. *Journal of Reproductive Medicine*, 1986, 31:1027-32.
3. Boring CC, Rochat RW, Becerra J. Sterilization regret among Puerto Rican women. *Fertility & Sterility*, 1988, 49:973-81.
4. Clarkson SE, Gillett WR. Psychological aspects of female sterilisation--assessment of subsequent regret. *New Zealand Medical Journal*, 1985, 98:748-50.
5. Grubb GS et al. Regret after decision to have a tubal sterilization. *Fertility & Sterility*, 1985, 44:248-53.
6. Hardy E et al. Risk factors for tubal sterilization regret, detectable before surgery. *Contraception*, 1996, 54:159-62.
7. Henshaw SK, Singh S. Sterilization regret among U.S. couples. *Family Planning Perspectives*, 1986, 18:238-40.
8. Hillis SD et al. Poststerilization regret: findings from the United States Collaborative Review of Sterilization. *Obstetrics & Gynecology*, 1999, 93:889-95.
9. Jamieson DJ et al. A comparison of women's regret after vasectomy versus tubal sterilization. *Obstetrics & Gynecology*, 2002, 99:1073-9.
10. Kariminia A, Saunders DM, Chamberlain M. Risk factors for strong regret and subsequent IVF request after having tubal ligation. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 2002, 42:526-9.
11. Leader A et al. A comparison of definable traits in women requesting reversal of sterilization and women satisfied with sterilization. *American Journal of Obstetrics & Gynecology*, 1983, 145:198-202.
12. Loaiza E. Sterilization regret in the Dominican Republic: looking for quality-of-care issues. *Studies in Family Planning*, 1995, 26:39-48.
13. Marcil-Gratton N. Sterilization regret among women in metropolitan Montreal. *Family Planning Perspectives*, 1988, 20:222-7.
14. Platz-Christensen JJ et al. Evaluation of regret after tubal sterilization. *International Journal of Gynaecology & Obstetrics*, 1992, 38:223-6.
15. Ramsay IN, Russell SA. Who requests reversal of female sterilisation? A retrospective study from a Scottish unit. *Scottish Medical Journal*, 1991, 36:44-6.
16. Schmidt JE et al. Requesting information about and obtaining reversal after tubal sterilization: findings from the U.S. Collaborative Review of Sterilization. *Fertility & Sterility*, 2000, 74:892-8.
17. Thranov I et al. Regret among 547 Danish sterilized women. *Scandinavian Journal of Social Medicine*, 1988, 16:41-8.
18. Trussell J, Guilbert E, Hedley A. Sterilization failure, sterilization reversal, and pregnancy after sterilization reversal in Quebec. *Obstetrics & Gynecology*, 2003, 101:677-84.
19. Wilcox LS et al. Risk factors for regret after tubal sterilization: 5 years of follow-up in a prospective study. *Fertility & Sterility*, 1991, 55:927-33.
20. Chi I, Mumford SD, Laufe LE. Technical failures in tubal ring sterilization: incidence, perceived reasons, outcome, and risk factors. *American Journal of Obstetrics & Gynecology*, 1980, 138:307-12.
21. Chi I, Kennedy KI. Early readmission following elective laparoscopic sterilization: a brief analysis of a rare event. *American Journal of Obstetrics & Gynecology*, 1984, 148:322-7.
22. Jamieson DJ et al. Complications of interval laparoscopic tubal sterilization: findings from the United States Collaborative Review of Sterilization. *Obstetrics & Gynecology*, 2000, 96:997-1002.
23. White MK, Ory HW, Goldenberg LA. A case-control study of uterine perforations documented at laparoscopy. *American Journal of Obstetrics & Gynecology*, 1977, 129:623-5.
24. Baggish MS et al. Complications of laparoscopic sterilization. Comparison of 2 methods. *Obstetrics & Gynecology*, 1979, 54:54-9.
25. Chi I, Feldblum PJ, Balogh SA. Previous abdominal surgery as a risk factor in interval laparoscopic sterilization. *American Journal of Obstetrics & Gynecology*, 1983, 145:841-6.
26. Feldblum PJ et al. Technical failures in female sterilization using the tubal ring: a case-control analysis. *Contraception*, 1986, 34:505-12.

PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY	3
Pregnancy.....	3
Age	3
Parity.....	3
Breastfeeding.....	3
Postpartum	3
Post-abortion	3
Past ectopic pregnancy.....	3
History of pelvic surgery.....	4
Smoking.....	4
Obesity.....	4
Blood pressure measurement unavailable.....	4
CARDIOVASCULAR DISEASE	4
Multiple risk factors for arterial cardiovascular disease.....	4
Hypertension.....	4
History of high blood pressure during pregnancy.....	5
Deep venous thrombosis (DVT)/Pulmonary embolism (PE)	5
Known thrombogenic mutations.....	5
Superficial venous thrombosis	5
Current and history of ischaemic heart disease	5
Stroke	5
Known hyperlipidaemias	5
Valvular heart disease	6
NEUROLOGIC CONDITIONS	6
Headaches.....	6
Epilepsy	6
DEPRESSIVE DISORDERS	6
Depressive disorders	6
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS	6
Vaginal bleeding patterns	6
Unexplained vaginal bleeding	6
Endometriosis	6
Benign ovarian tumours	6
Severe dysmenorrhoea.....	7
Trophoblast disease.....	7
Cervical ectropion	7
Cervical intraepithelial neoplasia (CIN).....	7
Cervical cancer	7
Breast disease	7
Endometrial cancer	7
Ovarian cancer.....	7
Uterine fibroids.....	7
Anatomical abnormalities.....	7
Pelvic inflammatory disease (PID)	8
STIs	8
HIV/AIDS.....	8
High risk of HIV	8
HIV-infected	8
AIDS	8
OTHER INFECTIONS	8
Schistosomiasis	8
Tuberculosis	8
Malaria	8

ENDOCRINE CONDITIONS	9
Diabetes.....	9
Thyroid disorders	9
GASTROINTESTINAL CONDITIONS	9
Gall-bladder disease	9
History of cholestasis	9
Viral hepatitis	9
Cirrhosis.....	9
Liver tumours	10
ANAEMIAS	10
Thalassaemia	10
Sickle cell disease.....	10
Iron-deficiency anaemia.....	10
DRUG INTERACTIONS.....	10
Drugs which affect liver enzymes.....	10
Antibiotics	10
Antiretroviral therapy.....	10

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
PERSONAL CHARACTERISTICS AND REPRODUCTIVE HISTORY								
PREGNANCY	NA*	NA*	NA*	NA*	NA*	NA*	4*	4*
AGE	Menarche to <40=1 ≥40=2	Menarche to <40=1 ≥40=2	Menarche to <18=1 18-45=1 >45=1	Menarche to <18=2 18-45=1 >45=2	Menarche to <18=1 18-45=1 >45=1	Menarche to <20=2 ≥20=1	Menarche to <20=2 ≥20=1	Menarche to <20=2 ≥20=1
PARITY								
a) Nulliparous	1	1	1	1	1	1	2	2
b) Parous	1	1	1	1	1	1	1	1
BREASTFEEDING								
a) < 6 weeks postpartum	4	4	4	3*	3*	3*		
b) 6 weeks to < 6 months (primarily breastfeeding)	3	3	3	1	1	1		
c) ≥ 6 months postpartum	2	2	2	1	1	1		
POSTPARTUM (non-breastfeeding women)								
a) < 21 days	3	3	3	1	1	1		
b) ≥ 21 days	1	1	1	1	1	1		
POSTPARTUM (breastfeeding or non-breastfeeding women, including post-caesarean section)								
a) < 48 hours							2	3
b) ≥ 48 hours to <4 weeks							3	3
c) ≥ 4 weeks							1	1
d) Puerperal sepsis							4	4
POST-ABORTION								
a) First trimester	1*	1*	1*	1*	1*	1*	1*	1*
b) Second trimester	1	1	1	1	1	1	2	2
c) Immediate post-septic abortion	1	1	1	1	1	1	4	4
PAST ECTOPIC PREGNANCY	1	1	1	2	1	1	1	1

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
HISTORY OF PELVIC SURGERY (including caesarean section) (see also postpartum section)	1	1	1	1	1	1	1	1
SMOKING								
a) Age < 35	2*	2	2	1	1	1	1	1
b) Age ≥ 35								
(i) <15 cigarettes/day	3*	2	3	1	1	1	1	1
(ii) ≥15 cigarettes/day	4*	3	4	1	1	1	1	1
OBESITY ≥30 kg/m ² body mass index (BMI)	2	2	2	1	1	1	1	1
BLOOD PRESSURE MEASUREMENT UNAVAILABLE	NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*
CARDIOVASCULAR DISEASE								
MULTIPLE RISK FACTORS FOR ARTERIAL CARDIOVASCULAR DISEASE (such as older age, smoking, diabetes and hypertension)	3/4*	3/4*	3/4*	2*	3*	2*	1	2
HYPERTENSION								
a) History of hypertension where blood pressure CANNOT be evaluated (including hypertension during pregnancy)	3*	3*	3*	2*	2*	2*	1	2
b) Adequately controlled hypertension, where blood pressure CAN be evaluated	3*	3*	3*	1*	2*	1*	1	1
c) Elevated blood pressure levels (properly taken measurements)								
(i) systolic 140-159 or diastolic 90-99	3	3	3	1	2	1	1	1
(ii) systolic >160 or diastolic >100	4	4	4	2	3	2	1	2
d) Vascular disease	4	4	4	2	3	2	1	2

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
HISTORY OF HIGH BLOOD PRESSURE DURING PREGNANCY (where current blood pressure is measurable and normal)	2	2	2	1	1	1	1	1
DEEP VENOUS THROMBOSIS (DVT)/ PULMONARY EMBOLISM (PE)								
a) History of DVT/PE	4	4	4	2	2	2	1	2
b) Current DVT/PE	4	4	4	3	3	3	1	3
c) Family history (first-degree relatives)	2	2	2	1	1	1	1	1
d) Major surgery								
(i) with prolonged immobilization	4	4	4	2	2	2	1	2
(ii) without prolonged immobilization	2	2	2	1	1	1	1	1
e) Minor surgery without immobilization	1	1	1	1	1	1	1	1
KNOWN THROMBOGENIC MUTATIONS (e.g. Factor V Leiden; Prothrombin mutation; Protein S, Protein C and Antithrombin deficiencies)	4*	4*	4*	2*	2*	2*	1*	2*
SUPERFICIAL VENOUS THROMBOSIS								
a) Varicose veins	1	1	1	1	1	1	1	1
b) Superficial thrombophlebitis	2	2	2	1	1	1	1	1
CURRENT AND HISTORY OF ISCHAEMIC HEART DISEASE				I C		I C		I C
	4	4	4	2 3	3	2 3	1	2 3
STROKE (history of cerebrovascular accident)				I C		I C		
	4	4	4	2 3	3	2 3	1	2
KNOWN HYPERLIPIDAEMIAS (screening is NOT necessary for safe use of contraceptive methods)	2/3*	2/3*	2/3*	2*	2*	2*	1*	2*

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES										
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD		
I = Initiation, C = Continuation										
VALVULAR HEART DISEASE										
a) Uncomplicated	2	2	2	1	1	1	1	1		
b) Complicated (pulmonary hypertension, atrial fibrillation, history of subacute bacterial endocarditis)	4	4	4	1	1	1	2*	2*		
NEUROLOGIC CONDITIONS										
HEADACHES	I	C	I	C	I	C	I	C	I	C
a) Non-migrainous (mild or severe)	1*	2*	1*	2*	1*	2*	1*	1*	1*	1*
b) Migraine										
(i) without aura										
Age <35	2*	3*	2*	3*	2*	3*	1*	2*	2*	2*
Age ≥35	3*	4*	3*	4*	3*	4*	1*	2*	2*	2*
(ii) with aura (at any age)	4*	4*	4*	4*	4*	4*	2*	3*	2*	3*
EPILEPSY	1*	1*	1*	1*	1*	1*	1*	1	1	
DEPRESSIVE DISORDERS										
DEPRESSIVE DISORDERS	1*	1*	1*	1*	1*	1*	1*	1*	1*	
REPRODUCTIVE TRACT INFECTIONS AND DISORDERS										
VAGINAL BLEEDING PATTERNS									I	C
a) Irregular pattern <i>without</i> heavy bleeding	1	1	1	2	2	2	1		1	1
b) Heavy or prolonged bleeding (includes regular and irregular patterns)	1*	1*	1*	2*	2*	2*	2*		1*	2*
UNEXPLAINED VAGINAL BLEEDING (suspicious for serious condition)									I	C
Before evaluation	2*	2*	2*	2*	3*	3*			4*	2*
ENDOMETRIOSIS	1	1	1	1	1	1	2		1	
BENIGN OVARIAN TUMOURS (including cysts)	1	1	1	1	1	1	1		1	

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
SEVERE DYSMENORRHOEA	1	1	1	1	1	1	2	1
TROPHOBLAST DISEASE								
a) Benign gestational trophoblastic disease	1	1	1	1	1	1	3	3
b) Malignant gestational trophoblastic disease	1	1	1	1	1	1	4	4
CERVICAL ECTROPION	1	1	1	1	1	1	1	1
CERVICAL INTRAEPITHELIAL NEOPLASIA (CIN)	2	2	2	1	2	2	1	2
CERVICAL CANCER (awaiting treatment)							I C	I C
	2	2	2	1	2	2	4 2	4 2
BREAST DISEASE								
a) Undiagnosed mass	2*	2*	2*	2*	2*	2*	1	2
b) Benign breast disease	1	1	1	1	1	1	1	1
c) Family history of cancer	1	1	1	1	1	1	1	1
d) Cancer								
(i) current	4	4	4	4	4	4	1	4
(ii) past and no evidence of current disease for 5 years	3	3	3	3	3	3	1	3
ENDOMETRIAL CANCER							I C	I C
	1	1	1	1	1	1	4 2	4 2
OVARIAN CANCER							I C	I C
	1	1	1	1	1	1	3 2	3 2
UTERINE FIBROIDS								
a) Without distortion of the uterine cavity	1	1	1	1	1	1	1	1
b) With distortion of the uterine cavity	1	1	1	1	1	1	4	4
ANATOMICAL ABNORMALITIES								
a) That distort the uterine cavity							4	4
b) That do not distort the uterine cavity							2	2

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES											
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD		LNG-IUD		
I = Initiation, C = Continuation											
PELVIC INFLAMMATORY DISEASE (PID)											
a) Past PID (assuming no current risk factors of STIs)							I	C	I	C	
(i) with subsequent pregnancy	1	1	1	1	1	1	1	1	1	1	
ii) without subsequent pregnancy	1	1	1	1	1	1	2	2	2	2	
b) PID - current	1	1	1	1	1	1	4	2*	4	2*	
STIs							I	C	I	C	
a) Current purulent cervicitis or chlamydial infection or gonorrhoea	1	1	1	1	1	1	4	2*	4	2*	
b) Other STIs (excluding HIV and hepatitis)	1	1	1	1	1	1	2	2	2	2	
c) Vaginitis (including trichomonas vaginalis and bacterial vaginosis)	1	1	1	1	1	1	2	2	2	2	
d) Increased risk of STIs	1	1	1	1	1	1	2/3*	2	2/3*	2	
HIV/AIDS											
HIGH RISK OF HIV							I	C	I	C	
	1	1	1	1	1	1	2	2	2	2	
HIV-INFECTED	1	1	1	1	1	1	2	2	2	2	
AIDS	1*	1*	1*	1*	1*	1*	3	2*	3	2*	
Clinically well on ARV therapy	See ANTIRETROVIRAL THERAPY below							2	2	2	2
OTHER INFECTIONS											
SCHISTOSOMIASIS											
a) Uncomplicated	1	1	1	1	1	1	1		1		
b) Fibrosis of the liver	1	1	1	1	1	1	1		1		
TUBERCULOSIS							I	C	I	C	
a) Non-pelvic	1*	1*	1*	1*	1*	1*	1	1	1	1	
b) Known pelvic	1*	1*	1*	1	1	1	4	3	4	3	
MALARIA	1	1	1	1	1	1	1		1		

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
ENDOCRINE CONDITIONS								
DIABETES								
a) History of gestational disease	1	1	1	1	1	1	1	1
b) Non-vascular disease								
(i) non-insulin dependent	2	2	2	2	2	2	1	2
(ii) insulin dependent	2	2	2	2	2	2	1	2
c) Nephropathy/ retinopathy/ neuropathy	3/4*	3/4*	3/4*	2	3	2	1	2
d) Other vascular disease or diabetes of >20 years' duration	3/4*	3/4*	3/4*	2	3	2	1	2
THYROID DISORDERS								
a) Simple goitre	1	1	1	1	1	1	1	1
b) Hyperthyroid	1	1	1	1	1	1	1	1
c) Hypothyroid	1	1	1	1	1	1	1	1
GASTROINTESTINAL CONDITIONS								
GALL-BLADDER DISEASE								
a) Symptomatic								
(i) treated by cholecystectomy	2	2	2	2	2	2	1	2
(ii) medically treated	3	2	3	2	2	2	1	2
(iii) current	3	2	3	2	2	2	1	2
b) Asymptomatic	2	2	2	2	2	2	1	2
HISTORY OF CHOLESTASIS								
a) Pregnancy-related	2	2	2	1	1	1	1	1
b) Past COC-related	3	2	3	2	2	2	1	2
VIRAL HEPATITIS								
a) Active	4	3/4*	4*	3	3	3	1	3
c) Carrier	1	1	1	1	1	1	1	1
CIRRHOSIS								
a) Mild (compensated)	3	2	3	2	2	2	1	2
b) Severe (decompensated)	4	3	4	3	3	3	1	3

* Please consult the tables in the text for a clarification to this classification

SUMMARY TABLES								
CONDITION	COC	CIC	P/R	POP	DMPA NET-EN	LNG/ ETG Implants	Cu-IUD	LNG-IUD
I = Initiation, C = Continuation								
LIVER TUMOURS								
a) Benign (adenoma)	4	3	4	3	3	3	1	3
b) Malignant (hepatoma)	4	3/4	4	3	3	3	1	3
ANAEMIAS								
THALASSAEMIA	1	1	1	1	1	1	2	1
SICKLE CELL DISEASE	2	2	2	1	1	1	2	1
IRON-DEFICIENCY ANAEMIA	1	1	1	1	1	1	2	1
DRUG INTERACTIONS								
DRUGS WHICH AFFECT LIVER ENZYMES								
a) Rifampicin	3*	2*	3*	3*	2*	3*	1	1
b) Certain anticonvulsants (phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine)	3*	2*	3*	3*	2	3	1	1
ANTIBIOTICS (excluding rifampicin)								
a) Griseofulvin	2	1	2	2	1	2	1	1
b) Other antibiotics	1	1	1	1	1	1	1	1
ANTIRETROVIRAL THERAPY								
							I	C
	2*	2*	2*	2*	2*	2*	2/3*	2
							I	C
							2/3*	2

* Please consult the tables in the text for a clarification to this classification

Annex 1. COCs and antiretroviral therapies

Few data from small, mostly unpublished studies suggest that the pharmacokinetics of a single dose of COCs may be altered by various antiretroviral (ARV) therapies. However, no clinical outcome studies have been conducted and the clinical significance of such changes, especially when the COCs have not been allowed to reach steady-state, is unknown. The following table summarizes the evidence to date regarding the effects of ARVs on contraceptive steroid levels and the effects of hormonal contraceptives on ARV levels.

Table 1. Pharmacokinetic COC-ARV drug interactions.

ARV	Contraceptive steroid levels	ARV levels
Protease inhibitors		
Nelfinavir	↓	No data
Ritonavir	↓	No data
Lopinavir/ritonavir	↓	No data
Atazanavir	↑	No data
Amprenavir	↑	↓
Indinavir	↑	No data
Saquinavir	No data	No change
Non-nucleoside reverse transcriptase inhibitors		
Nevirapine	↓	No change
Efavirenz	↑	No change
Delavirdine	?↑	No data

References:

Ouellet D et al. Effect of ritonavir on the pharmacokinetics of ethinyl oestradiol in healthy female volunteers. *British Journal of Clinical Pharmacology*, 1998, 46(2):111-6.

Mildvan D et al. Pharmacokinetic interaction between nevirapine and ethinyl estradiol/norethindrone when administered concurrently to HIV-infected women. *Journal of Acquired Immune Deficiency Syndromes: JAIDS*, 2002, 29(5):471-7.

Tackett D et al. Atazanavir: a summary of two pharmacokinetic drug interaction studies in healthy subjects (abstract). Presented at the 10th Retrovirus Conference, Boston, MA, February 10-14, 2003. Available on line at <http://www.retroconference.org/2003/Abstract/Abstract.aspx?AbstractID=649>. (Accessed July 31, 2003).

Mayer K et al. Efficacy, effect of oral contraceptives, and adherence in HIV infected women receiving Fortovase (Saquinavir) soft gel capsule (SQV-SGC; FTV) thrice (TID) and twice (BID) daily regimens (abstract). Presented at the XIII International AIDS Conference, Durban, 2000. Available on line at <http://www.iac2000.org/abdetail.asp?ID=TuPeB3226>. (Accessed July 31, 2003).

Merck & Co., Inc. Indinavir prescribing information, 2002.

Abbot Laboratories. Ritonavir prescribing information, 2001.

Abbot Laboratories. Lopinavir/ritonavir prescribing information, 2003.

Agouron Pharmaceuticals, Inc. Nelfinavir prescribing information, 2003.

Glaxo Smith Kline. Amprenavir prescribing information, 2002.

Pharmacia & Upjohn Co. Delavirdine prescribing information, 2001.

DuPont Pharmaceuticals Co. Efavirenz prescribing information, 2001.

Bristol-Myers Squibb Company. Atazanavir prescribing information, 2003

Annex 2. List of participants

Expert Working Group Meeting to update the Medical Eligibility Criteria for Contraceptive Use Salle C, World Health Organization, Geneva, 21-24 October 2003

Dr Halida Akhter
Managing Director
Health Promotion Limited (HPL)
H-310, Rd-3 Baitul Aman
Housing Society, Shyamoli
Dhaka, 1207
Bangladesh

Dr Willard Cates
President
Family Health International
P. O. Box 13950
Research Triangle Park, NC 27709
United States of America

Dr Tsungai Chipato
Department of OB/GYN
University of Zimbabwe
P.O. Box A 178
Harare
Zimbabwe

Dr Maria del Carmen Cravioto
Department of Reproductive Biology
National Institute of Nutrition,
Salvador Zubiran, Vaso de Quiroga 15
Delegacion Tlalpan
C. P. 14000
Mexico, D.F.
Mexico

Dr Juan Diaz
Medical Advisor for Latin
America and the Caribbean
The Population Council
Cidade Universitaria
Caixa postal 6509
13084-970 Campinas SP
Sao Paulo
Brazil

Dr Soledad Diaz
Consultorio de Planificacion Familiar
Instituto Chileno de Medicina Reproductiva
José Victorino Latarria 29
Depto 101, Correo 22, Casilla 96
Santiago
Chile

Dr Lindsay Edouard
Senior Reproductive Health Officer
Technical and Policy Division
United Nations Population Fund
220 East 42nd Street
New York, NY 10017
United States of America

Dr Anna Glasier
Family Planning and Well Woman Services
18 Dean Terrace
Edinburgh EH4 1NL
United Kingdom

Dr John Guillebaud
White Leas Mead
14 Hidscope Road
Cumnor Hill
Oxford OX2 9JJ
United Kingdom

Dr Kerstin Hagenfeldt
Department of OB/GYN
Karolinska Institute
Box 140
S-171 76 Stockholm
Sweden

Dr Ezzeldin Othman Hassan
The National Egyptian Fertility Care
Foundation
2(A) Mahrouky Street Mohandessen
POB 147 Orman
Giza
Egypt

Dr Mihai Horga
Directorate of Family and Social Asst.
Ministry of Health & Family
1-3 Cristian Popisteanu St.
70109 Bucharest
Romania

Dr Douglas Huber
Management Sciences for Health
165 Allandale Road
Boston, MA 02130-3400
United States of America

Dr Carlos Huezo
International Planned Parenthood Federation
Regent's College, Inner Circle, Regent's Park
London NW1 4NS
United Kingdom

Dr Roy Jacobstein
EngenderHealth
440 Ninth Avenue
New York, NY 1001
United States of America

Dr Victoria Jennings
Institute for Reproductive Health
Georgetown University Medical Center
Georgetown Center, 6th Floor
2115 Wisconsin Avenue, NW
Washington, D.C. 20007
United States of America

Dr Liliana Kapetanovic-Milosevic
Pharmaco-Toxico-Clinical Evaluation Unit
Agence Française de Sécurité Sanitaire des
Produits de Santé
143/147, Boulevard Anatole France
93285 Saint Denis Cedex
France

Dr Robert Leke
WHO Centre for Research in Human
Reproduction
Faculty of Medicine & Biological Sciences
University of Yaoundé 1, BP 1935
Yaoundé
Cameroon

Dr Enriquito R. Lu
STARH/FP Program
Tifa Building, 5th Floor, Suite 503
JI Kuningan Barat 26
Jakarta 12710
Indonesia

Dr Trent MacKay
Contraception and Reproductive Health
Branch, NICHD, NIH
6100 Executive Boulevard, Suite 8B13
Rockville, MD 20892
United States of America

Dr Maurizio Macaluso
Women's Health and Fertility Branch
CDC
Koger Rhodes Building Mailstop K-34
4770 Buford Highway, NE
Atlanta, GA 30341-3717
United States of America

Dr Olav Meirik
Instituto Chileno de Medicina Reproductiva
Jose Ramon Gutierrez 295, Depto 3
Santiago
Chile

Dr Suneeta Mittal
Head, Department of OB/GYN
Director-in-Charge, WHO CCR in Human
Reproduction
All India Institute of Medical Sciences
Ansari Nagar
New Delhi, 110 029
India

Dr Helen Randera-Rees
Reproductive Health Unit
Department of OB/GYN
Baragwanath Hospital
PO Betsam 2013
Johannesburg
South Africa

Mr Ward Rinehart
John Hopkins University
Center for Communications Program
111 Market Place, Suite 310
Baltimore, MD 21202
United States of America

Dr Roberto Rivera
Family Health International
P. O. Box 13950
Research Triangle Park, NC 27709
United States of America

Dr Fred Sai
P. O. Box 9983
Kotoka Airport, 4404
Accra
Ghana

Dr Pramilla Senanayake
International Planned Parenthood Federation
Regent's College, Inner Circle, Regent's Park
London NW1 4NS
United Kingdom

Dr James Shelton
Office of Population
Bureau of Science and Technology
USAID
1300 Pennsylvania Avenue, G/PHN
Washington, D.C. 20523
United States of America

Dr Connie Smith
Westside Contraceptive Services
Westminster PCT
Raymede Clinic
St Charles Hospital
Exmoor Street
London W10 6DZ
United Kingdom

Mr Irving Sivin
Centre for Biomedical Research
The Population Council
One Dag Hammarskjold Plaza
New York, N.Y. 10017
United States of America

Dr Bulbul Sood
Program Management Specialist, CEDPA
50 M Shantipath, Gate No 3,
Niti Marg, Chanakyapuri,
New Delhi-110021
India

Dr Marcel Vekemans
IntraHealth International
1700 Airport Road, Suite 300 CB# 8100,
Chapel Hill, NC 27514
United States of America

Dr Edith Weisberg
Sydney Centre for Reproductive Health
Research
328-336 Liverpool Road
Ashfield, NSW 2131
Australia

Dr Wu Shangchun
National Research Institute for Family Planning
12 Da Hui Si (Hai Dian Qu)
Beijing 100081
People's Republic of China

TEMPORARY ADVISERS

Dr Kate Curtis
Division of Reproductive Health
Koger Rhodes Building
CDC Mailstop K-34
4770 Buford Highway, NE
Atlanta, GA 30341-3717
United States of America

Ms Anshu Mohllajee
Division of Reproductive Health
Koger Rhodes Building
CDC Mailstop K-34
4770 Buford Highway, NE
Atlanta, GA 30341-3717
United States of America

Dr Kavita Nanda
Family Health International
P. O. Box 13950
Research Triangle Park, NC 27709
United States of America

Dr Jennifer Smith
1 bis Montée de Garillan
69005 Lyon
France

SECRETARIAT

Ms Kathryn Church, RHR
Dr MaryLyn Gaffied, RHR
Ms Sarah Johnson, RHR
Ms Cath Hamill, RHR
Mrs Gloria Lamptey, RHR
Dr Herbert Peterson, RHR
Dr Paul Van Look, RHR
Dr Effy Vayena, RHR

OBSERVERS

Dr Vanessa E. Cullins
Vice President for Medical Affairs
Planned Parenthood Federation of America
434 West 33rd Street
New York, N.Y. 10001
United States of America

Dr Fatiha Terki
International Planned Parenthood Federation
Regent's College, Inner Circle, Regent's Park
London NW1 4NS
United Kingdom

